

## *ad*-nilpotent ideals containing a fixed number of simple root spaces

PAOLA CELLINI – P. MÖSENER FRAJRIA – PAOLO PAPI

ABSTRACT: *We give formulas for the number of ad-nilpotent ideals of a Borel subalgebra of a Lie algebra of type B or D containing a fixed number of root spaces attached to simple roots. This result solves positively a conjecture of Panyushev [12, 3.5] and affords a complete knowledge of the above statistics for any simple Lie algebra. We also study the restriction of the above statistics to the abelian ideals of a Borel subalgebra, obtaining uniform results for any simple Lie algebra*

### 1 – Introduction

Let  $\mathfrak{g}$  be a complex finite-dimensional simple Lie algebra. Fix a Borel subalgebra  $\mathfrak{b}$  of  $\mathfrak{g}$ , and let  $\mathfrak{n}$  be its nilradical. If  $\mathfrak{g}$  is of type  $X$ , denote by  $\mathcal{I}(X)$  the set of *ad*-nilpotent ideals  $\mathfrak{i}$ , *i.e.* the ideals of  $\mathfrak{b}$  which are contained in  $\mathfrak{n}$ . Let  $\Delta^+$ ,  $\Pi$  denote respectively the positive and simple systems of the root system  $\Delta$  of  $\mathfrak{g}$  corresponding to  $\mathfrak{b}$ . Then  $\mathfrak{i} \in \mathcal{I}(X)$  if and only if  $\mathfrak{i} = \bigoplus_{\alpha \in \Phi_{\mathfrak{i}}} \mathfrak{g}_{\alpha}$ , where  $\mathfrak{g}_{\alpha}$  is the root space attached to  $\alpha$  and  $\Phi_{\mathfrak{i}} \subseteq \Delta^+$  is a dual order ideal of  $\Delta^+$  (w.r.t. the usual order:  $\alpha < \beta$  if  $\beta - \alpha$  is a sum of positive roots). *ad*-nilpotent ideals have been intensively investigated in recent literature: see references in [12]. The first goal of this short paper is to solve positively conjecture 3.5 of [12]. This conjecture regards the following statistics on  $\mathcal{I}(X)$ :

$$P_X(j) = |\{\mathfrak{i} \in \mathcal{I}(X) : |\Pi \cap \Phi_{\mathfrak{i}}| = j\}|$$

( $0 \leq j \leq n$ ). The formulas expressing  $P_X(j)$  for the classical Lie algebras are given in the following theorem. The result in type  $A$  has been proved in [12, Theorem 3.4], together with the equality  $P_{B_n} = P_{C_n}$ . The formulas for types  $B, D$  are conjecture 3.5 of the same paper.

THEOREM 1.1. *For  $0 \leq j \leq n$  we have*

$$\begin{aligned}
 P_{A_n}(j) &= \frac{j+1}{n+1} \binom{2n-j}{n}, \\
 P_{B_n}(j) &= P_{C_n}(j) = \binom{2n-j-1}{n-1}, \\
 P_{D_n}(j) &= \begin{cases} \binom{2n-2}{n-2} + \binom{2n-3}{n-3} & \text{if } j = 0 \\ \binom{2n-2-j}{n-2} + \binom{2n-3-j}{n-2} & \text{if } 1 \leq j \leq n. \end{cases}
 \end{aligned}$$

We remark that the numerical values of  $P_X(j)$  in the exceptional cases are easily calculated from the knowledge of  $P_X(0)$  using the inclusion-exclusion principle: see [12, §3]. On the other hand, the number  $P_X(0)$  can be uniformly described: see Remark 2.1. The relevance of the statistics  $P_X$  is motivated by the following discussion. It is known [4] that the cardinality of  $\mathcal{I}$  is given by the generalized Catalan number  $\frac{1}{|W|} \prod_{i=1}^n (e_i + h + 1)$  (see Remark 2.1 for undefined notation) as well as that of *clusters*, certain subsets of  $\Delta^+ \cup -\Pi$  which play a major role in Zelevinsky's theory of cluster algebras [7]. Panyushev noticed that  $P_X(j)$  also counts the number of clusters having  $j$  elements in  $-\Pi$ . Looking for a conceptual explanation of the interplay between *ad*-nilpotent ideals and clusters is an interesting open problem. Theorem 1.1 is proved in the next section. The final section deals with a formula for the same statistics on the subset  $\mathcal{I}^{ab}$  of  $\mathcal{I}$  consisting of abelian ideals. The study of  $\mathcal{I}^{ab}$ , pursued by Kostant, started an intense research activity which was later extended by considering *ad*-nilpotent ideals. Abelian ideals turn out to appear in several contexts, ranging from the structure of the exterior algebra of  $\mathfrak{g}$  [9], to affine algebras [2] and to difficult problems in classical invariant theory [11]. The key fact originating this activity is the following celebrated enumerative result by Dale Peterson, which we are going to exploit:

$$(1.1) \quad |\mathcal{I}^{ab}| = 2^{rk(\mathfrak{g})}.$$

Regarding our statistics, we obtain the following “uniform” result. Let  $P, Q$  denote the weight and root lattice of  $\Delta$  and let  $z(\mathfrak{g}) = |P/Q|$  be the connection index.

THEOREM 1.2. *The number  $P_X^{ab}(j)$  of abelian ideals of  $\mathfrak{b}$  in a Lie algebra  $\mathfrak{g}$  of type  $X$  and rank  $n$  containing  $j$  simple roots is given by*

$$P_X^{ab}(j) = \begin{cases} 2^n - z(\mathfrak{g}) + 1 & \text{if } j = 0, \\ z(\mathfrak{g}) - 1 & \text{if } j = 1, \\ 0 & \text{if } j > 1. \end{cases}$$

**2 – Proof of Theorem 1.1**

Our approach to Panyushev’s conjecture is based on Shi’s encoding [13] of *ad*-nilpotent ideals for classical Lie algebras via (possibly shifted) shapes as formulated in [3] . More precisely, consider a staircase diagram  $T_X$  of shape  $(n, n - 1, \dots, 1)$  in type  $A_n$  (respectively a shifted staircase diagram of shape  $(2n - 1, 2n - 3, \dots, 1)$  for  $B_n$  and  $C_n$ , and of shape  $(2n - 2, 2n - 4, \dots, 2)$  for  $D_n$ ). Arrange in the diagram the positive roots of  $\Delta$  according to the formulas

$$\begin{aligned} \tau_{i,j} &= \alpha_i + \dots + \alpha_{n-j+1}. \\ \tau_{i,j} &= \begin{cases} \alpha_i + \dots + \alpha_{j-1} + 2(\alpha_j + \dots + \alpha_{n-1}) + \alpha_n & \text{if } j \leq n - 1, \\ \alpha_i + \dots + \alpha_{2n-j} & \text{if } n \leq j \leq 2n - i. \end{cases} \\ \tau_{i,j} &= \begin{cases} \alpha_i + \dots + \alpha_j + 2(\alpha_{j+1} + \dots + \alpha_n) & \text{if } j \leq n - 1, \\ \alpha_i + \dots + \alpha_{2n-j} & \text{if } n \leq j \leq 2n - i. \end{cases} \\ \tau_{i,j} &= \begin{cases} \alpha_i + \dots + \alpha_j + 2(\alpha_{j+1} + \dots + \alpha_{n-2}) + \alpha_{n-1} + \alpha_n & \text{if } j \leq n - 2, \\ \alpha_i + \dots + \alpha_{n-2} + \alpha_n & \text{if } j = n - 1, \\ \alpha_i + \dots + \alpha_{2n-j-1} & \text{if } n \leq j \leq 2n - 1 - i. \end{cases} \end{aligned}$$

in types  $A_n, C_n, B_n, D_n$  respectively. *E.g.*, in types  $A_4, C_3, B_3, D_4$  we have, respectively

$\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4$	$\alpha_1 + \alpha_2 + \alpha_3$	$\alpha_1 + \alpha_2$	$\alpha_1$	
$\alpha_2 + \alpha_3 + \alpha_4$	$\alpha_2 + \alpha_3$	$\alpha_2$		
$\alpha_3 + \alpha_4$	$\alpha_3$			
$\alpha_4$				
$2\alpha_1 + 2\alpha_2 + \alpha_3$	$\alpha_1 + 2\alpha_2 + \alpha_3$	$\alpha_1 + \alpha_2 + \alpha_3$	$\alpha_1 + \alpha_2$	$\alpha_1$
	$2\alpha_2 + \alpha_3$	$\alpha_2 + \alpha_3$	$\alpha_2$	
		$\alpha_3$		

$\alpha_1 + 2\alpha_2 + 2\alpha_3$	$\alpha_1 + \alpha_2 + 2\alpha_3$	$\alpha_1 + \alpha_2 + \alpha_3$	$\alpha_1 + \alpha_2$	$\alpha_1$	
	$\alpha_2 + 2\alpha_3$	$\alpha_2 + \alpha_3$	$\alpha_2$		
		$\alpha_3$			
$\alpha_1 + 2\alpha_2 + \alpha_3 + \alpha_4$	$\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4$	$\alpha_1 + \alpha_2 + \alpha_4$	$\alpha_1 + \alpha_2 + \alpha_3$	$\alpha_1 + \alpha_2$	$\alpha_1$
	$\alpha_2 + \alpha_3 + \alpha_4$	$\alpha_2 + \alpha_4$	$\alpha_2 + \alpha_3$	$\alpha_2$	
		$\alpha_4$	$\alpha_3$		

Then  $\mathcal{I}(X)$  is in bijection with the set  $\mathcal{S}_X$  of subdiagrams of  $T_X$  when  $X = A, B, C$  whereas in type  $D$  one has to consider also the sets of boxes of  $T_D$  which become subdiagrams of  $T_D$  upon switching columns  $n - 1, n$  (see [13] or [3]). In turn to each subdiagram we can associate a lattice path of length  $2n$ , starting from the origin and never going under the  $x$ -axis, with step vectors  $(1, 1), (1, -1)$  (see [10]). The correspondence between subdiagrams and paths is best explained with an example at hand. Let  $n = 9$  and consider, for type  $B_n$  or  $C_n$ , the shifted partition  $(16, 13, 11, 8, 7, 5, 3)$ , see Figure 1 (here, as in Figure 3, the origin coincides with the left upper corner of the diagram, and the  $y$ -axis points downwards).




Figure 1

Fig. 1

Connect the point  $(2n, 0)$  to the border of the subdiagram with an horizontal segment, and consider the zig-zag line formed by the horizontal segment and the right border of the subdiagram. Rotate the figure by  $45^\circ$  in the positive direction and then flip it across a vertical line. After rescaling (in the obvious way) we obtain the desired lattice path. See Figure 2 for the path corresponding to the partition of Figure 1. (To make a comparison easy, the steps which correspond to thick segments in Figure 1 are also made thick in Figure 2.) So we have associated to any subdiagram of  $T_{B_n}$  (or  $T_{C_n}$ ) a lattice path of length  $2n$ . In a similar way we can associate to any subdiagram of  $T_{D_n}$  a lattice path of length  $2n - 1$ .




Fig. 2

Slight modifications are needed to define a correspondence in type  $A_n$ . Start from the point  $(n + 1, 0)$ , reach and follow the right border of the diagram. End in the point  $(0, n + 1)$ : see Figures 3,4 for the case of the partition  $(5, 3, 1, 1, 1, 0, 0)$ , relative to  $A_7$ .




Fig. 3

In type  $A_n$  this correspondence turns out to be a bijection between  $\mathcal{I}(A_n)$  and the set of Dyck paths of length  $2n + 2$ , whereas in types  $B_n, C_n$  one gets a bijection with the set of paths of length  $2n$  not necessarily ending on the  $x$ -axis.

Remark that in cases  $B_n, C_n$  our statistics  $P_X$  translates into the one which counts the number of *returns* of the paths, *i.e.* the number of contact points of the path with the  $x$ -axis minus one. In type  $A_n$  the statistics  $P_X$  counts the number of returns minus one (so the statistics has value 0 for the path of Figure 4).




Fig. 4

Denote by  $\mathcal{B}_{n,h,j}$  the set of paths of the previous type having length  $n$ , ending in the point  $(n, h)$  and having exactly  $j$  returns. The enumeration of such objects has been known since a long time (see 8, §2 for historical details and generalizations). As usual we set

$$\binom{n}{m} = 0 \quad \text{if } m < 0.$$

PROPOSITION 2.1.[6, 13, Cor. 3.2] *Assume  $n \equiv h, \text{ mod } 2$ . Then*

$$(2.1) \quad |\mathcal{B}_{n,h,j}| = \binom{n-(j+1)}{\frac{n+h}{2}-1} - \binom{n-(j+1)}{\frac{n+h}{2}}.$$

Note that if a path has length  $n$  and ends at height  $j$ , then  $n + j$  is even. In particular, if  $n + h$  is odd then  $\mathcal{B}_{n,h,j} = \emptyset$  for any  $j$ . We have immediately

$$P_{A_n}(j) = |\mathcal{B}_{2n+2,0,j+1}| = \frac{j+1}{n+1} \binom{2n-j}{n},$$

$$P_{B_n}(j) = P_{C_n}(j) = \sum_{h=0}^{2n} |\mathcal{B}_{2n,h,j}| = \binom{2n-j-1}{n-1}$$

which are the desired formulas in cases  $A_n, B_n, C_n$ . For type  $D$  we argue as follows. First observe that, in the diagramatic encoding, ideals can be counted as

$$(2.2) \quad 2|\mathcal{S}_{D_n}| - |\mathcal{D}_n|$$

$\mathcal{D}_n$  being the set of subdiagrams of  $T_{D_n}$  having columns  $n-1, n$  of equal length. So we have to understand our statistics on  $\mathcal{S}_{D_n}$  and on  $\mathcal{D}_n$ . Ideals corresponding to subdiagrams in  $\mathcal{S}_{D_n}$  give rise to paths starting from the origin and having length  $2n-1$ . The number of simple roots belonging to  $\Phi_i$  for such an ideal  $i$  is exactly the number of returns of the corresponding path precisely when the ideal does not contain  $\alpha_n$ . In this latter case to get the number of simple roots one has to add 1 to the number of returns. On the other hand the ideals containing  $\alpha_n$  are exactly the ones giving rise to paths ending at height 1. Therefore the piece in degree  $j$  of our statistics coming from  $\mathcal{S}_{D_n}$  is

$$\sum_{h=3}^{2n-1} |\mathcal{B}_{2n-1,h,j}| + |\mathcal{B}_{2n-1,1,j-1}|$$

$$= \binom{2n-j-2}{n} + \binom{2n-j-1}{n-1} - \binom{2n-j-1}{n}$$

$$= \binom{2n-j-2}{n-2}.$$

We have used relation (2.1) to evaluate the left hand side of the previous expression. Now remark that the contribution to the piece of degree  $j$  of our statistics coming from  $\mathcal{D}_n$  is

$$P_{B_{n-1}}(j) - P_{A_{n-2}}(j - 1) + P_{A_{n-2}}(j - 2).$$

Note in fact that to any diagram in  $\mathcal{D}_n$  we can associate a diagram in  $T_{B_{n-1}}$  by deleting the  $n$ -th column. In so doing our statistics counts:

- a) all paths for type  $B_{n-1}$  having  $j$  returns and end point not lying on the  $x$ -axis;
- (b) all paths for type  $B_{n-1}$  having  $j - 1$  returns and end point on the  $x$ -axis.

It is clear that paths for  $B_{n-1}$  having  $k$  returns and end point on the  $x$ -axis are the same as paths for  $A_{n-2}$  with  $k - 1$  returns. Hence contribution (a) is  $P_{B_{n-1}}(j) - P_{A_{n-2}}(j - 1)$ , and contribution (b) is  $P_{A_{n-2}}(j - 2)$ . Relation (3) and some elementary calculations yield the last formula in the Theorem.

REMARK 2.1. It is worth recalling that the value  $P_X(0)$  has a special geometric meaning. Indeed, *ad*-nilpotent ideals correspond to connected components in the dominant chamber of  $\mathfrak{h}_{\mathbb{R}}$  ( $\mathfrak{h}$  being a Cartan subalgebra of  $\mathfrak{g}$ ) determined by the hyperplanes  $(\alpha, x) = 0, (\alpha, x) = 1, \alpha \in \Delta^+$ . More precisely, the open region associated to the ideal  $\mathfrak{i}$  is determined by the inequalities  $0 < (\alpha, x) < 1$  if  $\mathfrak{g}_{\alpha} \not\subset \mathfrak{i}$ , and  $(\alpha, x) > 1$  if  $\mathfrak{g}_{\alpha} \subset \mathfrak{i}$ . Panyushev proved that an ideal in  $\mathcal{I}$  does not contain a simple root space if and only if the corresponding region is bounded (see [12, Proposition 3.7]). He also found the following remarkable formula (see [12, Proposition 3.10]):

$$P_X(0) = \frac{1}{|W|} \prod_{i=1}^n (h + e_i - 1).$$

Here  $W$  is the Weyl group,  $h$  the Coxeter number and  $e_1, \dots, e_n$  the exponents of  $\mathfrak{g}$ .  $P_X(0)$  is also the number of positive clusters.

### 3 – Proof of Theorem 1.2

LEMMA 3.1. *An abelian ideal  $\mathfrak{i} \in \mathcal{I}^{ab}$  may contain at most one simple root space.*

PROOF. Let  $\alpha, \alpha' \in \Pi$  such that  $\mathfrak{g}_{\alpha}, \mathfrak{g}_{\alpha'} \subset \mathfrak{i}$ . Consider a minimal length path from  $\alpha$  to  $\alpha'$  in the Dynkin diagram of  $\mathfrak{g}$ . By Corollaire 3 in [1, VI, 1.7] the sum  $\gamma$  of the simple roots in the path belongs to  $\Delta^+$  as well as  $\gamma - \alpha$ . Moreover  $\gamma > \alpha, \gamma - \alpha > \alpha'$ . Therefore  $\mathfrak{g}_{\gamma} \subset \mathfrak{i}, \mathfrak{g}_{\gamma - \alpha} \subset \mathfrak{i}$ . But  $[\mathfrak{g}_{\alpha}, \mathfrak{g}_{\gamma - \alpha}] = \mathfrak{g}_{\gamma}$ , hence  $\mathfrak{i}$  is not abelian.  $\square$

Recall that an  $ad$ -nilpotent ideal is nilpotent, *i.e.* its descending central series

$$\mathfrak{i} \supset [\mathfrak{i}, \mathfrak{i}] \supset [[\mathfrak{i}, \mathfrak{i}], \mathfrak{i}] \supset [[[ \mathfrak{i}, \mathfrak{i}], \mathfrak{i}], \mathfrak{i}] \supset \dots$$

has a finite number  $n(\mathfrak{i})$  of non zero terms. In particular,  $\mathfrak{i}$  is an abelian ideal if and only if  $n(\mathfrak{i}) \leq 1$ . Also recall that  $ad$ -nilpotent ideals are in canonical bijection with antichains (*i.e.*, subset formed by mutually non-comparable elements) in the root poset. The correspondence is given by mapping an ideal to its minimal roots w.r.t  $<$ , and the inverse map associates to an antichain  $A$  the ideal  $\bigoplus_{\beta \in A} \bigoplus_{\alpha \geq \beta} \mathfrak{g}_\alpha$ . If  $\Pi = \{\alpha_1, \dots, \alpha_n\}$ , denote by  $\theta = \sum_{i=1}^n a_i \alpha_i$  the highest root of  $\Delta$ .

LEMMA 3.2. *Let  $\mathfrak{i}_j = \bigoplus_{\beta \geq \alpha_j} \mathfrak{g}_\alpha$ ,  $1 \leq j \leq n$ . Then*

$$n(\mathfrak{i}_j) = a_j.$$

PROOF. We use the following result of Chari, Dolbin and Ridenour [5, Theorem 1]. Let  $\mathfrak{i}$  an  $ad$ -nilpotent ideal corresponding to the antichain  $A = \{\beta_1, \dots, \beta_k\}$ . Then  $n(\mathfrak{i}) = s$  if and only if  $s$  is the minimal non-negative integer such that  $\beta_{i_1} + \dots + \beta_{i_s} \not\leq \theta$  (repetitions in the  $\beta$  are allowed). The claim follows immediately, because the antichain attached to  $\mathfrak{i}_j$  consists only of  $\alpha_j$ , and  $\theta - a_j \alpha_j = \sum_{i=1}^{j-1} a_i \alpha_i + \sum_{i=j+1}^n a_i \alpha_i$  belongs to the positive root lattice, whereas

$$\theta - (a_j + 1)\alpha_j = \sum_{i=1}^{j-1} a_i \alpha_i - \alpha_j + \sum_{i=j+1}^n a_i \alpha_i$$

does not. □

We are ready to prove Theorem 1.2. The result follows combining (1.1) and Lemma 3.1 if we prove that  $P_X^{ab}(1) = z(\mathfrak{g}) - 1$ . On the other hand Lemma 3.2 implies that  $P_X^{ab}(1)$  equals the number of indices  $i$  such that  $a_i = 1$ . The latter number is known to coincide with  $z(\mathfrak{g}) - 1$  (see [1, VI, §2.3]).

### Acknowledgements

The authors wish to thank Christian Krattenthaler for a useful remark on the paper.



## REFERENCES

- [1] N. BOURBAKI: *Groupes et algèbres de Lie, Chapitres 4-6*, Hermann, Paris (1968).
- [2] P. CELLINI, V. G. KAC, P. MSENEDER FRAJRIA AND P. PAPI: *Decomposition rules for conformal pairs associated to symmetric spaces and abelian subalgebras of  $\mathbb{Z}_2$ -graded Lie algebras*, Adv. Math. **207** (2006), 156-204.
- [3] P. CELLINI AND P. PAPI: *ad-nilpotent ideals of a Borel subalgebra*, J. Algebra **225** (2000), 130-141.
- [4] P. CELLINI AND P. PAPI: *ad-nilpotent ideals of a Borel subalgebra II*, J. Algebra **258** (2002), 112-121.
- [5] V. CHARI, R. J. DOLBIN AND T. RIDENOUR: *Ideals in Parabolic Subalgebras of Simple Lie Algebras*, arXiv:0809.0245 (2008).
- [6] O. ENGELBERG: *On some problems concerning a restricted random walk*, J. Appl. Probab. **2** (1965), 396-404.
- [7] S.V. FOMIN, A.V. ZELEVINSKY: *Y -system and generalized associahedra*, Ann. Math. **158** (2003), 977-1018.
- [8] C. KRATTENTHALER: *Watermelon configurations with wall interaction: exact and asymptotic results*, J. Physics Conf. Series **42** (2006), 179-212.
- [9] B. KOSTANT: *The set of abelian ideals of a Borel subalgebra, Cartan decompositions, and discrete series representations*, Internat. Math. Res. Notices (1998), n. 5, 225-252.
- [10] C. KRATTENTHALER, L. ORSINA, P. PAPI: *Enumeration of ad-nilpotent  $\mathfrak{b}$ -ideals for simple Lie algebras*, Adv. Appl. Math. **28** (2002), 478-522.
- [11] S. KUMAR: *On the Cachazo-Douglas-Seiberg-Witten conjecture for simple Lie algebras*, J. Amer. Math. Soc. **21** (2008), n. 3, 797-808.
- [12] D. PANYUSHEV: *ad-nilpotent ideals: generators and duality*, J. Algebra **274** (2004), 822-846.
- [13] J. SHI: *The number of  $\oplus$ -sign types*, Quart. J. Math. Oxford **48** (1997), 93-105.

*Lavoro pervenuto alla redazione il 20 maggio 2009  
ed accettato per la pubblicazione il 15 giugno 2009.  
Bozze licenziate il 01 ottobre 2009*

## INDIRIZZO DEGLI AUTORI:

Paola Cellini – Dipartimento di Scienze – Università di Chieti-Pescara – Viale Pindaro 42,  
65127 Pescara  
E-mail: cellini@sci.unich.it

Pierluigi Mösener Frajria – Politecnico di Milano – Polo regionale di Como – Via Valleggio  
11, 22100 Como  
E-mail: pierluigi.moseneder@polimi.it

Paolo Papi – Dipartimento di Matematica – Sapienza Università di Roma – P.le A. Moro 2,  
00185, Roma, Italy  
E-mail: papi@mat.uniroma1.it