

# An introduction to Algebraic Geometry - Varieties

Kieran G. O'Grady

5th April 2024

## Contents

<b>Contents</b>	<b>1</b>
<b>0 Introduction</b>	<b>3</b>
<b>1 Quasi projective varieties</b>	<b>7</b>
1.1 Zariski's topology on affine space . . . . .	7
1.2 Zariski's topology on projective space . . . . .	9
1.3 Decomposition into irreducibles . . . . .	10
1.4 The Nullstellensatz . . . . .	12
1.5 Regular maps . . . . .	13
1.6 Regular functions on affine varieties . . . . .	18
1.7 Quasi-projective varieties defined over a subfield of $\mathbb{K}$ . . . . .	20
1.8 Geometry and Algebra . . . . .	23
1.9 Exercises . . . . .	24
<b>2 Algebraic varieties</b>	<b>29</b>
2.1 Introduction . . . . .	29
2.2 Algebraic prevarieties . . . . .	29
2.3 Products and algebraic varieties . . . . .	33
2.4 Complete varieties . . . . .	40
2.5 Algebraic vector bundles . . . . .	42
2.6 Exercises . . . . .	43
<b>3 Rational maps, dimension</b>	<b>47</b>
3.1 Introduction . . . . .	47
3.2 Rational maps . . . . .	47
3.3 Blow-up . . . . .	49
3.4 The field of rational functions . . . . .	53
3.5 Dimension . . . . .	56

<b>4</b>	<b>Tangent space, smooth points</b>	<b>59</b>
4.1	Introduction . . . . .	59
4.2	The local ring of a variety at a point . . . . .	59
4.3	The Zariski tangent and cotangent space . . . . .	60
4.4	Smooth points . . . . .	65
4.5	Criterion for local invertibility of regular maps . . . . .	68
4.6	Differential forms . . . . .	70
<b>5</b>	<b>Curves</b>	<b>71</b>
5.1	Introduction . . . . .	71
5.2	Birational models of curves . . . . .	71
5.3	Divisors on curves and linear equivalence . . . . .	75
5.4	The genus of a curve and the Riemann-Roch Theorem . . . . .	75
5.5	Proof of the Riemann-Roch Theorem . . . . .	76
<b>A</b>	<b>Algebra à la carte</b>	<b>79</b>
A.1	Introduction . . . . .	79
A.2	Unique factorization . . . . .	79
A.3	Noetherian rings . . . . .	79
A.4	Extensions of fields . . . . .	81
A.5	The key to the Nullstellensatz . . . . .	83
A.6	Descent . . . . .	83
A.7	Derivations . . . . .	84
A.8	Nakayama's Lemma . . . . .	86
A.9	Order of vanishing . . . . .	87
	<b>Bibliography</b>	<b>89</b>

# Chapter 0

## Introduction

### Motivation

We will describe some problems and results in order to whet your appetite. Some (or most) of the statements below might leave you puzzled, do not worry, they will become clear later on. In fact one of the goals of reading the book is to be able to understand what is written in the paragraphs below.

We start from the following well known indefinite integral:

$$\int \frac{dx}{\sqrt{1-x^2}} = \arcsin x.$$

What if we ask

$$\int \frac{dx}{\sqrt{1-x^3}} = ?$$

Note that one gets the first integral by writing out the formula for the length of arcs of a circle. Similarly, one gets the second integral, or more generally integrals of functions  $p(x)^{-1/2}$ , where  $p$  is a polynomial of degree 3 (or 4), if one sets out to compute the length of arcs of ellipses. There is no way to express the second integral starting from elementary functions. What Fagnano discovered for similar integrals, and what Euler amplified, is that, although we cannot express the integral via elementary functions, there is a rational addition formula, i.e. there exists a rational function  $F$  of four variables such that for fixed  $l_0$  and varying  $a, b$  we have

$$\int_{l_0}^a \frac{dx}{\sqrt{1-x^3}} + \int_{l_0}^b \frac{dx}{\sqrt{1-x^3}} = \int_{l_0}^c \frac{dx}{\sqrt{1-x^3}} + \text{const},$$

where

$$c = F(a, b, \sqrt{1-a^3}, \sqrt{1-b^3}).$$

Let us sketch a geometric explanation of the addition formula. First of all it is convenient to allow  $x, y$  to be complex numbers. Since couples  $(x, \sqrt{1-x^3})$  are solutions of the equation  $x^3 + y^2 = 1$ , we consider the curve  $C_0 \subset \mathbb{A}^2(\mathbb{C})$  whose equation is  $x^3 + y^2 = 1$ , where  $\mathbb{A}^2(\mathbb{C}) = \mathbb{C}^2$  is the standard complex affine plane. Now  $C_0$  is a complex submanifold of  $\mathbb{A}^2(\mathbb{C})$ , hence a 1-dimensional complex manifold. Since it is not compact, we consider its closure  $C \subset \mathbb{P}^2(\mathbb{C})$  in the projective complex plane. This means adding a single point “at infinity”, namely  $[0, 0, 1]$  (we let  $[T, X, Y]$  be homogeneous coordinates, and  $x = X/T, y = Y/T$ ). Note that by integrating the 1-form  $dx/y$  on  $C$  (as we will do) we do not have to pay attention to which of the two square roots of  $1-x^3$  we choose. A fundamental observation is that  $dx/y$  is holomorphic on all of  $C_0$ , including the points  $(e^{2\pi mi/3}, 0)$  where the denominator vanishes), and moreover it extends to a holomorphic 1-form on all of  $C$ . In order to show that there is an addition formula we fix a line  $R_0 \subset \mathbb{P}^2(\mathbb{C})$  intersecting  $C$  in 3 points  $\bar{p}_1, \bar{p}_2, \bar{p}_3$  and, given another line  $R$  intersecting  $C$  in 3 points  $p_1, p_2, p_3$ , we let

$$\int_{R_0}^R \frac{dx}{y} := \int_{\bar{p}_1}^{p_1} \frac{dx}{y} + \int_{\bar{p}_2}^{p_2} \frac{dx}{y} + \int_{\bar{p}_3}^{p_3} \frac{dx}{y}.$$

Of course in order to make sense of the right hand side one needs to choose paths starting at  $\bar{p}_i$  and ending at  $p_i$  for  $i \in \{1, 2, 3\}$ . By Goursat's Theorem the integrals do not vary if the paths are homotopically equivalent. Hence if we let  $R$  move in a small open subset of  $\mathbb{P}^2(\mathbb{C})^\vee$  we may choose well defined homotopy classes of such paths and the integral above defines a well defined holomorphic function on the open set. There is no way to define a holomorphic function

$$R \xrightarrow{\Phi} \int_{R_0}^R \frac{dx}{y}.$$

on all of  $\mathbb{P}^2(\mathbb{C})^\vee$ : if we define it locally and then we move around, when we come back the value of the function will change by an additive constant. Since it changes by an additive constant, the differential  $d\Phi$  is a well defined holomorphic 1-form  $\omega$  on all of  $\mathbb{P}^2(\mathbb{C})^\vee$  although  $\Phi$  is only well defined locally. Since every holomorphic 1-form on a complex projective space is zero, we get that  $\omega = 0$ , i.e. the (locally defined) function  $\Phi$  is constant. Now notice that the given points  $p_1, p_2 \in C$  there is a unique line  $R$  containing  $p_1, p_2$  (if  $p_1 = p_2$  we let  $R$  be the tangent to  $C$  at  $p_1$ ), and that the coordinates of the third point of intersection of  $R$  and  $C$ , i.e.  $p_3$ , are rational functions of the coordinates of the first two points. This gives the validity of the formula

$$\int_{l_0}^a \frac{dx}{\sqrt{1-x^3}} + \int_{l_0}^b \frac{dx}{\sqrt{1-x^3}} = - \int_{l_0}^c \frac{dx}{\sqrt{1-x^3}} + \text{const},$$

where  $c$  is a rational function of  $(a, b, \sqrt{1-a^3}, \sqrt{1-b^3})$ . With a little more work one gets from this the addition formula as formulated above.

Next we ask more in general what can be said about integrals of the form

$$\int \frac{dx}{\sqrt{D(x)}}, \tag{0.0.1}$$

where  $D(x)$  is a polynomial. For simplicity we assume that  $D(x)$  has no multiple roots. If  $D(x)$  has degree 3, then the arguments above apply verbatim to give an addition formula. In general, the first step is to consider the curve  $C_0 \subset \mathbb{A}^2(\mathbb{C})$  whose equation is  $y^2 = D(x)$ . This is a 1-dimensional complex submanifold of  $\mathbb{A}^2(\mathbb{C})$ . Since it is not compact it is convenient to compactify. The closure of  $C_0$  in  $\mathbb{P}^2(\mathbb{C})$  is compact, but if the degree of  $D(x)$  is greater than 3 then the closure of  $C_0$  is not a submanifold of  $\mathbb{P}^2(\mathbb{C})$  at its unique "point at infinity" (i.e.  $[0, 0, 1]$ ). Nonetheless there is 1-dimensional complex manifold  $C$  containing  $C_0$  as an open dense subset, in fact  $C \setminus C_0$  consists of a single point if  $D(x)$  has odd degree, and consists of two points if  $D(x)$  has even degree. The qualitative behaviour of the integral that we set out to study is determined by the topology of  $C$ . The  $C^\infty$  manifold underlying  $C$  is connected, compact and orientable surface. By the classification compact surfaces it is homeomorphic to a connected sum of  $g$  tori. In fact one show that

$$g = \left\lfloor \frac{\deg D - 1}{2} \right\rfloor. \tag{0.0.2}$$

For example, if  $D$  has degree 3 then  $g = 1$ , i.e.  $C$  is a torus. Suppose that  $g > 1$ . Then there exists an addition formula, but it involves the addition of vectors in  $\mathbb{C}^g$  obtained by integrating the  $g$  linearly independent holomorphic 1-forms

$$\frac{dx}{y}, \frac{xdx}{y}, \dots, \frac{x^{g-1}dx}{y}. \tag{0.0.3}$$

Lastly we discuss how the topological quantity  $g$  (the genus of  $C$ ) controls the arithmetic of  $C$ . Suppose that the polynomial  $p(x)$  has integer coefficients. If  $p$  is a prime we let  $\bar{D}(x) \in \mathbb{F}_p[x]$  be the polynomial whose coefficients are the equivalence classes of the coefficients of  $D$  - we say that  $\bar{D}(x)$  is obtained from  $D$  reducing modulo  $p$ . We suppose that  $\bar{D}(x)$  has the same degree as  $D$  (i.e.  $p$  does not divide the leading coefficient of  $D$ ), and that  $\bar{D}(x)$  does not have multiple roots in the algebraic closure of  $\mathbb{F}_p$ . We also assume that  $p \neq 2$ . For  $n \geq 1$  let  $\mathbb{F}_{p^n}$  be the finite field of cardinality  $p^n$ , and let  $C(\mathbb{F}_{p^n})$  be

---

the set of solutions in  $\mathbb{F}_{p^n}$  of the equation  $y^2 = \overline{D}(x)$ . We view the points at infinity (there is one if  $\deg D$  is odd and two if  $\deg D$  is even) as solutions “in  $\mathbb{F}_{p^n}$ ”. A convenient generating function for the cardinalities  $|C(\mathbb{F}_{p^n})|$  is given by Weil’s zeta function

$$Z(C, T) := \exp \left( \sum_{n=1}^{\infty} \frac{|C(\mathbb{F}_{p^n})|}{n} T^n \right). \quad (0.0.4)$$

A famous theorem of Weil states that

$$Z(C, T) = \frac{\prod_{i=1}^{2g} (1 - a_i T)}{(1 - T)(1 - pT)}, \quad (0.0.5)$$

where each  $a_i$  is an algebraic integer of modulus  $p^{1/2}$  (the last statement is an analogue of Riemann’s hypothesis). This shows that the topological genus  $g$  can be extracted from the number of solutions  $(x, y) \in \mathbb{A}^2(\mathbb{F}_{p^n})$  of the equation  $y^2 = \overline{D}(x)$ . We also see that there is an explicit formula giving the cardinality  $|C(\mathbb{F}_{p^n})|$  for all  $n$  once we know the cardinalities  $|C(\mathbb{F}_p)|, |C(\mathbb{F}_{p^2})|, \dots, |C(\mathbb{F}_{p^{2g}})|$ . The function of  $s$  obtained by making the substitution  $T = p^{-s}$ , i.e.  $Z(C, p^{-s})$ , is a precise analogue of Riemann’s zeta function  $\zeta(s)$ , and the statement that each  $a_i$  has modulus  $p^{1/2}$  is the analogue of the Riemann Hypothesis. It is very compelling evidence in favour of the validity of the Riemann Hypothesis.



# Chapter 1

## Quasi projective varieties

Throughout the book  $\mathbb{K}$  is an algebraically closed field, e.g.  $\mathbb{K} = \mathbb{C}$  or  $\overline{\mathbb{Q}}$ , the algebraic closure of the rational field  $\mathbb{Q}$ , or  $\overline{\mathbb{F}_p}$ , the algebraic closure of the finite field  $\mathbb{F}_p$  where  $p$  is a prime. We are interested in understanding the set of solutions  $(z_1, \dots, z_n) \in \mathbb{K}^n$  of a family of polynomial equations

$$f_1(z_1, \dots, z_n) = 0, \dots, f_r(z_1, \dots, z_n) = 0.$$

“Polynomial equations” means each  $f_i$  is an element of the polynomial ring  $\mathbb{K}[z_1, \dots, z_n]$ .

In order to understand the geometry of a set of solutions of polynomial equations, it is convenient to replace affine space  $\mathbb{A}^n(\mathbb{K})$  by projective space  $\mathbb{P}^n(\mathbb{K})$ , and consider the set of points in  $\mathbb{P}^n(\mathbb{K})$  which are solutions of homogeneous polynomial equations in the homogeneous coordinates. As motivation for this step we recall that results in projective geometry are usually cleaner than in affine geometry - for example two distinct lines in a projective plane have exactly one point of intersection, while two distinct lines in an affine line may intersect in one point or be disjoint. If  $\mathbb{K} = \mathbb{C}$  we may guess that passing to projective space makes life simpler because  $\mathbb{P}^n(\mathbb{C})$  with the classical topology is compact, while  $\mathbb{A}^n(\mathbb{C})$  is not (unless  $n = 0$ ).

Whenever there is no possibility of a misunderstanding we omit  $\mathbb{K}$  from the notation for affine and projective space, i.e.  $\mathbb{A}^n$  is  $\mathbb{A}^n(\mathbb{K})$  and  $\mathbb{P}^n$  is  $\mathbb{P}^n(\mathbb{K})$ .

### 1.1 Zariski’s topology on affine space

If  $f_1, \dots, f_r \in \mathbb{K}[z_1, \dots, z_n]$ , we let

$$V(f_1, \dots, f_r) := \{z \in \mathbb{A}^n \mid f_i(z) = 0 \ \forall i \in \{1, \dots, r\}\}. \quad (1.1.1)$$

More generally, if  $I \subset \mathbb{K}[z_1, \dots, z_n]$  is an ideal (note: the inclusion sign  $\subset$  does not mean strict inclusion, and similarly for  $\supset$ ) we let

$$V(I) := \{z \in \mathbb{A}^n \mid f(z) = 0 \ \forall f \in I\}. \quad (1.1.2)$$

Unless  $n = 0$  or  $I = 0$  an ideal  $I$  of  $\mathbb{K}[z_1, \dots, z_n]$  has an infinite number of elements so that  $V(I)$  is the set of solutions of an infinite set of polynomial equations. However  $I$  has a finite set of generators  $f_1, \dots, f_r$  by Hilbert’s basis Theorem A.3.6, and it follows that  $V(I) = V(f_1, \dots, f_r)$ . In fact it is clear that  $V(I) \subset V(f_1, \dots, f_r)$ . For the reverse inclusion  $V(f_1, \dots, f_r) \subset V(I)$  notice that if  $z \in V(f_1, \dots, f_r)$  and  $f \in I$ , then  $f = \sum_{i=1}^r g_i f_i$  for suitable  $g_1, \dots, g_r \in \mathbb{K}[z_1, \dots, z_n]$  and hence  $f(z) = \sum_{i=1}^r g_i(z) f_i(z) = 0$ .

An elementary observation is that passing from ideals to their zero sets reverses inclusion, i.e. if  $I, J \subset \mathbb{K}[z_1, \dots, z_n]$  are ideals then

$$I \subset J \text{ implies that } V(I) \supset V(J). \quad (1.1.3)$$

**Proposition 1.1.1.** *The collection of subsets  $V(I) \subset \mathbb{A}^n$ , where  $I$  runs through the collection of ideals of  $\mathbb{K}[z_1, \dots, z_n]$ , satisfies the axioms for the closed subsets of a topological space.*

*Proof.* We have  $\emptyset = V((1))$ ,  $\mathbb{A}^n = V((0))$ .

Let  $I, J \subset \mathbb{K}[z_1, \dots, z_n]$  be ideals. We claim that  $V(I) \cup V(J) = V(I \cap J)$ . We have  $V(I), V(J) \subset V(I \cap J)$ , because  $I, J \supset I \cap J$ . Thus  $V(I) \cup V(J) \subset V(I \cap J)$ . Hence it suffices to show that if  $z \in V(I \cap J)$  and  $z \notin V(I)$ , then  $z \in V(J)$ . Since  $z \notin V(I)$ , there exists  $f \in I$  such that  $f(z) \neq 0$ . If  $g \in J$ , then  $f \cdot g \in I \cap J$ , and thus  $(f \cdot g)(z) = 0$  because  $z \in V(I \cap J)$ . Since  $f(z) \neq 0$ , it follows that  $g(z) = 0$ . This proves that  $z \in V(J)$ .

Lastly, let  $\{I_t\}_{t \in T}$  be a family of ideals of  $\mathbb{K}[z_1, \dots, z_n]$ . Then

$$\bigcap_{t \in T} V(I_t) = V(\langle \{I_t\}_{t \in T} \rangle),$$

where  $\langle \{I_t\}_{t \in T} \rangle$  is the ideal generated by the collection of the  $I_t$ 's. □

**Definition 1.1.2.** The *Zariski topology* of  $\mathbb{A}^n$  is the topology whose closed sets are the sets  $V(I)$ , where  $I$  runs through the collection of ideals of  $\mathbb{K}[z_1, \dots, z_n]$ . The Zariski topology of a subset  $A \subset \mathbb{A}^n$  is the topology induced by the Zariski topology of  $\mathbb{A}^n$ .

*Remark 1.1.3.* If  $\mathbb{K} = \mathbb{C}$ , the Zariski topology is weaker than the classical topology of  $\mathbb{A}^n$ . In fact, unless  $n = 0$ , the Zariski is much weaker than the classical topology, in particular it is *not* Hausdorff.

*Example 1.1.4.* A subset  $X \subset \mathbb{A}^n$  is a *hypersurface* if it is equal to  $V(f)$ , where  $f$  is a non constant homogeneous polynomial.

A picture of a hypersurface in  $\mathbb{A}^2$  is in Figure 1.1. Notice that  $(x, y)$  are the affine coordinates - in general, whenever we consider affine or projective space of small dimension, we will denote affine or homogeneous coordinates by letters  $x, y, z, \dots$  and  $X, Y, Z, \dots$  respectively.

What is the field  $\mathbb{K}$ ? The picture shows points with real coordinates. We can view the picture as a “slice” of the corresponding hypersurface over  $\mathbb{C}$ , or as the closure (either in the Zariski or the classical topology) of the corresponding hypersurface over the algebraic closure of the rationals  $\overline{\mathbb{Q}}$ .




Figure 1.1:  $(x^2 + 2y^2 - 1)(3x^2 + y^2 - 1) + \frac{3}{100} = 0$

Given a subset  $X \subset \mathbb{A}^n$ , let

$$I(X) := \{f \in \mathbb{K}[z_1, \dots, z_n] \mid f(z) = 0 \text{ for all } z \in X\}. \quad (1.1.4)$$

Clearly  $I(X)$  is an ideal of  $\mathbb{K}[z_1, \dots, z_n]$  and  $X$  is contained in the closed set  $V(I(X))$ . Moreover  $V(I(X))$  is the closure of  $X$  in the Zariski topology. In fact suppose that  $V(J) \subset \mathbb{A}^n$  is a closed



subset containing  $X$ . Then  $f(z) = 0$  for all  $f \in J$  and  $z \in X$ , and hence  $J \subset I(X)$ . This shows that  $V(J) \supset V(I(X))$  (recall (1.1.3)).

*Remark 1.1.5.* Let  $\mathcal{A}$  be a finite dimensional affine space over  $\mathbb{K}$  of dimension  $n$ . Then the Zariski topology on  $\mathcal{A}$  may be defined by analogy with the case of  $\mathbb{A}^n$ , simply replacing  $\mathbb{K}[z_1, \dots, z_n]$  by the  $\mathbb{K}$  algebra of polynomial functions on  $\mathcal{A}$  (which is isomorphic to  $\mathbb{K}[z_1, \dots, z_n]$ ). Another way of putting it is that an affine transformation of  $\mathbb{A}^n$  is a homeomorphism for the Zariski topology.

## 1.2 Zariski's topology on projective space

Let  $F \in \mathbb{K}[Z_0, \dots, Z_n]_d$  be homogeneous of degree  $d$  (to be correct we should say that  $F$  belongs to the homogeneous summand of degree  $d$ , because the degree of 0 is  $-\infty$ ). Let  $x = [Z] \in \mathbb{P}^n$ . Then  $F(Z) = 0$  if and only if  $F(\lambda Z) = 0$  for every  $\lambda \in \mathbb{K}^*$ , because  $F(\lambda Z) = \lambda^d F(Z)$ . Hence, although  $F(x)$  is not defined, it makes to state that  $F(x) = 0$  or  $F(x) \neq 0$ . Thus if  $F_1, \dots, F_r \in \mathbb{K}[Z_0, \dots, Z_n]$  are homogeneous (of possibly different degrees) it makes sense to let

$$V(F_1, \dots, F_r) := \{x \in \mathbb{P}^n \mid F_1(x) = \dots = F_r(x) = 0\}. \quad (1.2.1)$$

As in the case of affine space, it is convenient to consider the zero locus of ideals, but we need to consider homogeneous ideals. An ideal  $I \subset \mathbb{K}[Z_0, \dots, Z_n]$  is *homogeneous* if

$$I = \bigoplus_{d=0}^{\infty} I \cap \mathbb{K}[Z_0, \dots, Z_n]_d, \quad (1.2.2)$$

i.e. if it is generated by homogeneous elements. Let  $I \subset \mathbb{K}[Z_0, \dots, Z_n]$  be a homogeneous ideal; we let

$$V(I) := \{x \in \mathbb{P}^n \mid F(x) = 0 \ \forall \text{ homogeneous } F \in I\}.$$

By Hilbert's basis Theorem A.3.6  $I$  is generated by a finite set of homogeneous polynomials  $F_1, \dots, F_r$ , and hence  $V(I) = V(F_1, \dots, F_r)$ . Notice that if  $I \subset \mathbb{K}[Z_0, \dots, Z_n]$  is a homogeneous ideal we have two different meanings for  $V(I)$ , namely the subset of  $\mathbb{P}^n$  defined above and the subset of  $\mathbb{A}^{n+1}$  defined in (1.1.2). The context will indicate which of the two we mean.

Proceeding as in the proof of Proposition 1.1.1 one gets the following result.

**Proposition 1.2.1.** *The collection of subsets  $V(I) \subset \mathbb{P}^n$ , where  $I$  runs through the collection of homogeneous ideals of  $\mathbb{K}[Z_0, \dots, Z_n]$ , satisfies the axioms for the closed subsets of a topological space.*

**Definition 1.2.2.** The *Zariski topology* of  $\mathbb{P}^n$  is the topology whose closed sets are the sets  $V(I) \subset \mathbb{P}^n$ , where  $I$  runs through the collection of homogeneous ideals of  $\mathbb{K}[Z_0, \dots, Z_n]$ . The Zariski topology of a subset  $A \subset \mathbb{P}^n$  is the topology induced by the Zariski topology of  $\mathbb{P}^n$ .

*Remark 1.2.3.* Let  $\pi: (\mathbb{K}^{n+1} \setminus \{0\}) \rightarrow \mathbb{P}^n$  be the map defined by  $\pi(Z) = [Z]$ , so that  $\mathbb{P}^n$  is identified as the quotient of  $\mathbb{K}^{n+1} \setminus \{0\}$  for the action by homotheties. The Zariski topology of  $\mathbb{P}^n$  is the quotient of the Zariski topology on  $\mathbb{K}^{n+1} \setminus \{0\}$ .

*Remark 1.2.4.* If  $F \in \mathbb{K}[Z_0, \dots, Z_n]$  is homogeneous we let

$$\mathbb{P}_F^n := \mathbb{P}^n \setminus V(F). \quad (1.2.3)$$

Thus  $\mathbb{P}_F^n$  is an open subset of  $\mathbb{P}^n$ .

From now on we make the identification

$$\begin{array}{ccc} \mathbb{A}^n & \longleftrightarrow & \mathbb{P}_{Z_0}^n \\ (z_1, \dots, z_n) & \mapsto & [1, z_1, \dots, z_n] \end{array}$$

The Zariski topology of  $\mathbb{A}^n$  induced by the Zariski topology on  $\mathbb{P}^n$  is the same as the Zariski topology of Definition 1.1.2. In fact let  $X \subset \mathbb{A}^n$ . Suppose first that  $X$  is closed for the topology induced

from the Zariski topology of  $\mathbb{P}^n$ , i.e.  $X = (\mathbb{P}_{Z_0}^n) \cap V(F_1, \dots, F_r)$ , where each  $F_j \in \mathbb{K}[Z_0, Z_1, \dots, Z_n]$  is homogeneous. Then  $X = V(f_1, \dots, f_r)$ , where

$$f_j(z_1, \dots, z_n) := F(1, z_1, \dots, z_n).$$

Next suppose that  $X$  is closed for the Zariski topology of Definition 1.1.2, i.e.  $X = V(f_1, \dots, f_r)$  where  $f_1, \dots, f_r \in \mathbb{K}[z_1, \dots, z_n]$ . We may assume that all  $f_j$  are non zero because  $\mathbb{A}^n$  is clearly closed for the induced topology, and hence each  $f_j$  has a well defined degree  $d_j$ . For  $j \in \{1, \dots, r\}$  let

$$F_j(Z_0, \dots, Z_n) := Z_0^{d_j} f\left(\frac{Z_1}{Z_0}, \dots, \frac{Z_n}{Z_0}\right).$$

Then  $F_j$  is a homogeneous polynomial of degree  $d_j$  and hence  $V(F_1, \dots, F_r) \subset \mathbb{P}^n$  is a closed subset. Since

$$V(f_1, \dots, f_r) = (\mathbb{P}_{Z_0}^n) \cap V(F_1, \dots, F_r),$$

we get that  $V(f_1, \dots, f_r)$  is closed for the induced topology.

*Example 1.2.5.* A subset  $X \subset \mathbb{P}^n$  is a *hypersurface* if it is equal to  $V(F)$ , where  $F$  is a non constant homogeneous polynomial. Notice that  $V(F) \cap \mathbb{A}^n$  is a hypersurface unless  $F = cZ_0^d$  for some  $c \in \mathbb{K}^*$ .

Given a subset  $A \subset \mathbb{P}^n$ , let

$$I(A) := \langle F \in \mathbb{K}[Z_0, \dots, Z_n] \mid F \text{ is homogeneous and } F(p) = 0 \text{ for all } p \in A \rangle, \quad (1.2.4)$$

where  $\langle \cdot \rangle$  means “the ideal generated by”. Clearly  $I(A)$  is a homogeneous ideal of  $\mathbb{K}[Z_0, \dots, Z_n]$ , and  $V(I(A))$  is the closure of  $A$  in the Zariski topology.

**Definition 1.2.6.** A *quasi-projective* variety is a Zariski locally closed subset of a projective space, i.e.  $X \subset \mathbb{P}^n$  such that  $X = U \cap Y$ , where  $U, Y \subset \mathbb{P}^n$  are Zariski open and Zariski closed respectively.

*Example 1.2.7.* By Remark 1.2.4, every closed subset of  $\mathbb{A}^n$  is a quasi projective variety.

*Remark 1.2.8.* If  $V$  is a finite dimensional complex vector space, the Zariski topology on  $\mathbb{P}(V)$  is defined by imitating what was done for  $\mathbb{P}^n$ : one associates to a homogeneous ideal  $I \subset \text{Sym } V^\vee$  the set of zeroes  $V(I)$ , etc. Everything that we do in the present chapter applies to this situation, but for the sake of concreteness we formulate it for  $\mathbb{P}^n$ .

### 1.3 Decomposition into irreducibles

A proper closed subset  $X \subset \mathbb{P}^1$  (or  $X \subset \mathbb{A}^1$ ) is a finite set of points. In general, a quasi projective variety is a finite union of closed subsets which are irreducible, i.e. are not the union of proper closed subsets. In order to formulate the relevant result, we give a few definitions.

**Definition 1.3.1.** Let  $X$  be a topological space. We say that  $X$  is *reducible* if either  $X = \emptyset$  or there exist proper closed subsets  $Y, W \subset X$  such that  $X = Y \cup W$ . We say that  $X$  is *irreducible* if it is not reducible.

*Example 1.3.2.* A subset  $A \subset \mathbb{R}^n$  with the euclidean (classical) topology is irreducible if and only if it is a singleton.

*Example 1.3.3.* Projective space  $\mathbb{P}^n$  with the Zariski topology is irreducible. In fact suppose that  $\mathbb{P}^n = X \cup Y$  with  $X$  and  $Y$  proper closed subsets. Then there exist homogeneous  $F \in I(X)$  and  $G \in I(Y)$  such that  $F(y) \neq 0$  for one (at least)  $y \in Y$  and  $G(x) \neq 0$  for one (at least)  $x \in X$ . In particular both  $F$  and  $G$  are non zero, and hence  $FG \neq 0$  because  $\mathbb{K}[Z_0, \dots, Z_n]$  is an integral domain. On the other hand  $FG = 0$  because  $\mathbb{P}^n = Y \cup W$ . This is a contradiction, and hence  $\mathbb{P}^n$  is irreducible.

*Remark 1.3.4.* Since the field  $\mathbb{K}$  is algebraically closed it is infinite, and hence there is no distinction between the polynomial ring  $\mathbb{K}[z_1, \dots, z_n]$  and the ring of polynomial functions in  $z_1, \dots, z_n$ . That is implicit in the argument given in Example 1.3.3, and it will appear repeatedly.

**Definition 1.3.5.** Let  $X$  be a topological space. An *irreducible decomposition* of  $X$  consists of a decomposition (possibly empty)

$$X = X_1 \cup \dots \cup X_r \tag{1.3.1}$$

where each  $X_i$  is a closed irreducible subset of  $X$  (irreducible with respect to the induced topology) and moreover  $X_i \not\subset X_j$  for all  $i \neq j$ .

We will prove the following result.

**Theorem 1.3.6.** *Let  $A \subset \mathbb{P}^n$  with the (induced) Zariski topology. Then  $A$  admits an irreducible decomposition, and such a decomposition is unique up to reordering of components.*

The key step in the proof of Theorem 1.3.6 is the following remarkable consequence of Hilbert's basis Theorem A.3.6.

**Proposition 1.3.7.** *Let  $A \subset \mathbb{P}^n$ , and let  $A \supset X_0 \supset X_1 \supset \dots \supset X_m \supset \dots$  be a descending chain of Zariski closed subsets of  $A$ , i.e.  $X_m \supset X_{m+1}$  for all  $m \in \mathbb{N}$ . Then the chain is stationary, i.e. there exists  $m_0 \in \mathbb{N}$  such that  $X_m = X_{m_0}$  for  $m \geq m_0$ .*

*Proof.* Let  $\bar{X}_i$  be the closure of  $X_i$  in  $\mathbb{P}^n$ . Then  $X_i = A \cap \bar{X}_i$ , because  $X_i$  is closed in  $A$ . Hence we may replace  $X_i$  by  $\bar{X}_i$ , or equivalently we may suppose that the  $X_i$  are closed in  $\mathbb{P}^n$ . Let  $I_m = I(X_m)$ . Then  $I_0 \subset I_1 \subset \dots \subset I_m \subset \dots$  is an ascending chain of (homogeneous) ideals of  $\mathbb{K}[Z_0, \dots, Z_n]$ . By Hilbert's basis Theorem and Lemma A.3.3 the ascending chain of ideals is stationary, i.e. there exists  $m_0 \in \mathbb{N}$  such that  $I_m = I_{m_0}$  for  $m \geq m_0$ . Thus  $X_{m_0} = V(I_{m_0}) = V(I_m) = X_m$  for  $m \geq m_0$ .  $\square$

*Proof of Theorem 1.3.6.* If  $A$  is empty, then it is the empty union (of irreducibles). Next, suppose that  $A$  is not empty and that it does not admit an irreducible decomposition; we will arrive at a contradiction. First  $A$  is reducible, i.e.  $A = X_0 \cup W_0$  with  $X_0, W_0 \subset A$  proper closed subsets. If both  $X_0$  and  $W_0$  have an irreducible decomposition, then  $A$  is the union of the irreducible components of  $X_0$  and  $W_0$ , contradicting the assumption that  $A$  does not admit an irreducible decomposition. Hence one of  $X_0, W_0$ , say  $X_0$ , does *not* have an irreducible decomposition. In particular  $X_0$  is reducible. Thus  $X_0 = X_1 \cup W_1$  with  $X_1, W_1 \subset X_0$  proper closed subsets, and arguing as above, one of  $X_1, W_1$ , say  $X_1$ , does not admit a decomposition into irreducibles. Iterating, we get a strictly descending chain of closed subsets

$$A \supset X_0 \supset X_1 \supset \dots \supset X_m \supset X_{m+1} \supset \dots$$

This contradicts Proposition 1.3.7. This proves that  $X$  has a decomposition into irreducibles  $X = X_1 \cup \dots \cup X_r$ .

By discarding  $X_i$ 's which are contained in  $X_j$  with  $i \neq j$ , we may assume that if  $i \neq j$ , then  $X_i$  is not contained in  $X_j$ .

Lastly, let us prove that such a decomposition is unique up to reordering, by induction on  $r$ . The case  $r = 1$  is trivially true. Let  $r \geq 2$ . Suppose that  $X = Y_1 \cup \dots \cup Y_s$ , where each  $Y_j$  is Zariski closed irreducible, and  $Y_j \not\subset Y_k$  if  $j \neq k$ . Since  $Y_s$  is irreducible, there exists  $i$  such that  $Y_s \subset X_i$ . We may assume that  $i = r$ . By the same argument, there exists  $j$  such that  $X_r \subset Y_j$ . Thus  $Y_s \subset X_r \subset Y_j$ . It follows that  $j = s$ , and hence  $Y_s = X_r$ . It follows that  $X_1 \cup \dots \cup X_{r-1} = Y_1 \cup \dots \cup Y_{s-1}$ , and hence the decomposition is unique up to reordering by the inductive hypothesis.  $\square$

**Definition 1.3.8.** Let  $X$  be a quasi projective variety, and let

$$X = X_1 \cup \dots \cup X_r$$

be an irreducible decomposition of  $X$ . The  $X_i$ 's are the *irreducible components* of  $X$  (this makes sense because, by Theorem 1.3.6, the collection of the  $X_i$ 's is uniquely determined by  $X$ ).

We notice the following consequence of Proposition 1.3.7.

**Corollary 1.3.9.** *A quasi projective variety  $X$  (with the Zariski topology) is quasi compact, i.e. every open covering of  $X$  has a finite subcover.*

The following result makes a connection between irreducibility and algebra.

**Proposition 1.3.10.** *A subset  $X \subset \mathbb{P}^n$  is irreducible if and only if  $I(X)$  is a prime ideal.*

*Proof.* The proof has essentially been given in Example 1.3.3. Suppose that  $X$  is irreducible. In particular  $X \neq \emptyset$  (by definition), and hence  $I(X)$  is a proper ideal of  $\mathbb{K}[Z_0, \dots, Z_n]$ . We must prove that  $\mathbb{K}[Z_0, \dots, Z_n]/I(X)$  is an integral domain. Suppose the contrary. Then there exist

$$F, G \in \mathbb{K}[Z_0, \dots, Z_n], \quad F \notin I(X), \quad G \notin I(X), \quad (1.3.2)$$

such that

$$F \cdot G \in I(X). \quad (1.3.3)$$

By (1.3.2) both  $X \cap V(F)$  and  $X \cap V(G)$  are proper closed subsets of  $X$ , and by (1.3.3) we have  $X = (X \cap V(F)) \cup (X \cap V(G))$ . This is a contradiction, hence  $I(X)$  is a prime ideal.

Next, assume that  $X$  is reducible; we must prove that  $I(X)$  is not prime. If  $X = \emptyset$ , then  $I(X) = \mathbb{K}[Z_0, \dots, Z_n]$  and hence  $I(X)$  is not prime. Thus we may assume that  $X \neq \emptyset$ , and hence there exist proper closed subset  $Y, W \subset X$  such that  $X = Y \cup W$ . Since  $Y \not\subset W$  and  $W \not\subset Y$ , there exist  $F \in (I(Y) \setminus I(W))$  and  $G \in (I(W) \setminus I(Y))$ . It follows that both (1.3.2) and (1.3.3) hold, and hence  $I(X)$  is not prime.  $\square$

*Remark 1.3.11.* Let  $I := (Z_0^2) \subset \mathbb{K}[Z_0, Z_1]$ . Then  $V(I) = \{[0, 1]\}$  is irreducible although  $I$  is *not* prime. Of course  $I(V(I))$  is prime, it equals  $(Z_0)$ .

*Remark 1.3.12.* Let  $X \subset \mathbb{A}^n$ . Let  $I(X) \subset \mathbb{K}[z_1, \dots, z_n]$  be the ideal of polynomials vanishing on  $X$ . Then  $X$  is irreducible if and only if  $I(X)$  is a prime ideal. The proof is analogous to the proof of Proposition 1.3.10. One may also directly relate  $I(X)$  with the ideal  $J \subset \mathbb{K}[Z_0, \dots, Z_n]$  generated by homogeneous polynomials vanishing on  $X$  (as subset of  $\mathbb{P}^n$ ), and argue that  $I(X)$  is prime if and only if  $J$  is.

## 1.4 The Nullstellensatz

Let an ideal  $I$  in a ring  $R$ . The *radical* of  $I$ , denoted by  $\sqrt{I}$ , is the set of elements  $a \in R$  such that  $a^m \in I$  for some  $m \in \mathbb{N}$ . As is easily checked,  $\sqrt{I}$  is an ideal. It is clear that  $\sqrt{I} \subset I(V(I))$ . The Nullstellensatz states that we have equality.

**Theorem 1.4.1** (Hilbert's Nullstellensatz). *Let  $I \subset \mathbb{K}[z_1, \dots, z_n]$  be an ideal. Then  $I(V(I)) = \sqrt{I}$ .*

Before discussing the proof of the Nullstellensatz, we introduce some notation. For  $a = (a_1, \dots, a_n)$  an element of  $\mathbb{A}^n$ , let

$$\mathfrak{m}_a := (z_1 - a_1, \dots, z_n - a_n) = \{f \in \mathbb{K}[z_1, \dots, z_n] \mid f(a_1, \dots, a_n) = 0\}. \quad (1.4.1)$$

Notice that  $\mathfrak{m}_a$  is the kernel of the surjective homomorphism

$$\begin{array}{ccc} \mathbb{K}[z_1, \dots, z_n] & \xrightarrow{\phi} & \mathbb{K} \\ f & \mapsto & f(a_1, \dots, a_n), \end{array}$$

and hence is a maximal ideal. The Nullstellensatz is a consequence of the following result.

**Proposition 1.4.2.** *An ideal  $\mathfrak{m} \subset \mathbb{K}[z_1, \dots, z_n]$  is maximal if and only if there exists  $(a_1, \dots, a_n) \in \mathbb{A}^n$  such that  $\mathfrak{m} = \mathfrak{m}_a$ .*

*Proof.* We have shown that  $\mathfrak{m}_a$  is maximal. Now suppose that  $\mathfrak{m} \subset \mathbb{K}[z_1, \dots, z_n]$  is a maximal ideal. Let  $F := \mathbb{K}[z_1, \dots, z_n]/\mathfrak{m}$ . Then  $F$  is an algebraic extension of  $\mathbb{K}$  by Corollary A.5.2. Since  $\mathbb{K}$  is algebraically closed  $F = \mathbb{K}$ , and hence the quotient map is

$$\mathbb{K}[z_1, \dots, z_n] \xrightarrow{\phi} \mathbb{K}[z_1, \dots, z_n]/\mathfrak{m} = \mathbb{K}.$$

For  $i \in \{1, \dots, n\}$  let  $a_i := \phi(z_i)$ . Then  $(z_i - a_i) \in \ker \phi$ . Since  $\mathfrak{m}_a$  is generated by  $(z_1 - a_1), \dots, (z_n - a_n)$  it follows that  $\mathfrak{m}_a \subset \mathfrak{m}$ . Since both  $\mathfrak{m}_a$  and  $\mathfrak{m}$  are maximal it follows that  $\mathfrak{m} = \mathfrak{m}_a$ .  $\square$

**Corollary 1.4.3** (Weak Nullstellensatz). *Let  $I \subset \mathbb{K}[z_1, \dots, z_n]$  be an ideal. Then  $V(I) = \emptyset$  if and only if  $I = (1)$ .*

*Proof.* If  $I = (1)$ , then  $V(I) = \emptyset$ . Assume that  $V(I) = \emptyset$ . Suppose that  $I \neq (1)$ . Then there exists a maximal ideal  $\mathfrak{m} \subset \mathbb{K}[z_1, \dots, z_n]$  containing  $I$ . Since  $I \subset \mathfrak{m}$ ,  $V(I) \supset V(\mathfrak{m})$ . By Proposition 1.4.2 there exists  $a \in \mathbb{K}^n$  such that  $\mathfrak{m} = \mathfrak{m}_a$  and hence  $V(\mathfrak{m}) = V(\mathfrak{m}_a) = \{(a_1, \dots, a_n)\}$ . Thus  $a \in V(I)$  and hence  $V(I) \neq \emptyset$ . This is a contradiction, and hence  $I = (1)$ .  $\square$

*Proof of Hilbert's Nullstellensatz (Rabinowitz's trick).* Let  $f \in I(V(I))$ . By Hilbert's basis theorem  $I = (g_1, \dots, g_s)$  for  $g_1, \dots, g_s \in \mathbb{K}[z_1, \dots, z_n]$ . Let  $J \subset \mathbb{K}[z_1, \dots, z_n, w]$  be the ideal

$$J := (g_1, \dots, g_s, f \cdot w - 1).$$

Since  $f \in I(V(I))$  we have  $V(J) = \emptyset$  and hence by the Weak Nullstellensatz  $J = (1)$ . Thus there exist  $h_1, \dots, h_s, h \in \mathbb{K}[x_1, \dots, x_n, y]$  such that

$$\sum_{i=1}^s h_i g_i + h(f \cdot w - 1) = 1.$$

Replacing  $w$  by  $1/f(z)$  in the above equality we get

$$\sum_{i=1}^s h_i \left( z, \frac{1}{f(z)} \right) g_i(z) = 1. \tag{1.4.2}$$

Let  $d \gg 0$ : multiplying both sides of (1.4.2) by  $f^d$  we get that

$$\sum_{i=1}^s \bar{h}_i(z) g_i(z) = f^d(z), \quad \bar{h}_i \in \mathbb{K}[z_1, \dots, z_n].$$

Thus  $f \in \sqrt{I}$ .  $\square$

*Example 1.4.4.* Let  $V(F) \subset \mathbb{P}^n$  be a hypersurface, and let  $F_1, \dots, F_r$  be the distinct prime factors of the decomposition of  $F$  into a products of primes (recall that  $\mathbb{K}[Z_0, \dots, Z_n]$  is a UFD, by Corollary A.2.2). The irreducible decomposition of  $V(F)$  is

$$V(F) = V(F_1) \cup \dots \cup V(F_r).$$

In fact, each  $V(F_i)$  is irreducible by Proposition 1.3.10. What is not obvious is that  $V(F_i) \not\subset V(F_j)$  if  $F_i, F_j$  are non associated primes. This follows from Hilbert's Nullstellensatz.

## 1.5 Regular maps

Let  $U \subset \mathbb{P}^n$  be a locally closed subset. Suppose that  $F_0, \dots, F_m \in \mathbb{K}[Z_0, \dots, Z_n]_d$  are homogeneous polynomials of the same degree, and that for all  $[Z] \in U$  we have  $(F_0(Z), \dots, F_m(Z)) \neq (0, \dots, 0)$ . Let  $[Z] \in U$ . Then  $[F_0(Z), \dots, F_m(Z)] \in \mathbb{P}^m$  and if  $\lambda \in \mathbb{K}^*$  we have

$$[F_0(\lambda Z), \dots, F_m(\lambda Z)] = [\lambda^d F_0(Z), \dots, \lambda^d F_m(Z)] = [F_0(Z), \dots, F_m(Z)].$$

Hence we may define

$$\begin{array}{ccc} U & \longrightarrow & \mathbb{P}^m \\ [Z] & \longrightarrow & [F_0(Z), \dots, F_m(Z)] \end{array} \tag{1.5.1}$$

Maps as above are the local models for regular maps between quasi projective varieties.

**Definition 1.5.1.** Let  $X \subset \mathbb{P}^n$  and  $Y \subset \mathbb{P}^m$  be locally closed subsets (hence  $X$  and  $Y$  are quasi projective varieties), and let  $\varphi: X \rightarrow Y$  be a map. Then  $\varphi$  is *regular at*  $a \in X$  if there exist an open  $U \subset X$  containing  $a$  such that the restriction of  $\varphi$  to  $U$  is described as in (1.5.1). (We assume that  $(F_0(Z), \dots, F_m(Z)) \neq (0, \dots, 0)$  for all  $[Z] \in U$ .) The map  $\varphi$  is *regular* if it is regular at each point of  $X$ .

*Remark 1.5.2.* Let  $\varphi: X \rightarrow Y$  be a map between quasi projective varieties. Suppose that  $Y = \bigcup_{i \in I} U_i$  is an open cover, that  $\varphi^{-1}U_i$  is open in  $X$  for each  $i \in I$  and that the restriction

$$\begin{array}{ccc} \varphi^{-1}(U_i) & \longrightarrow & U_i \\ x & \longmapsto & \varphi(x) \end{array}$$

is regular for each  $i \in I$ . Then  $\varphi$  is regular. In other words regularity of a map is a local notion.

**Proposition 1.5.3.** *A regular map of quasi projective varieties is Zariski continuous.*

*Proof.* Let  $X \subset \mathbb{P}^n$  and  $Y \subset \mathbb{P}^m$  be Zariski locally closed, and let  $f: X \rightarrow Y$  be a regular map. We must prove that if  $C \subset Y$  is Zariski closed, then  $f^{-1}(C)$  is Zariski closed in  $X$ . Let  $U \subset W$  be an open subset such that (1.5.1) holds. Let us show that  $\phi^{-1}(C) \cap U$  is closed in  $U$ . Since  $C$  is closed  $C = V(I) \cap Y$  where  $I \subset \mathbb{K}[T_0, \dots, T_m]$  is a homogeneous ideal. Thus

$$\phi^{-1}(C) \cap U = \{[Z] \in U \mid P(F_0(Z), \dots, F_m(Z)) = 0 \forall P \in I\}.$$

Since each  $P(F_0(Z), \dots, F_m(Z))$  is a homogeneous polynomial, we get that  $\phi^{-1}(C) \cap U$  is closed in  $U$ .

By definition of regular map  $X$  can be covered by Zariski open sets  $U_\alpha$  such that (1.5.1) holds with  $U$  replaced by  $U_\alpha$ . We have proved that  $C_\alpha := \phi^{-1}(C) \cap U_\alpha$  is closed in  $U_\alpha$  for all  $\alpha$ . It follows that  $\phi^{-1}(C)$  is closed. In fact let  $\overline{C}_\alpha \subset X$  be the closure of  $C_\alpha$  and  $D_\alpha := X \setminus U_\alpha$ . Since  $C_\alpha$  is closed in  $U_\alpha$  we have

$$\overline{C}_\alpha \cap U_\alpha = C_\alpha = \phi^{-1}(C) \cap U_\alpha. \tag{1.5.2}$$

Moreover  $D_\alpha$  is closed in  $X$  because  $U_\alpha$  is open. By (1.5.2) we have

$$\phi^{-1}(C) = \bigcap_{\alpha} (\overline{C}_\alpha \cup D_\alpha).$$

Thus  $\phi^{-1}(C)$  is an intersection of closed sets and hence is closed. □

It is convenient to unravel the condition of being regular for maps with domain a subset of an affine space or both domain and codomain subsets of an affine space.

*Example 1.5.4.* Let  $X \subset \mathbb{A}^n (= \mathbb{P}_{Z_0}^n)$  and  $Y \subset \mathbb{P}^m$  be locally closed subsets, and let  $\varphi: X \rightarrow Y$  be a map. Then  $\varphi$  is a regular map if and only if, given any  $a \in X$ , there exist  $f_0, \dots, f_m \in \mathbb{K}[z_1, \dots, z_n]$  (in general *not* homogeneous) such that on an open subset  $U \subset X$  containing  $a$  we have

$$\varphi(z) = [f_0(z), \dots, f_m(z)]. \tag{1.5.3}$$

(This includes the statement that  $V(f_1, \dots, f_m) \cap U = \emptyset$ .) In fact, if  $\varphi$  is regular there exist homogeneous  $F_0, \dots, F_m \in \mathbb{K}[Z_0, \dots, Z_n]_d$  such that  $\varphi([1, z]) = [F_0(1, z), \dots, F_m(1, z)]$ , and it suffices to let  $f_j(z) := F_j(1, z)$ . Conversely, if (1.5.3) holds, then

$$\varphi([Z_0, Z_1, \dots, Z_n]) = [Z_0^d, Z_0^d f_1 \left( \frac{Z_1}{Z_0}, \dots, \frac{Z_n}{Z_0} \right), \dots, Z_0^d f_m \left( \frac{Z_1}{Z_0}, \dots, \frac{Z_n}{Z_0} \right)], \tag{1.5.4}$$

and for  $d$  is large enough, each of the rational functions appearing in (1.5.4) is actually a homogeneous polynomial of degree  $d$ .

*Example 1.5.5.* Let  $X \subset \mathbb{A}^n$  and  $Y \subset \mathbb{A}^m$  be locally closed subsets and let  $\varphi: X \rightarrow Y$  be a map. Recall that  $\mathbb{A}^n = \mathbb{P}_{Z_0}^n$  and  $\mathbb{A}^m = \mathbb{P}_{T_0}^m$ . Then  $\varphi$  is regular if and only if locally there exist  $f_0, \dots, f_m \in \mathbb{K}[z_1, \dots, z_n]$  (in general *not* homogeneous) such that

$$f(z) = \left( \frac{f_1(z)}{f_0(z)}, \dots, \frac{f_m(z)}{f_0(z)} \right). \quad (1.5.5)$$

Here it is understood that  $f_0(z) \neq 0$  for all  $z$  in the relevant open subset  $U$  of  $X$ . In fact this follows from (1.5.3) if we divide the homogeneous coordinates of  $\varphi(z)$  by  $f_0(z)$  (by hypothesis it does not vanish for  $z \in U$ ).

The identity map of a quasi projective variety is regular (choose  $F_j(Z) = Z_j$ ). If  $\varphi: X \rightarrow Y$  and  $\psi: Y \rightarrow W$  are regular maps of quasi projective varieties, the composition  $\psi \circ \varphi: X \rightarrow W$  is regular because the composition of homogeneous polynomial functions is a homogeneous polynomial function. Thus we have the *category of quasi projective varieties*. In particular we have the notion of isomorphism between quasi projective varieties.

**Definition 1.5.6.** A quasi projective variety is

- an *affine variety* if it is isomorphic to a closed subset of an affine space (as usual  $\mathbb{A}^n = \mathbb{P}_{Z_0}^n \subset \mathbb{P}^n$ ),
- a *projective variety* if it is isomorphic to a closed subset of a projective space.

*Remark 1.5.7.* Let  $X$  be an affine variety. If  $Y \subset X$  is closed then it is an affine variety. In fact by hypothesis there exist a closed subset  $W \subset \mathbb{A}^n$  and an isomorphism  $\varphi: X \xrightarrow{\sim} W$ . Since  $\varphi$  is an isomorphism it is a homeomorphism (see Proposition 1.5.3), and hence  $\varphi(Y)$  is a closed subset of  $W$ . Since  $W$  is closed in  $\mathbb{A}^n$ , it follows that  $\varphi(Y)$  is a closed subset of  $\mathbb{A}^n$ . The isomorphism  $Y \xrightarrow{\sim} \varphi(Y)$  shows that  $Y$  is an affine variety. Similarly one shows that if  $X$  is a projective variety and  $Y \subset X$  is closed, then  $Y$  is a projective variety.

The example below gives open (and non closed) subsets of an affine space which are affine varieties.

*Example 1.5.8.* Let  $f \in \mathbb{K}[z_1, \dots, z_n]$ . We let

$$\mathbb{A}_f^n := \mathbb{A}^n \setminus V(f). \quad (1.5.6)$$

Let  $Y := V(f(z_1, \dots, z_n) \cdot w - 1) \subset \mathbb{A}^{n+1}$ . The regular map

$$\begin{array}{ccc} \mathbb{A}_f^n & \xrightarrow{\varphi} & Y \\ (z_1, \dots, z_n) & \mapsto & (z_1, \dots, z_n, \frac{1}{f(z_1, \dots, z_n)}) \end{array}$$

is an isomorphism. In fact the inverse of  $\varphi$  is given by

$$\begin{array}{ccc} Y & \xrightarrow{\psi} & \mathbb{A}_f^n \\ (z_1, \dots, z_n, w) & \mapsto & (z_1, \dots, z_n) \end{array}$$

*Example 1.5.9.* Let

$$\mathcal{C}_d = \left\{ [\xi_0, \dots, \xi_d] \in \mathbb{P}^d \mid \text{rk} \begin{pmatrix} \xi_0 & \xi_1 & \cdots & \xi_{d-1} \\ \xi_1 & \xi_2 & \cdots & \xi_d \end{pmatrix} \leq 1 \right\}. \quad (1.5.7)$$

Since a matrix has rank at most 1 if and only if all the determinants of its  $2 \times 2$  minors vanish it follows that  $\mathcal{C}_d$  is closed. We have a regular map

$$\begin{array}{ccc} \mathbb{P}^1 & \xrightarrow{\varphi_d} & \mathcal{C}_d \\ [s, t] & \mapsto & [s^d, s^{d-1}t, \dots, t^d] \end{array} \quad (1.5.8)$$

Let us prove that  $\varphi_d$  is an isomorphism. Let  $\psi_d: \mathcal{C}_d \rightarrow \mathbb{P}^1$  be defined as follows:

$$\psi_d([\xi_0, \dots, \xi_d]) = \begin{cases} [\xi_0, \xi_1] & \text{if } [\xi_0, \dots, \xi_d] \in \mathcal{C}_d \cap \mathbb{P}_{\xi_0}^d \\ [\xi_{d-1}, \xi_d] & \text{if } [\xi_0, \dots, \xi_d] \in \mathcal{C}_d \cap \mathbb{P}_{\xi_d}^d \end{cases}$$

Of course in order for this to make sense one has to check the following:

1. The subset  $\mathcal{C}_d$  is the union of the open subsets  $\mathcal{C}_d \cap \mathbb{P}_{\xi_0}^d$  and  $\mathcal{C}_d \cap \mathbb{P}_{\xi_d}^d$ .
2. The two expressions for  $\psi_d$  coincide for points in  $\mathcal{C}_d \cap \mathbb{P}_{\xi_0}^d \cap \mathbb{P}_{\xi_d}^d$ .

To prove (1) suppose that  $[\xi] \in \mathcal{C}_d$  and  $\xi_0 = 0$ . By the equations defining  $\mathcal{C}_d$  it follows that  $\xi_1 = 0$ ,  $\xi_2 = 0$ , etc. up to  $\dots = \xi_{d-1}$ . Hence if  $\xi_0 = 0$  then  $\xi_d \neq 0$ , and this prove that Item (1) holds. To prove Item (2) suppose that  $[\xi] \in \mathcal{C}_d \cap \mathbb{P}_{\xi_0}^d \cap \mathbb{P}_{\xi_d}^d$ . By the equations defining  $\mathcal{C}_d$  it follows that  $\xi_0 \cdot \xi_n - \xi_1 \xi_{n-1} = 0$  and hence  $[\xi_0, \xi_1] = [\xi_{d-1}, \xi_d]$ . This prove that Item (2) holds.

One checks easily that  $\psi_d \circ \varphi_d = \text{Id}_{\mathbb{P}^1}$  and  $\varphi_d \circ \psi_d = \text{Id}_{\mathcal{C}_d}$ . Thus  $\varphi_d$  is an isomorphism, as claimed.

**Definition 1.5.10.** The closed subset  $\mathcal{C}_d \subset \mathbb{P}^d$  defined in (1.5.7) or any  $X \subset \mathbb{P}^d$  projectively equivalent to  $\mathcal{C}_d$  (i.e. given by  $g(\mathcal{C}_d)$  where  $g \in \text{PGL}_n(\mathbb{K})$ ) is a *rational normal curve in  $\mathbb{P}^d$* .

In the above definition “rational” refers to the fact that  $\mathcal{C}_d$  (and hence also any  $X$  projectively equivalent to  $\mathcal{C}_d$ ) is isomorphic to  $\mathbb{P}^1$ , “curve” refers to the fact that  $\mathbb{P}^1$  (and hence also  $\mathcal{C}_d$ ) has dimension 1 (we will define the dimension of a quasi projective variety later on), the attribute “normal” will be explained later in the book.

The remark below shows that, in the definition of regular map, we cannot require that  $\varphi$  is given globally by homogeneous polynomials.

*Remark 1.5.11.* Unless we are in the trivial case  $d = 1$ , it is *not* possible to define  $\psi_d$  globally as

$$\psi_d([\xi_0, \dots, \xi_d]) = [P(\xi_0, \dots, \xi_d), Q(\xi_0, \dots, \xi_d)], \quad (1.5.9)$$

with  $P, Q \in \mathbb{K}[\xi_0, \dots, \xi_d]_e$  not vanishing simultaneously on  $\mathcal{C}_d$ . In fact suppose that (1.5.9) holds, and let

$$p(s, t) := P(s^d, \dots, t^d), \quad q(s, t) := Q(s^d, \dots, t^d).$$

The polynomials  $p(s, t), q(s, t)$  are homogeneous of degree  $de$ , they do not vanish simultaneously on a non zero  $(s_0, t_0) \in \mathbb{K}^2$ , and for all  $[s, t] \in \mathbb{P}^1$  we have  $[p(s, t), q(s, t)] = [s, t]$ . The last equality means that  $tp(s, t) = sq(s, t)$ . It follows that  $p(s, t) = s \cdot r(s, t)$  and  $q(s, t) = t \cdot r(s, t)$  where  $r(s, t)$  has no non trivial zeroes. Thus  $r(s, t)$  is constant. In particular  $de = \deg p = \deg q = 1$ , and hence  $d = 1$ .

The example below extends Example 1.5.9 to arbitrary dimension.

*Example 1.5.12.* We recall the formula

$$\dim \mathbb{K}[Z_0, \dots, Z_n]_d = \binom{d+n}{n}. \quad (1.5.10)$$

(See Exercise 1.9.9 for a proof.) Let  $N(n; d) := \binom{d+n}{n} - 1$ . Let

$$\begin{array}{ccc} \mathbb{P}^n & \xrightarrow{\nu_d^n} & \mathbb{P}^{N(n; d)} \\ [Z] & \mapsto & [Z_0^d, Z_0^{d-1}Z_1, \dots, Z_n^d] \end{array} \quad (1.5.11)$$

be defined by all homogeneous monomials of degree  $d$  - this is a *Veronese map*. Clearly  $\nu_d^n$  is regular. Note that for  $n = 1$  we get back the map  $\varphi_d$  in (1.5.8).

The homogeneous coordinates on  $\mathbb{P}^{N(n; d)}$  appearing in (1.5.11) are indexed by length  $n + 1$  multiindices  $I = (i_0, \dots, i_n) \in \mathbb{N}^{n+1}$  such that  $\deg I := i_0 + \dots + i_n = d$ ; we denote them by  $[\dots, \xi_I, \dots]$ . Let  $\mathcal{V}_d^n \subset \mathbb{P}^{N(n; d)}$  be the closed subset defined by

$$\mathcal{V}_d^n := V(\dots, \xi_I \cdot \xi_J - \xi_K \cdot \xi_L, \dots),$$

where  $I, J, L, K$  run through all multiindices such that  $I + J = K + L$ . Clearly  $\nu_d^n(\mathbb{P}^n) \subset \mathcal{V}_d^n$ . Let us show that  $\nu_d^n$  is an isomorphism onto  $\mathcal{V}_d^n$ .



Let  $s \in \{0, \dots, n\}$ , and let  $H \in \mathbb{N}^{n+1}$  be a multiindex of degree  $(d-1)$ . We let  $e_s \in \mathbb{N}^{n+1}$  be the element all of whose entries are equal to 0 except for the entry at place  $s+1$ , which is equal to 1, and  $H_s := H + e_s$ . Also let

$$\begin{array}{ccc} \mathcal{V}_d^n \setminus V(\xi_{H_0}, \dots, \xi_{H_n}) & \xrightarrow{\varphi_d^n(H)} & \mathbb{P}^n \\ [\dots, \xi_I, \dots] & \mapsto & [\xi_{H_0}, \dots, \xi_{H_n}] \end{array}$$

Clearly  $\varphi_d^n(H)$  is regular. Moreover if  $[\dots, \xi_I, \dots] \in \mathcal{V}_d^n$  then there exist a multiindex  $H \in \mathbb{N}^{n+1}$  of degree  $(d-1)$  such that  $x$  belongs to  $\mathcal{V}_d^n \setminus V(\xi_{H_0}, \dots, \xi_{H_n})$  for  $H \in \mathbb{N}^{n+1}$  (there exists  $I \in \mathbb{N}^{n+1}$  of degree  $d$  such that  $\xi_I \neq 0$  and  $I = H + e_s$  where  $s$  is such that  $i_s \neq 0$ ). Moreover we claim that if  $[\dots, \xi_I, \dots] \in \mathcal{V}_d^n$  belong both to the domain of  $\varphi_d^n(H)$  and to the domain of  $\varphi_d^n(H')$ , then

$$\varphi_d^n(H)([\dots, \xi_I, \dots]) = [\xi_{H_0}, \dots, \xi_{H_n}] = [\xi_{H'_0}, \dots, \xi_{H'_n}] = \varphi_d^n(H')([z]). \quad (1.5.12)$$

In fact for  $s, t \in \{0, \dots, n\}$  we have  $H_s + H'_t = H + H' + e_s + e_t = H_t + H'_s$ , thus  $\xi_{H_s} \cdot \xi_{H'_t} - \xi_{H'_s} \cdot \xi_{H_t} = 0$  by the equations defining  $\mathcal{V}_d^n$ , and this proves that the equality in (1.5.12) holds. This shows that the maps  $\varphi_d^n(H)$ 's define a regular map

$$\mathcal{V}_d^n \xrightarrow{\varphi_d^n} \mathbb{P}^n. \quad (1.5.13)$$

We claim that

$$\varphi_d^n \circ \nu_d^n = \text{Id}_{\mathbb{P}^n} \quad (1.5.14)$$

$$\nu_d^n \circ \varphi_d^n = \text{Id}_{\mathcal{V}_d^n}. \quad (1.5.15)$$

The first equality is easily checked. In order to check the second equality it suffices to show that  $\nu_d^n$  is surjective. One may proceed as follows. Let  $x = [\dots, \xi_I, \dots] \in \mathcal{V}_d^n$  be a point such that  $\xi_{de_s} \neq 0$  for some  $s \in \{0, \dots, n\}$ . Thus  $x \in (\mathcal{V}_d^n \setminus V(\xi_{H_0}, \dots, \xi_{H_n}))$  where  $H = (d-1)e_0$ . It is not difficult to show that  $x = \nu_d^n([\xi_{H_0}, \dots, \xi_{H_n}])$ . Hence it suffices to prove that if  $x = [\dots, \xi_I, \dots] \in \mathcal{V}_d^n$ , then there exists  $s \in \{0, \dots, n\}$  such that  $\xi_{de_s} \neq 0$ . Equivalently, we must show that the following statement holds: if  $\xi := (\dots, \xi_I, \dots)$  is such that  $\xi_{de_s} = 0$  for all  $s \in \{0, \dots, n\}$  and  $\xi_I \cdot \xi_J = \xi_K \cdot \xi_L$  whenever  $I + J = K + L$ , then  $\xi_I = 0$  for all multiindices  $I$ . This is easily proved by “descending induction” on the maximum of  $i_0, \dots, i_n$ . If the maximum is  $d$ , then  $\xi_I = 0$  by hypothesis. Suppose that the maximum is at least  $d/2$ , i.e. that there exists  $s \in \{0, \dots, n\}$  be such that  $2i_s \geq d$ . Then  $2I = de_s + J$  where  $J \in \mathbb{N}^{n+1}$  is a multiindex of degree  $d$  and hence  $\xi_I^2 = \xi_{de_s} \cdot \xi_J = 0$  by the equations defining  $\mathcal{V}_d^n$ . Thus  $\xi_I = 0$ . This proves that if the maximum is at least  $d/2$  then  $\xi_I = 0$ . Iterating the argument we get that if the maximum is at least  $d/4$  then  $\xi_I = 0$  etc.

The Veronese map allows us to show that the open affine subsets of a quasi projective variety form a basis for the Zariski topology. First we need a definition.

**Definition 1.5.13.** Let  $X \subset \mathbb{P}^n$  be a closed subset. A *principal open subset* of  $X$  is an open  $U \subset X$  which is equal to

$$X_F := X \setminus V(F),$$

where  $F \in \mathbb{K}[Z_0, \dots, Z_n]$  is a homogeneous polynomial of *strictly positive degree*.

**Claim 1.5.14.** Let  $X \subset \mathbb{P}^n$  be closed. A *principal open subset* of  $X$  is an affine variety.

*Proof.* First we prove the claim for  $X = \mathbb{P}^n$ . Let  $F \in \mathbb{K}[Z_0, \dots, Z_n]$  be a homogeneous polynomial of strictly positive degree  $d$ . In order to prove that  $\mathbb{P}_F^n$  is affine we consider the Veronese map  $\nu_d^n: \mathbb{P}^n \rightarrow \mathbb{P}^{N(n,d)}$ , see (1.5.11). Let  $\mathcal{V}_d^n := \text{im}(\nu_d^n)$  be the corresponding Veronese variety. As shown in Example 1.5.12 the map  $\mathbb{P}^n \rightarrow \mathcal{V}_d^n$  defined by  $\nu_d^n$  is an isomorphism. Let  $F = \sum_I a_I Z^I$ , and let  $H \subset \mathbb{P}^{N(n,d)}$  be the hyperplane  $H = V(\sum_I a_I \xi_I)$ . Then we have the isomorphism

$$\begin{array}{ccc} \mathbb{P}_F^n & \xrightarrow{\sim} & (\mathcal{V}_d^n \setminus H) \\ x & \mapsto & \nu_d^n(x) \end{array} \quad (1.5.16)$$

But  $\mathbb{P}^{N(n,d)} \setminus H$  is the affine space  $\mathbb{A}^{N(n,d)}$ , and hence  $(\mathcal{V}_d^n \setminus H)$  is a closed subset of  $\mathbb{A}^{N(n,d)}$ . Hence the map in (1.5.16) is an isomorphism between  $\mathbb{P}_F^n$  and closed subset of  $\mathbb{A}^{N(n,d)}$ , and therefore  $\mathbb{P}_F^n$  is an affine variety.

In general, let  $X \subset \mathbb{P}^n$  be closed, and let  $F$  be as above. Then  $X_F$  is a closed subset of the affine variety  $\mathbb{P}_F^n$ , and hence it is an affine variety, see Remark `rmk:trapano`.  $\square$

**Proposition 1.5.15.** *The open affine subsets of a quasi projective variety form a basis of the Zariski topology.*

*Proof.* Since a quasi-projective variety is an open subset of a projective variety, it suffices to prove the result for projective varieties. Let  $X \subset \mathbb{P}^n$  be closed. Let  $U \subset X$  be open. If  $U = X$  then

$$U = X = X_{Z_0} \cup X_{Z_1} \cup \dots \cup X_{Z_n}, \quad (1.5.17)$$

and each of the  $X_{Z_i}$ 's is an open affine subset by Claim 1.5.14.

Next assume that  $U \neq X$ . Then  $U = X \setminus V(F_1, \dots, F_r)$ , where each  $F_j$  is a non constant homogeneous polynomial, and  $r \geq 1$ . Then

$$U = X_{F_1} \cup \dots \cup X_{F_r},$$

and each of the  $X_{F_j}$ 's is an open affine subset by Claim 1.5.14.  $\square$

## 1.6 Regular functions on affine varieties

**Definition 1.6.1.** A *regular function* on a quasi projective variety  $X$  is a regular map  $X \rightarrow \mathbb{K}$ .

Let  $X$  be a non empty quasi projective variety. The set of regular functions on  $X$  with pointwise addition and multiplication is a  $\mathbb{K}$ -algebra, named the *ring of regular functions* of  $X$ . We denote it by  $\mathbb{K}[X]$ .

If  $X$  is a projective variety, then it has few regular functions. In fact we will prove (see Corollary 2.4.8) that every regular function on  $X$  is locally constant. On the other hand, affine varieties have plenty of functions. In fact if  $X \subset \mathbb{A}^n$  is closed we have an inclusion

$$\mathbb{K}[z_1, \dots, z_n]/I(X) \hookrightarrow \mathbb{K}[X]. \quad (1.6.1)$$

**Theorem 1.6.2.** *Let  $X \subset \mathbb{A}^n$  be closed. Then the homomorphism in (1.6.1) is an isomorphism, i.e. every regular function on  $X$  is the restriction of a polynomial function on  $\mathbb{A}^n$ .*

Theorem 1.6.2 follows from the Nullstellensatz. Before giving the proof we discuss a particular instance of Theorem 1.6.2, which shows the relation with the Nullstellensatz. Let  $X \subset \mathbb{A}^n$  be closed. Suppose that  $g \in \mathbb{K}[z_1, \dots, z_n]$  and that  $g(a) \neq 0$  for all  $a \in X$ . Then  $1/g \in \mathbb{K}[X]$  and hence Theorem 1.6.2 predicts the existence of  $f \in \mathbb{K}[z_1, \dots, z_n]$  such that  $g^{-1} = f|_X$ . Such an  $f$  exists by the Nullstellensatz. In fact let  $X = V(g_1, \dots, g_r)$  where  $g_1, \dots, g_r \in \mathbb{K}[z_1, \dots, z_n]$ . By our hypothesis on  $g$  we have  $V(g_1, \dots, g_r, g) = \emptyset$ , and hence  $(g_1, \dots, g_r, g) = (1)$  by the Nullstellensatz. Hence there exist  $f_1, \dots, f_r, f \in \mathbb{K}[z_1, \dots, z_n]$  such that

$$f_1 \cdot g_1 + \dots + f_r \cdot g_r + f \cdot g = 1.$$

Restricting to  $X$  we get that  $f(x) = g(x)^{-1}$  for all  $x \in X$ , as claimed.

Before proving Theorem 1.6.2, we notice that, if  $X \subset \mathbb{A}^n$  is closed, the Nullstellensatz for  $\mathbb{K}[z_1, \dots, z_n]$  implies a Nullstellensatz for  $\mathbb{K}[z_1, \dots, z_n]/I(X)$ . First a definition: given an ideal  $J \subset (\mathbb{K}[z_1, \dots, z_n]/I(X))$  we let

$$V(J) := \{a \in X \mid f(a) = 0 \quad \forall f \in J\}.$$

The following result follows at once from the Nullstellensatz.

**Proposition 1.6.3** (Nullstellensatz for a closed subset of  $\mathbb{A}^n$ ). *Let  $X \subset \mathbb{A}^n$  be closed, and let  $J \subset (\mathbb{K}[z_1, \dots, z_n]/I(X))$  be an ideal. Then*

$$\{f \in (\mathbb{K}[z_1, \dots, z_n]/I(X)) \mid f|_{V(J)} = 0\} = \sqrt{J}.$$

(The radical  $\sqrt{J}$  is taken inside  $\mathbb{K}[z_1, \dots, z_n]/I(X)$ .) In particular  $V(J) = \emptyset$  if and only if  $J = (1)$ .

We introduce notation that is useful in the proof of Theorem 1.6.2. Given a quasi projective variety  $X$ , and  $f \in \mathbb{K}[X]$ , let

$$X_f := X \setminus V(f), \quad (1.6.2)$$

where  $V(f) := \{x \in X \mid f(x) = 0\}$ . Note the similarity with the notation for principal open subsets of projective varieties.

*Remark 1.6.4.* Assume that  $X$  is affine, hence we may assume that  $X \subset \mathbb{A}^n$  is closed. The collection of open subsets  $\{X_f\}$  is a basis for the Zariski topology of  $X$ . In fact let  $U$  be an open subset of  $X$ . Then  $U = X \setminus V(g_1, \dots, g_r)$  where  $g_i \in \mathbb{K}[z_1, \dots, z_n]$  for  $i \in \{1, \dots, r\}$ . Let  $f_i := g_i|_X$ . Then  $U = X_{g_1} \cup \dots \cup X_{g_r}$ .

*Proof of Theorem 1.6.2.* The proof is simpler if  $X$  is irreducible. We first give the proof under this hypothesis. Let  $\varphi \in \mathbb{K}[X]$ . We claim that there exist  $f_i, g_i \in \mathbb{K}[z_1, \dots, z_n]$  for  $1 \leq i \leq d$  with  $g_i \notin I(X)$  such that

$$(a) \quad X = \bigcup_{1 \leq i \leq d} X_{g_i}, \text{ i.e. } V(g_1, \dots, g_d) \cap X = \emptyset,$$

$$(b) \quad \text{for all } x \in X_{g_i} \text{ we have } \varphi(x) = \frac{f_i(x)}{g_i(x)},$$

In fact by definition of regular function (see Example 1.5.5) there exist an open cover  $X = \bigcup_{\alpha \in A} U_\alpha$  and  $f_\alpha, g_\alpha \in \mathbb{K}[z_1, \dots, z_n]$  for each  $\alpha \in A$  such that  $U_\alpha \subset X_{g_\alpha}$  and  $\varphi(x) = \frac{f_\alpha(x)}{g_\alpha(x)}$  for each  $x \in U_\alpha$ . Since the Zariski topology is quasi compact (see Corollary 1.3.9) we may assume that index set  $A$  is finite, say  $A = \{1, \dots, d\}$ . Of course we may assume that  $g_i \neq 0$  for all  $i \in \{1, \dots, d\}$ . Since  $X$  is irreducible so is  $X_{g_i}$  and hence  $U_i$  is dense in  $X_{g_i}$ . This implies that  $\varphi(x) = \frac{f_i(x)}{g_i(x)}$  on all of  $X_{g_i}$  because regular functions are Zariski continuous (see Proposition 1.5.3). This proves the claim.

In the rest of the proof we adopt the following notation: for  $f \in \mathbb{K}[z_1, \dots, z_n]$  we let  $\bar{f} := f|_X$ .

For  $i = 1, \dots, d$  the equality  $\bar{g}_i \varphi = \bar{f}_i$  holds on  $X_{g_i}$  by Item (2). Since  $X$  is irreducible and  $X_{g_i}$  is a non empty subset of  $X$  it is dense in  $X$ , and hence  $\bar{g}_i \varphi = \bar{f}_i$  on all of  $X$  (this is where the hypothesis that  $X$  is irreducible simplifies the proof). By Proposition 1.6.3 we have that  $(\bar{g}_1, \dots, \bar{g}_d) = (1)$ , i.e. there exist  $h_1, \dots, h_d \in \mathbb{K}[z_1, \dots, z_n]$  such that

$$1 = \bar{h}_1 \bar{g}_1 + \dots + \bar{h}_d \bar{g}_d.$$

where  $\bar{h}_i := h_i|_X$ . Multiplying by  $\varphi$  both sides of the above equality we get that

$$\varphi = \bar{h}_1 \bar{g}_1 \varphi + \dots + \bar{h}_d \bar{g}_d \varphi = \bar{h}_1 \bar{f}_1 + \dots + \bar{h}_d \bar{f}_d = (h_1 f_1 + \dots + h_d f_d)|_X. \quad (1.6.3)$$

This shows that  $\varphi$  is the restriction to  $X$  of a polynomial function on  $\mathbb{A}^n$ .

Now we give the proof for arbitrary (closed)  $X$ . Let  $\varphi \in \mathbb{K}[X]$ . This time we claim that there exist  $f_i, g_i \in \mathbb{K}[z_1, \dots, z_n]$  for  $i \in \{1, \dots, d\}$  such that

$$1. \quad X = \bigcup_{1 \leq i \leq d} X_{g_i}, \text{ i.e. } V(g_1, \dots, g_d) \cap X = \emptyset,$$

$$2. \quad \text{for all } a \in X_{g_i} \text{ we have } \varphi(a) = \frac{f_i(a)}{g_i(a)},$$

$$3. \quad \text{for } 1 \leq i \leq j \text{ we have } (g_j f_i - g_i f_j)|_X = 0.$$

We start proving the claim as in the case of  $X$  irreducible. There is a finite open cover  $X = \bigcup_{\alpha \in A} U_\alpha$  and  $f_\alpha, g_\alpha \in \mathbb{K}[z_1, \dots, z_n]$  for each  $\alpha \in A$  such that  $U_\alpha \subset X_{g_\alpha}$  and  $\varphi(x) = \frac{f_\alpha(x)}{g_\alpha(x)}$  for each  $x \in U_\alpha$ . We may cover  $U_\alpha$  by open affine sets  $X_{\gamma_{\alpha,1}}, \dots, X_{\gamma_{\alpha,r}}$ , see Remark 1.6.4. Since  $V(\bar{g}_\alpha) \subset \bigcap_{j=1}^r V(\bar{\gamma}_{\alpha,j})$  (recall that  $\bar{g}_\alpha$  and  $\bar{\gamma}_{\alpha,j}$  are the restrictions to  $X$  of  $g_\alpha$  and  $\gamma_{\alpha,j}$  respectively), the Nullstellensatz for  $X$  gives that, for each  $\alpha, j$ , there exist  $N_{\alpha,j} > 0$  and  $\mu_{\alpha,j} \in \mathbb{K}[z_1, \dots, z_n]$  such that  $\bar{\gamma}_{\alpha,j}^{N_{\alpha,j}} = \bar{\mu}_{\alpha,j} \cdot \bar{g}_\alpha$ . Hence  $\varphi(x) = \mu_{\alpha,j}(x)f_\alpha(x)/\gamma_{\alpha,j}(x)^{N_{\alpha,j}}$  for all  $x \in X_{\gamma_{\alpha,j}}$ . Since  $V(\gamma_{\alpha,j}) = V(\gamma_{\alpha,j}^{N_{\alpha,j}})$  it follows that there exist  $f'_i, g'_i \in \mathbb{K}[z_1, \dots, z_n]$  for  $i \in \{1, \dots, d\}$  such that  $X = \bigcup_{i=1}^d X_{g'_i}$  and  $\varphi(x) = f'_i(x)/g'_i(x)$  for all  $x \in X_{g'_i}$ . For  $i \in \{1, \dots, d\}$  let

$$f_i := f'_i g'_i, \quad g_i := (g'_i)^2.$$

Clearly Items (1) and (2) hold. In order to check Item (3) we write

$$(g_j f_i - g_i f_j)|_X = ((g'_j)^2 f'_j g'_i - (g'_i)^2 f'_i g'_j)|_X = ((g'_i g'_j)(f'_i g'_j - f'_j g'_i))|_X.$$

Since  $\varphi(z) = f'_i(z)/g'_i(z) = f'_j(z)/g'_j(z)$  for all  $x \in X_{g'_i} \cap X_{g'_j}$  the last term vanishes on  $X_{g'_i} \cap X_{g'_j}$ . On the other hand the last term vanishes also on  $(X \setminus X_{g'_i} \cap X_{g'_j}) = X \cap V(g'_i g'_j)$  because of the factor  $(g'_i g'_j)$ . This finishes the proof that there exist  $f_i, g_i \in \mathbb{K}[z_1, \dots, z_n]$  for  $i \in \{1, \dots, d\}$  such that (1), (2) and (3) hold.

Next, for  $i = 1, \dots, d$  let  $\bar{g}_i := g_i|_X$  and  $\bar{f}_i := f_i|_X$ . Then

$$\bar{g}_i \varphi = \bar{f}_i. \tag{1.6.4}$$

In fact by Item (1) it suffices to check that (1.6.4) holds on  $X_{g_j}$  for  $j = 1, \dots, d$ . For  $j = i$  it holds by Item (2), for  $j \neq i$  it holds by Item (3). Given the equalities in (1.6.4), one finishes the proof proceeding as in the case when  $X$  is irreducible.  $\square$

*Example 1.6.5.* Let  $X$  be an affine variety, thus we may assume that  $X \subset \mathbb{A}^n$  is closed. If  $f \in \mathbb{K}[X]$  then  $X_f$  is a principal open subset of  $\bar{X}$ . In fact by Theorem 1.6.2 there exists  $g \in \mathbb{K}[z_1, \dots, z_n]$  such that  $f = g|_X$ . If  $d \gg 0$  then

$$G(Z_0, \dots, Z_n) := Z_0^d g\left(\frac{Z_1}{Z_0}, \dots, \frac{Z_n}{Z_0}\right)$$

is a homogeneous polynomial whose zero locus (in  $\mathbb{P}^n$ ) is equal to the union of  $V(Z_0)$  and  $V(g)$  (which is contained in  $\mathbb{A}^n$ ). Hence  $\bar{X}_G = (\bar{X} \setminus V(G)) = (X \setminus V(g)) = X_f$ . An explicit isomorphism between  $X_f$  and a closed subset of an affine space is obtained as follows. Let  $Y := V(J) \subset \mathbb{A}^{n+1}$  where  $J$  is the ideal generated by  $I(X)$  and the polynomial  $g(z_1, \dots, z_n) \cdot z_{n+1} - 1$ . Then the map

$$\begin{aligned} X_f &\longrightarrow Y \\ (z_1, \dots, z_n) &\longmapsto \left(z_1, \dots, z_n, \frac{1}{g(z_1, \dots, z_n)}\right) \end{aligned}$$

is an isomorphism (see Example 1.5.8). Note that by Theorem 1.6.2 every regular function on  $X_f$  is given by the restriction to  $X_f$  of  $\frac{h}{f^m}$ , where  $h \in \mathbb{K}[X]$  and  $m \in \mathbb{N}$ .

## 1.7 Quasi-projective varieties defined over a subfield of $\mathbb{K}$

Let  $F \subset \mathbb{K}$  be a subfield, for example  $\mathbb{R} \subset \mathbb{C}$ ,  $\mathbb{Q} \subset \mathbb{C}$  or  $\mathbb{F}_q \subset \bar{\mathbb{F}}_q$  where  $q = p^r$  with  $p$  a prime.

**Definition 1.7.1.** A locally closed subset  $X \subset \mathbb{P}^n(\mathbb{K})$  is *defined over  $F$*  if both the homogeneous ideals  $I(\bar{X}) \subset \mathbb{K}[Z_0, \dots, Z_n]$  and  $I(\bar{X} \setminus X) \subset \mathbb{K}[Z_0, \dots, Z_n]$  admit sets of generators belonging to  $F[Z_0, \dots, Z_n]$ .

Trivially  $\mathbb{P}^n(\mathbb{K})$  and  $\mathbb{A}^n(\mathbb{K}) = \mathbb{P}^n(\mathbb{K})_{Z_0}$  are defined over the prime field, i.e. over  $\mathbb{Q}$  if  $\text{char } \mathbb{K} = 0$  and over  $\mathbb{F}_p$  if  $\text{char } \mathbb{K} = p$ .

*Remark 1.7.2.* A locally closed subset  $X \subset \mathbb{A}^n(\mathbb{K}) = \mathbb{P}^n(\mathbb{K})_{Z_0}$  is defined over  $F$  if both the ideals  $I(\overline{X}) \subset \mathbb{K}[z_1, \dots, z_n]$  and  $I(\overline{X} \setminus X) \subset \mathbb{K}[z_1, \dots, z_n]$  (in general non homogeneous) admit sets of generators which belong to  $F[z_1, \dots, z_n]$ . This is so because a polynomial  $p \in \mathbb{K}[z_1, \dots, z_n]$  of degree  $d$  vanishes on  $X$  if and only if the homogeneous polynomial  $P := Z_0^d \cdot f(Z_1/Z_0, \dots, Z_n/Z_0)$  vanishes on  $\overline{X}$ , and conversely a homogeneous  $P \in \mathbb{K}[Z_0, \dots, Z_n]$  vanishes on  $\overline{X}$  if and only if  $P(1, z_1, \dots, z_n) \in \mathbb{K}[z_1, \dots, z_n]$  vanishes on  $X$ .

*Example 1.7.3.* Let  $a = (a_1, \dots, a_n) \in \mathbb{A}^n$ . If  $a_i$  belongs to  $F$  for all  $i \in \{1, \dots, n\}$  then  $\{a\}$  is defined over  $F$  because its ideal is generated by  $(z_1 - a_1, \dots, z_n - a_n)$ . The converse is true if we make a hypothesis on the field extension  $F \subset \mathbb{K}$ . Let  $\text{Aut}(\mathbb{K}, F)$  be the group of automorphisms of  $\mathbb{K}$  fixing every element of  $F$ . Assume that the field of elements of  $\mathbb{K}$  fixed by  $\text{Aut}(\mathbb{K}, F)$  is equal to  $F$ . (Since  $\mathbb{K}$  is algebraically closed this holds if  $\text{char } \mathbb{K} = 0$  or, in case  $\text{char } \mathbb{K} = p$  if  $F$  is perfect, i.e. every element of  $F$  has a  $p$ -th root in  $F$  (necessarily unique).) With this hypothesis, suppose that  $\{a\}$  is defined over  $F$ , and let  $p_1, \dots, p_r \in F[z_1, \dots, z_n]$  be generators of  $I(\{a\}) \subset \mathbb{K}[z_1, \dots, z_n]$ . For  $j \in \{1, \dots, r\}$  let  $p_j = \sum_I c_{j,I} z^I$  where  $c_{j,I} \in F$  for each multiindex  $I$ . If  $\sigma \in \text{Aut}(\mathbb{K}, F)$  we have

$$0 = \sigma(0) = \sigma(p_j(a)) = p_j(\sigma(a_1), \dots, \sigma(a_n)) = \sum_I c_{j,I} \sigma(a_1)^{i_1} \dots \sigma(a_n)^{i_n} = p_j(\sigma(a)). \quad (1.7.1)$$

(The third equality holds because  $p_j$  has coefficients in  $F$ .) Since the above equality holds for generators of the ideal of  $\{a\}$ , we get that  $(\sigma(a_1), \dots, \sigma(a_n)) = (a_1, \dots, a_n)$  for all  $\sigma \in \text{Aut}(\mathbb{K}, F)$ . By our hypothesis on  $\text{Aut}(\mathbb{K}, F)$  it follows that  $a_i \in F$  for all  $i$ .

*Example 1.7.4.* Let  $Q \in \mathbb{R}[Z_0, \dots, Z_n]_2$  be a non zero quadratic form. Then  $Z := V(Q) \subset \mathbb{P}^n(\mathbb{C})$  is a projective variety defined over  $\mathbb{R}$ . In fact if  $Q$  has rank at least 2 then  $Q$  generates  $I(Z)$ , and if  $Q$  has rank 1, i.e.  $Q = L^2$  for  $L \in \mathbb{C}[Z_0, \dots, Z_n]_1$  then either  $L \in \mathbb{R}[Z_0, \dots, Z_n]_1$  or  $\sqrt{-1}L \in \mathbb{R}[Z_0, \dots, Z_n]_1$ .

*Example 1.7.5.* The Fermat hypersurface  $X := V(\sum_{i=0}^n Z_i^d)$  is defined over the prime field. In order to check this one must show that  $I(X)$ , i.e. the radical of  $(\sum_{i=0}^n Z_i^d)$  is generated by a polynomial with coefficients in the prime field. If  $\text{char } \mathbb{K}$  does not divide  $d$  then the polynomial  $\sum_{i=0}^n Z_i^d$  generates a radical ideal in  $\mathbb{K}[Z_0, \dots, Z_n]$  (to see this take the formal partial derivative with respect to one of its variables), and hence it generates  $I(X)$ . Since the coefficients of  $\sum_{i=0}^n Z_i^d$  belong to the prime field we are done. If  $\text{char } \mathbb{K} = p > 0$  write  $d = p^r d_0$  where  $p$  does not divide  $d_0$ . Then  $\sum_{i=0}^n Z_i^d = (\sum_{i=0}^n Z_i^{d_0})^{p^r}$  and hence  $I(X)$  is generated by  $\sum_{i=0}^n Z_i^{d_0}$  (see above). Since the coefficients of  $\sum_{i=0}^n Z_i^{d_0}$  belong to the prime field we are done.

*Remark 1.7.6.* Let  $F \subset F' \subset \mathbb{K}$  be an inclusion of fields, and let  $X \subset \mathbb{P}^n(\mathbb{K})$  be a locally closed subset defined over  $F$ . Then  $X$  is also defined over  $F'$ . In particular if  $X$  is defined over the prime field it is defined over every subfield of  $\mathbb{K}$ .

**Definition 1.7.7.** Let  $X \subset \mathbb{P}^n(\mathbb{K})$  be a locally closed subset defined over  $F$ . We let  $X(F) \subset X$  be the set of points represented by  $(n+1)$ -tuples  $(Z_0, Z_1, \dots, Z_n) \in F^{n+1} \setminus \{(0, \dots, 0)\}$ .

*Remark 1.7.8.* Let  $X \subset \mathbb{A}^n(\mathbb{K})$  be a locally closed subset defined over  $F$ . Then  $X(F) \subset X$  is equal to  $X \cap \mathbb{A}^n(F)$ .

*Remark 1.7.9.* Let  $F \subset F' \subset \mathbb{K}$  be an inclusion of fields, and let  $X \subset \mathbb{P}^n(\mathbb{K})$  be a locally closed subset defined over  $F$ . Then  $X$  is also defined over  $F'$  and hence  $X(F')$  is also defined. In particular  $X(\mathbb{K})$  is defined and equals  $X$ .

*Remark 1.7.10.* Let  $p$  be a prime, and suppose that  $\mathbb{F}_q \subset \mathbb{K}$  where  $q = p^r$ . Let  $X \subset \mathbb{P}^n(\overline{\mathbb{F}}_p)$  be a locally closed subset defined over  $F_q$ . For each  $m \in \mathbb{N}_+$  there is a unique inclusion  $\mathbb{F}_q \subset \mathbb{F}_{q^m} \subset \mathbb{K}$ , and hence we have  $X(\mathbb{F}_{q^m})$ . Clearly  $X(\mathbb{F}_{q^m})$  is a finite set.

**Definition 1.7.11.** Let  $X \subset \mathbb{P}^n(\overline{\mathbb{F}}_p)$  be a locally closed subset defined over  $F_q$ , where  $q = p^r$ . The *Weil Zeta function* of  $X$  is defined to be formal power series in the variable  $T$  given by

$$Z(X, T) := \exp \left( \sum_{m=1}^{\infty} \frac{|X(\mathbb{F}_{q^m})|}{m} T^m \right) \quad (1.7.2)$$

**Definition 1.7.12.** Let  $X \subset \mathbb{P}^n(\mathbb{K})$  and  $Y \subset \mathbb{P}^m(\mathbb{K})$  be locally closed subset, both defined over a subfield  $F \subset \mathbb{K}$ . A map  $\varphi := X \rightarrow Y$  is *defined over  $F$*  if for each  $a \in X$  there exist an open  $U \subset X$  containing  $a$  and  $P_j \in F[Z_0, \dots, Z_n]_d$  for  $j \in \{0, \dots, m\}$  ( $d$  depends on  $U$ ), such that the restriction of  $\varphi$  to  $U$  is

$$\begin{array}{ccc} U & \longrightarrow & \mathbb{P}^m \\ [Z] & \longmapsto & [P_0(Z), \dots, P_m(Z)] \end{array} \quad (1.7.3)$$

(of course  $(P_0(Z), \dots, P_m(Z)) \neq (0, \dots, 0)$  for all  $[Z] \in U$ ).

Let  $F \subset \mathbb{K}$  be a subfield. If  $X \subset \mathbb{P}^n(\mathbb{K})$  is a locally closed subset defined over  $F$  then the identity map  $\text{Id}_X: X \rightarrow X$  is clearly defined over  $F$ . If  $X \subset \mathbb{P}^n(\mathbb{K})$ , and  $Y \subset \mathbb{P}^m(\mathbb{K})$ ,  $W \subset \mathbb{P}^l(\mathbb{K})$  are locally closed subsets defined over  $F$  and  $\varphi: X \rightarrow Y$ ,  $\psi: Y \rightarrow W$  are regular maps defined over  $F$  then the composition  $\psi \circ \varphi: X \rightarrow W$  is also defined over  $F$ . In fact this holds because if  $P \in F[Z_0, \dots, Z_m]_d$  and  $Q_0, \dots, Q_m \in F[T_0, \dots, T_n]_e$  then  $P(Q_0, \dots, Q_m) \in F[T_0, \dots, T_n]_{de}$ .

Hence we have the category of quasi projective varieties defined over  $F$ . In particular we have the notion of isomorphism over  $F$  of varieties defined over  $F$ .

*Remark 1.7.13.* Let  $X \subset \mathbb{P}^n(\mathbb{K})$  and  $Y \subset \mathbb{P}^m(\mathbb{K})$  be locally closed subsets defined over  $F$ . If  $\varphi: X \rightarrow Y$  is a regular map defined over  $F$  then  $\varphi(X(F)) \subset Y(F)$  because the value of a polynomial with coefficients in  $F$  at  $(A_0, \dots, A_n) \in F^{n+1}$  belongs to  $F$ .

*Example 1.7.14.* Let  $Q_1, Q_2 \in \mathbb{R}[Z_0, \dots, Z_n]_2$  be non degenerate quadratic forms, and let  $X_i := V(Q_i)$  for  $i \in \{1, 2\}$ . Then  $X_i \subset \mathbb{P}^n(\mathbb{C})$  is a projective variety defined over  $\mathbb{R}$ . Since  $Q_i$  is diagonalizable in suitable coordinates, there exists a projectivity  $\varphi: \mathbb{P}^n(\mathbb{C}) \rightarrow \mathbb{P}^n(\mathbb{C})$  whose restriction to  $X_1$  defines an isomorphism  $X_1 \xrightarrow{\sim} X_2$ . In particular  $X_1$  is isomorphic to  $X_2$  (over  $\mathbb{C}$ ). On the other hand  $X_1$  is not necessarily isomorphic to  $X_2$  over  $\mathbb{R}$ . In fact let  $Q_1 := \sum_{j=0}^n Z_j^2$  and  $Q_2 := Z_0^2 - \sum_{j=1}^n Z_j^2$ . Thus  $X_1(\mathbb{R})$  is empty while  $X_2(\mathbb{R})$  is not empty. Since a regular map  $\varphi: X_1 \rightarrow X_2$  defined over  $\mathbb{R}$  maps  $X_1(\mathbb{R})$  to  $X_2(\mathbb{R})$  it follows that  $X_1$  is not isomorphic to  $X_2$  over  $\mathbb{R}$  (we assume that  $n \geq 1$ ).

Under a suitable hypothesis we can avoid computing the radical of ideals if we wish to decide whether a locally closed subset  $X \subset \mathbb{P}^n(\mathbb{K})$  is defined over a subfield  $F \subset \mathbb{K}$ . Let  $\text{Aut}(\mathbb{K}/F)$  be the group of automorphisms of  $\mathbb{K}$  which are the identity on  $F$ .

**Proposition 1.7.15.** *Suppose that the fixed field of  $\text{Aut}(\mathbb{K}/F)$  is equal to  $F$ . Let  $X \subset \mathbb{P}^n(\mathbb{K})$  be a locally closed subset given by  $V(I) \setminus V(J)$  where  $I, J \subset \mathbb{K}[Z_0, \dots, Z_n]$  are homogeneous ideals which admit sets of generators belonging to  $F[Z_0, \dots, Z_n]$ . Then  $X$  is defined over  $F$ .*

Before proving Proposition 1.7.15 we go through a few preliminaries. The group  $\text{Aut}(\mathbb{K})$  of field automorphisms of  $\mathbb{K}$  acts on  $\mathbb{P}^n$  as follows: for  $\sigma \in \text{Aut}(\mathbb{K})$

$$\begin{array}{ccc} \text{Aut}(\mathbb{K}) \times \mathbb{P}^n & \longrightarrow & \mathbb{P}^n \\ (\sigma, [Z_0, \dots, Z_n]) & \longmapsto & [\sigma(Z_0), \dots, \sigma(Z_n)] \end{array} \quad (1.7.4)$$

Note that if  $X \subset \mathbb{A}^n (= \mathbb{P}_{Z_0}^n)$  then  $\sigma(z_1, \dots, z_n) = (\sigma(z_1), \dots, \sigma(z_n))$ .

*Remark 1.7.16.* In general the map  $\mathbb{P}^n \rightarrow \mathbb{P}^n$  that one gets by fixing a non trivial  $\sigma \in \text{Aut}(\mathbb{K})$  in (1.7.4) is not regular. For example if  $F = \mathbb{R} \subset \mathbb{C}$  and  $\sigma$  is complex conjugation the map is not regular.

**Proposition 1.7.17.** *Let  $X \subset \mathbb{P}^n(\mathbb{K})$  be a locally closed subset defined over  $F$ . If  $\sigma \in \text{Aut}(\mathbb{K}/F)$  then  $\sigma(X) = X$ .*

*Proof.* It suffices to prove that  $\sigma(X) = X$  for every closed subset  $X \subset \mathbb{P}^n(\mathbb{K})$  defined over  $F$ . Let  $P \in F[Z_0, \dots, Z_n] \cap I(X)$  be homogeneous. Thus  $P = \sum_I c_I Z^I$  where each  $c_I$  belongs to  $F$ . If  $[A_0, \dots, A_n] \in X$  then  $P(A_0, \dots, A_n) = 0$  and hence

$$0 = \sigma(P(A_0, \dots, A_n)) = \sum_I \sigma(c_I) \sigma(A_0)^{i_0} \dots \sigma(A_n)^{i_n} = \sum_I c_I \sigma(A_0)^{i_0} \dots \sigma(A_n)^{i_n} = P(\sigma(A_0), \dots, \sigma(A_n)).$$

This proves that  $\sigma(X) \subset X$  because the ideal  $I(X) \subset \mathbb{K}[Z_0, \dots, Z_n]$  is generated by homogeneous elements of  $F[Z_0, \dots, Z_n] \cap I(X)$ . Thus we also have  $\sigma^{-1}(X) \subset X$  and hence  $X \subset \sigma(X)$ .  $\square$

*Proof of Proposition 1.7.15.* The group  $\text{Aut}(\mathbb{K}/F)$  acts on  $\mathbb{K}[Z_0, \dots, Z_n]$  by acting on the coefficients of polynomials. We claim that  $\text{Aut}(\mathbb{K}/F)$  maps  $I(X)$  to itself. In fact let  $\sigma \in \text{Aut}(\mathbb{K}/F)$  and let  $P \in I(X)$  be a homogeneous polynomial,  $P = \sum_I c_I Z^I$ . By Proposition 1.7.17 we have  $\sigma^{-1}(X) = X$ , hence

$$\sigma(P)(A) = \sum_I \sigma(c_I) A^I = \sigma \left( \sum_I c_I \sigma^{-1}(A_0)^{i_0} \dots \sigma^{-1}(A_n)^{i_n} \right) = \sigma(P(\sigma^{-1}(A))) = 0.$$

We have an obvious isomorphism  $\mathbb{K}[Z_0, \dots, Z_n] \cong \mathbb{K}_F F[Z_0, \dots, Z_n]$  which is equivariant for the action of  $\text{Aut}(\mathbb{K}/F)$  that we have just defined and the action considered in Section A.6. By Proposition A.6.3 it follows that  $I(X)$  is generated (as  $\mathbb{K}$  vector space by its intersection with  $F[Z_0, \dots, Z_n]$ . This proves Proposition 1.7.15.  $\square$

*Example 1.7.18.* Assume that  $\text{char } \mathbb{K} = p > 0$ . Let  $F: \mathbb{K} \rightarrow \mathbb{K}$  be the Frobenius automorphism:  $F(a) := a^p$ . Let  $r$  be a positive natural number. Of course  $F^r$  is also an automorphism of  $\mathbb{K}$ . Note that  $F^r(a) = a^{p^r}$  and that  $F^r \in \text{Aut}(\mathbb{K}/\mathbb{F}_q)$ . There exists a unique embedding  $\mathbb{F}_q \subset \mathbb{K}$ . Suppose that  $X \subset \mathbb{P}^n$  is a locally closed subset defined over  $\mathbb{F}_q$ . Proposition 1.7.17 gives that we have the bijective map

$$\begin{array}{ccc} X & \xrightarrow{\pi} & X \\ [Z] & \mapsto & [Z_0^q, \dots, Z_n^q]. \end{array}$$

This is the *Frobenius map* of  $X$ . Note the exceptional feature of the Frobenius map: it is regular (see remark 1.7.16) and even defined over the prime field. Note also that  $X(\mathbb{F}_q)$  is equal to the fixed locus of  $\pi$ :

$$X(\mathbb{F}_q) = \text{Fix}(\pi). \tag{1.7.5}$$

## 1.8 Geometry and Algebra

Below is a remarkable consequences of Theorem 1.6.2.

**Proposition 1.8.1.** *Let  $R$  be a finitely generated  $\mathbb{K}$  algebra without nilpotents. There exists an affine variety  $X$  such that  $\mathbb{K}[X] \cong R$  (as  $\mathbb{K}$  algebras).*

*Proof.* Let  $\alpha_1, \dots, \alpha_n$  be generators (over  $\mathbb{K}$ ) of  $R$ , and let  $\varphi: \mathbb{K}[z_1, \dots, z_n] \rightarrow R$  be the surjection of algebras mapping  $z_i$  to  $\alpha_i$ . The kernel of  $\varphi$  is an ideal  $I \subset \mathbb{K}[z_1, \dots, z_n]$ , which is radical because  $R$  has no nilpotents. Let  $X := V(I) \subset \mathbb{A}^n$ . Then  $\mathbb{K}[X] \cong R$  by Theorem 1.6.2.  $\square$

The Nullstellensatz allows one to construct  $X$  abstractly from the  $\mathbb{K}$  algebra as follows. Let

$$\text{Spec}_m(R) := \{ \mathfrak{m} \subset R \mid \mathfrak{m} \text{ is a maximal ideal of } R \}$$

be the *maximal spectrum* of  $R$ . Hilbert's Nullstellensatz gives a bijection

$$\begin{array}{ccc} X & \leftrightarrow & \text{Spec}_m(R) \\ p & \mapsto & \{ f \in R \mid f(p) = 0 \} \end{array}$$

Thus  $X$  may be identified with  $\text{Spec}_m(R)$ . Moreover  $f \in R$  defines a function  $\text{Spec}_m(R) \rightarrow \mathbb{K}$  by setting  $f(\mathfrak{m}) := f \pmod{\mathfrak{m}}$ . This makes sense because the composition

$$\mathbb{K} \longrightarrow R \longrightarrow R/\mathfrak{m} \tag{1.8.1}$$

is an isomorphism.

Actually we get a contravariant equivalence between the category of affine varieties over  $\mathbb{K}$  and that of finitely generated  $\mathbb{K}$ -algebras. First we give a definition.

**Definition 1.8.2.** Let  $\varphi: X \rightarrow Y$  be a regular map of non empty quasi projective varieties. The pull-back  $\varphi^*: \mathbb{K}[Y] \rightarrow \mathbb{K}[X]$  is the homomorphism of  $\mathbb{K}$  algebras defined by

$$\begin{array}{ccc} \mathbb{K}[Y] & \xrightarrow{\varphi^*} & \mathbb{K}[X] \\ f & \mapsto & f \circ \varphi \end{array} \quad (1.8.2)$$

**Proposition 1.8.3.** Let  $Y$  be an affine variety, and let  $X$  be a quasi projective variety. The map

$$\begin{array}{ccc} \{X \xrightarrow{\varphi} Y \mid \varphi \text{ regular}\} & \longrightarrow & \{\mathbb{K}[Y] \xrightarrow{\alpha} \mathbb{K}[X] \mid \alpha \text{ homom. of } \mathbb{K}\text{-algebras}\} \\ \varphi & \mapsto & \varphi^* \end{array} \quad (1.8.3)$$

is a bijection.

*Proof.* We may assume that  $Y \subset \mathbb{A}^n$  is closed; for  $i \in \{1, \dots, n\}$  let  $\bar{z}_i := z_{i|X}$ . Suppose that  $f, g: X \rightarrow Y$  are regular maps, and that  $f^* = g^*$ . Then  $f^*(\bar{z}_i) = g^*(\bar{z}_i)$  for  $i \in \{1, \dots, n\}$ , and hence  $f = g$ . This proves injectivity of the map in (1.8.3).

In order to prove surjectivity, let  $\alpha: \mathbb{K}[Y] \rightarrow \mathbb{K}[X]$  be a homomorphism of  $\mathbb{K}$  algebras. Let  $f_i := \alpha(\bar{z}_i)$ , and let  $\varphi: X \rightarrow \mathbb{A}^n$  be the regular map defined by  $\varphi(x) := (f_1(x), \dots, f_n(x))$  for  $x \in X$ . We claim that  $\varphi(x) \in Y$  for all  $x \in X$ . In fact, since  $Y$  is closed, it suffices to show that  $g(\varphi(x)) = 0$  for all  $g \in I(X)$ . Now

$$g(\varphi(x)) = g(f_1(x), \dots, f_n(x)) = g(\alpha(\bar{z}_1), \dots, \alpha(\bar{z}_n)) = \alpha(g(\bar{z}_1), \dots, \bar{z}_n) = \alpha(0) = 0.$$

(The third equality holds because  $\alpha$  is a homomorphism of  $\mathbb{K}$ -algebras.) Thus  $\varphi$  is a regular map  $f: X \rightarrow Y$  such that  $\varphi^*(\bar{z}_i) = \alpha(\bar{z}_i)$  for  $i \in \{1, \dots, n\}$ . By Theorem 1.6.2 the  $\mathbb{K}$ -algebra  $\mathbb{K}[Y]$  is generated by  $\bar{z}_1, \dots, \bar{z}_n$ ; it follows that  $\varphi^* = \alpha$ .  $\square$

**Corollary 1.8.4.** In Proposition 1.8.1, the affine variety  $X$  such that  $\mathbb{K}[X] \cong R$  is unique up to isomorphism.

Proposition 1.8.3 shows that by associating to an affine variety over  $\mathbb{K}$  the  $\mathbb{K}$ -algebra of its regular functions we get a contravariant equivalence between the category of affine varieties over  $\mathbb{K}$  (with maps the regular maps) and the category of finitely generated  $\mathbb{K}$ -algebras with no non-zero nilpotent elements. Note that if  $\varphi: S \rightarrow R$  is a morphism of finitely generated  $\mathbb{K}$ -algebras with no non-zero nilpotent elements the corresponding map (in the reverse direction) between the associated affine varieties is given by

$$\begin{array}{ccc} \text{Spec}_m(R) & \longrightarrow & \text{Spec}_m(S) \\ \mathfrak{m} & \mapsto & \varphi^{-1}(\mathfrak{m}) =: \mathfrak{m}^c \end{array}$$

(notice that  $\varphi^{-1}(\mathfrak{m})$  is maximal because  $\varphi$  is a morphism of  $\mathbb{K}$ -algebras).

## 1.9 Exercises

**Exercise 1.9.1.** Which of the following subsets of  $\mathbb{A}^2$  are locally closed? Which are closed?

- (a)  $X := \{(x, y) \mid \exp(2\pi\sqrt{-1}x) = 1\} \subset \mathbb{A}^2(\mathbb{C})$ .
- (b)  $Y := \{(t, t^2) \mid t \in \mathbb{K}\} \subset \mathbb{A}^2(\mathbb{K})$ .
- (c)  $W := \left\{ \left( \frac{2t}{t^2+1}, \frac{t^2-1}{t^2+1} \right) \mid t \in \mathbb{C} \setminus \{\pm\sqrt{-1}\} \right\} \subset \mathbb{A}^2(\mathbb{C})$ .
- (d)  $V := \{(t, tu) \mid (t, u) \in \mathbb{K}^2\} \subset \mathbb{A}^2(\mathbb{K})$ .

**Exercise 1.9.2.** Compute  $I(Z)$  for

1.  $Z = V(x^2 + 1) \subset \mathbb{A}^1(\mathbb{K})$ ,



$$2. Z = \mathbb{Z}^2 \subset \mathbb{A}^2(\mathbb{C}),$$

$$3. Z = V(x^2 - y^2, x^2 - xy) \subset \mathbb{A}^2(\mathbb{K}).$$

**Exercise 1.9.3.** Let  $M_{2,2}(\mathbb{C})$  be the complex vector-space of  $2 \times 2$  complex matrices. Let  $n > 0$  and let  $U_n \subset M_{2,2}(\mathbb{C})$  be the set of matrices  $T$  such that  $T^n = 1$  (here  $1 \in M_{2,2}(\mathbb{C})$  is the unit matrix).

1. Prove that  $U_n$  is a closed subset (for the Zariski Topology) of  $M_{2,2}(\mathbb{C})$ .

2. Describe the irreducible components of  $U_n$  and show that there are  $\binom{n+1}{2}$  of them.

**Exercise 1.9.4.** Let  $f_1, \dots, f_r \in \mathbb{K}[x, y]$  and suppose that

$$\gcd\{f_1, \dots, f_r\} = 1.$$

Show that  $V(f_1, \dots, f_r) \subset \mathbb{A}^2(\mathbb{K})$  is finite.

**Exercise 1.9.5.** Let  $X \subset \mathbb{A}^2(\mathbb{K})$  be a proper closed irreducible subset. Show that  $Z$  is either a singleton or an irreducible hypersurface.

**Exercise 1.9.6.** Let  $M_n(\mathbb{K})$  be the vector-space of  $n \times n$  matrices with entries in  $\mathbb{K}$ , and let  $M_n(\mathbb{K})_- \subset M_n(\mathbb{K})$  be the subspace of skew-symmetric matrices. Let  $X \in M_n(\mathbb{K})_-$ : then

$$X = \begin{bmatrix} 0 & x_{1,2} & \dots & \dots & x_{1,n} \\ -x_{1,2} & 0 & x_{2,3} & \dots & x_{2,n} \\ -x_{1,3} & -x_{1,3} & 0 & \dots & x_{3,n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -x_{1,n} & -x_{2,n} & \dots & \dots & 0 \end{bmatrix}$$

Thus  $\{x_{1,2}, \dots, x_{1,n}, x_{2,3}, \dots, x_{n-1,n}\}$  is a basis of the dual of  $M_n(\mathbb{K})_-$ , and hence  $\mathbb{K}[x_{1,2}, \dots, x_{1,n}, x_{2,3}, \dots, x_{n-1,n}]$  is the  $\mathbb{K}$  algebra of polynomial functions on  $M_n(\mathbb{K})_-$ . Let  $\Delta_n \subset M_n(\mathbb{K})_-$  be the set of  $n \times n$  singular skew-symmetric matrices, and let  $\delta_n$  be the polynomial on  $M_n(\mathbb{K})_-$  given by  $\delta_n(X) := \det X$ . Then  $\Delta_n$  is closed in  $M_n(\mathbb{K})_-$  because  $\Delta_n = V(\delta_n)$ . Prove the following:

(1.9.6a) If  $n$  is odd then  $\Delta_n = M_n(\mathbb{K})_-$ .

(1.9.6b) If  $n$  is even then  $\Delta_n$  is a hypersurface and  $I(\Delta_n) \neq (\delta_n)$ .

**Exercise 1.9.7.** An affine map

$$\begin{array}{ccc} \mathbb{A}^n & \longrightarrow & \mathbb{A}^n \\ Z & \longmapsto & A \cdot Z + B \end{array}$$

(here  $Z, B$  are column vectors with  $n$  entries and  $A \in \mathrm{GL}_n(\mathbb{K})$ ) is an automorphism of  $\mathbb{A}^n$ .

(1.9.7a) Show that every automorphism of  $\mathbb{A}^1$  is an affine map.

(1.9.7b) Let  $n \geq 2$ . Show that if  $f \in \mathbb{K}[z_1, \dots, z_{n-1}]$  then

$$\begin{array}{ccc} \mathbb{A}^n & \xrightarrow{\Phi_f} & \mathbb{A}^n \\ z & \longmapsto & (z_1, \dots, z_{n-1}, z_n + f(z_1, \dots, z_{n-1})) \end{array} \quad (1.9.4)$$

is an automorphism. Prove that  $\Phi_f$  is an affine map if and only if  $\deg f \leq 1$ .

**Exercise 1.9.8.** Show that one can prove the validity of Theorem 1.6.2 for  $\mathbb{A}^n$  by invoking unique factorization in  $\mathbb{K}[z_1, \dots, z_n]$ , without using the Nullstellensatz.

**Exercise 1.9.9.** Let  $K$  be a field. Given a finite-dimensional  $K$ -vector space  $V$  define the formal power series  $p_V \in \mathbb{Z}[[t]]$  as

$$P_V := \sum_{d=0}^{\infty} (\dim_K \operatorname{Sym}^d V) t^d$$

where  $\operatorname{Sym}^d V$  is the symmetric product of  $V$ . Thus if  $V = K[x_1, \dots, x_n]_1$  then  $S^d(K[x_1, \dots, x_n]_1) = K[x_1, \dots, x_n]_d$ .

1. Prove that if  $V = U \oplus W$  then  $P_V = P_U \cdot P_W$ .
2. Prove that if  $\dim_K V = n$  then  $P_V = (1-t)^{-n}$  and hence the equality in (1.5.10) holds.

**Exercise 1.9.10.** The purpose of the present exercise is to give a different proof of the properties of the Veronese map  $\nu_d^n$  discussed in Example 1.5.12, *valid if  $\operatorname{char} \mathbb{K} = 0$ , or more generally  $\operatorname{char} \mathbb{K}$  does not divide  $d!$* . Let

$$\begin{array}{ccc} \mathbb{P}(\mathbb{K}[T_0, \dots, T_n]_1) & \xrightarrow{\mu_d^n} & \mathbb{P}(\mathbb{K}[T_0, \dots, T_n]_d) \\ [L] & \mapsto & [L^d] \end{array} \quad (1.9.5)$$

and let  $\mathscr{W}_d^n = \operatorname{im}(\mu_d^n)$ . The above map can be identified with the Veronese map  $\nu_d^n$ . In fact, writing  $L \in \mathbb{K}[T_0, \dots, T_n]_1$  as  $L = \sum_{i=0}^n \alpha_i T_i$ , we see that  $[\alpha_0, \dots, \alpha_n]$  are coordinates on  $\mathbb{P}(\mathbb{K}[T_0, \dots, T_n]_1)$ , and they give an identification  $\mathbb{P}^n \xrightarrow{\sim} \mathbb{P}(\mathbb{K}[T_0, \dots, T_n]_1)$ . Moreover, let

$$\begin{array}{ccc} \mathbb{P}^{\binom{d+n}{n}-1} & \xrightarrow{\sim} & \mathbb{P}(\mathbb{K}[T_0, \dots, T_n]_d), \\ [\dots, \xi_I, \dots] & \mapsto & \sum_{\substack{I=(i_0, \dots, i_n) \\ i_0 + \dots + i_n = d}} \frac{d!}{i_0! \dots i_n!} \xi_I T^I \end{array}$$

where  $T^I = T_0^{i_0} \dots T_n^{i_n}$ . By Newton's formula  $(\sum_{i=0}^n \alpha_i T_i)^d = \sum_I \frac{d!}{i_0! \dots i_n!} \alpha^I T^I$ , we see that, modulo the above isomorphisms, the Veronese map  $\nu_d^n$  is identified with  $\mu_d^n$ , and hence  $\mathscr{V}_d^n$  is identified with  $\mathscr{W}_d^n$ .

Now let us show that  $\mathscr{W}_d^n$  is closed. The key observation is that  $[F] \in \mathscr{W}_d^n$  if and only if  $\frac{\partial F}{\partial Z_0}, \dots, \frac{\partial F}{\partial Z_n}$  span a 1-dimensional subspace of  $\mathbb{K}[Z_0, \dots, Z_n]$ . This may be proved by induction on  $\deg F$  and *Euler's identity*

$$\sum_{j=0}^n Z_j \frac{\partial F}{\partial Z_j} = (\deg F) \cdot F, \quad (1.9.6)$$

valid for  $F$  homogeneous. Now, the condition that  $\frac{\partial F}{\partial Z_0}, \dots, \frac{\partial F}{\partial Z_n}$  span a 1-dimensional subspace of  $\mathbb{K}[Z_0, \dots, Z_n]$  is equivalent to the vanishing of determinants of all  $2 \times 2$  minors of the matrix whose entries are the coordinates of  $\frac{\partial F}{\partial Z_0}, \dots, \frac{\partial F}{\partial Z_n}$ ; thus  $\mathscr{W}_d^n$  is closed.

In order to show that  $\mu_d^n$  is an isomorphism, we notice that if  $F = L^d$ , where  $L \in \mathbb{P}(\mathbb{K}[T_0, \dots, T_n]_1)$  is non zero, then for each  $i \in \{0, \dots, n\}$  the partial derivative  $\frac{\partial^{n-1} F}{\partial Z_i^{n-1}}$  is a multiple of  $L$  (eventually equal to 0 if  $\frac{\partial L}{\partial Z_i} = 0$ ), and that one at least of such  $(n-1)$ -th partial derivative is non zero. Thus, the inverse of  $\mu_d^n$  is the regular map  $\theta_d^n: \mathscr{W}_d^n \rightarrow \mathbb{P}(\mathbb{K}[T_0, \dots, T_n]_1)$  defined by

$$\theta_d^n([F]) := \begin{cases} \left[ \frac{\partial^{n-1} F}{\partial Z_0^{n-1}} \right] & \text{if } \frac{\partial^{n-1} F}{\partial Z_0^{n-1}} \neq 0, \\ \dots & \dots \\ \left[ \frac{\partial^{n-1} F}{\partial Z_n^{n-1}} \right] & \text{if } \frac{\partial^{n-1} F}{\partial Z_n^{n-1}} \neq 0. \end{cases} \quad (1.9.7)$$

**Exercise 1.9.11.** Let  $X \subset \mathbb{P}^n(\mathbb{C})$  and  $Y \subset \mathbb{P}^m(\mathbb{C})$  be complex quasi projective varieties defined over  $\mathbb{R}$ , and let  $\varphi: X \rightarrow Y$  be a regular map defined over  $\mathbb{R}$ . Note that the map  $X(\mathbb{R}) \rightarrow Y(\mathbb{R})$  defined by the restriction of  $\varphi$  to  $X(\mathbb{R})$  is continuous for the euclidean topologies of  $X(\mathbb{R})$  and  $Y(\mathbb{R})$ . Using this prove that the real quadrics

$$V(Z_0^2 - Z_1^2 - Z_2^2 - Z_3^2) \subset \mathbb{P}^3(\mathbb{C}), \quad V(Z_0^2 + Z_1^2 - Z_2^2 - Z_3^2) \subset \mathbb{P}^3(\mathbb{C}) \quad (1.9.8)$$

are not isomorphic over  $\mathbb{R}$  although they are isomorphic (actually projectively equivalent) over  $\mathbb{C}$ .

**Exercise 1.9.12.** Let  $R$  be an integral domain, and let  $(m, n) \in (\mathbb{N}^2 \setminus \{0\})$ . Let  $F \in R[X, Y]_m$  and  $G \in R[X, Y]_n$ . The *resultant*  $\mathscr{R}(F, G)$  is the element of  $R$  defined as follows. Consider the map of free  $R$ -modules

$$\begin{array}{ccc} R[X, Y]_{n-1} \oplus R[X, Y]_{m-1} & \xrightarrow{L(F, G)} & R[X, Y]_{m+n-1} \\ (\Phi, \Psi) & \mapsto & \Phi \cdot F + \Psi \cdot G \end{array} \quad (1.9.9)$$

and let  $S(F, G)$  be the matrix of  $L(F, G)$  relative to the basis

$$(X^{n-1}, 0), (X^{n-2}Y, 0), \dots, (Y^{n-1}, 0), (0, X^{m-1}), (0, X^{m-2}Y), \dots, (0, Y^{m-1}) \quad (1.9.10)$$

of the domain and the basis

$$X^{m+n-1}, X^{m+n-2}Y, \dots, XY^{m+n-2}, Y^{m+n-1} \quad (1.9.11)$$

of the codomain. Then  $\mathcal{R}(F, G)$  is defined by

$$\mathcal{R}(F, G) := \det S(F, G). \quad (1.9.12)$$

Explicitly: if

$$F = \sum_{i=0}^m a_i X^{m-i} Y^i, \quad G = \sum_{j=0}^n b_j X^{n-j} Y^j \quad (1.9.13)$$

then

$$\mathcal{R}(F, G) = \det \begin{pmatrix} a_0 & 0 & \cdots & 0 & b_0 & 0 & \cdots & 0 \\ a_1 & a_0 & \cdots & 0 & b_1 & b_0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \cdots & a_0 & \vdots & \vdots & \cdots & b_0 \\ a_m & a_{m-1} & \cdots & \vdots & b_n & b_{n-1} & \cdots & \vdots \\ 0 & a_m & \cdots & \vdots & 0 & b_n & \cdots & \vdots \\ 0 & 0 & \cdots & \vdots & 0 & 0 & \cdots & \vdots \\ \vdots & \vdots & \cdots & \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & \cdots & a_m & 0 & 0 & \cdots & b_n \end{pmatrix}. \quad (1.9.14)$$

Now let  $K$  be a field and  $K \subset \bar{K}$  be an algebraic closure of  $K$ . Let  $F \in K[X, Y]_m$  and  $G \in K[X, Y]_n$ .

- (a) Prove that  $\mathcal{R}(F, G) = 0$  if and only if there exists  $H \in K[X, Y]_d$  with  $d > 0$  which divides both  $F$  and  $G$  (in  $K[X, Y]$ ).
- (b) Prove that  $\mathcal{R}_{m,n}(F, G) = 0$  if and only if there exists a common non-trivial root of  $F$  and  $G$  in  $\bar{k}^2$ , i.e.  $[X_0, Y_0] \in \mathbb{P}_{\bar{k}}^1$  such that  $F(X_0, Y_0) = G(X_0, Y_0) = 0$ .
- (c) Let  $f(t, x) \in K[t_1, \dots, t_m][x]$  and  $g(t, x) \in K[t_1, \dots, t_m][x]$  (here  $t = t_1, \dots, t_m$ ) be polynomials of degrees  $m$  and  $n$  in  $x$  respectively, i.e.

$$f(t, x) = \sum_{i=1}^m a_i(t)x^{m-i}, \quad g(t, x) = \sum_{j=1}^n b_j(t)x^{n-j} \quad a_i(t), b_j(t) \in K[t_1, \dots, t_m], \quad a_0(t) \neq 0 \neq b_0(t).$$

We let

$$D(f, g) := \{\bar{t} \in \mathbb{A}^m(\bar{K}) \mid \exists x \in \bar{K} \text{ such that } f(\bar{t}, x) = g(\bar{t}, x) = 0\}.$$

Using the properties of the resultant proved above show that if  $f, g$  are both monic, i.e.  $a_0(t) = b_0(t) = 1$ , then there exists  $\varphi \in K[t_1, \dots, t_m]$  such that  $D(f, g) = V(\varphi)$ .

- (d) Give examples of  $f(t, x) \in K[t_1, \dots, t_m][x]$  and  $g(t, x) \in K[t_1, \dots, t_m][x]$  for which there exists no  $\varphi \in K[t_1, \dots, t_m]$  such that  $D(f, g) = V(\varphi)$ .

**Exercise 1.9.13.** We recall that if  $\phi: B \rightarrow A$  is a homomorphism of rings, and  $I \subset A, J \subset B$  are ideals, the contraction  $I^c \subset B$  and the extension  $J^e \subset A$  are the ideals defined as follows:

$$I^c := \phi^{-1}(I), \quad J^e := \left\{ \sum_{i=1}^r \lambda_i \phi(b_i) \mid \lambda_i \in A, b_i \in J \forall i = 1, \dots, r \right\} \quad (1.9.15)$$

(In other words,  $J^e$  is the ideal of  $A$  generated by  $\phi(J)$ .)

Let  $f: X \rightarrow Y$  be a regular map between affine varieties and suppose that  $f^*: \mathbb{K}[Y] \rightarrow \mathbb{K}[X]$  is injective.

- 1. Let  $p \in X$ . Prove that  $\mathfrak{m}_p^c = \mathfrak{m}_{f(p)}$ , in particular it is maximal.

1. QUASI PROJECTIVE VARIETIES

---

2. Let  $q \in Y$ . Prove that

$$f^{-1}(q) = \{p \in X \mid \mathfrak{m}_p \supset \mathfrak{m}_q^e\},$$

and conclude, by the Nulstellensatz, that  $f^{-1}(q)$  is not empty if and only if  $\mathfrak{m}_q^e \neq \mathbb{K}[X]$ .

**Exercise 1.9.14.** The left action of  $\mathrm{GL}_n(\mathbb{K})$  on  $\mathbb{A}^n$  defines a left action of  $\mathrm{GL}_n(\mathbb{K})$  on  $\mathbb{K}[z_1, \dots, z_n]$  as follows. Let  $\phi \in \mathbb{K}[z_1, \dots, z_n]$  and  $g \in \mathrm{GL}_n(\mathbb{K})$ . Let  $z$  be the column vector with entries  $z_1, \dots, z_n$ : we define  $g\phi \in \mathbb{K}[z_1, \dots, z_n]$  by letting

$$g\phi(X) := \phi(g^{-1} \cdot z).$$

Now let  $G < \mathrm{GL}_n(\mathbb{K})$  be a subgroup. The algebra of  $G$ -invariant polynomials is

$$\mathbb{K}[z_1, \dots, z_n]^G := \{\phi \in \mathbb{K}[z_1, \dots, z_n] \mid g\phi = \phi \forall g \in G\}.$$

(it is clearly a  $\mathbb{K}$ -algebra). Now suppose that  $G$  is finite. One identifies  $\mathbb{A}^n/G$  with an affine variety proceeding as follows.

1. Define the Reynolds operator as

$$\begin{aligned} \mathbb{K}[z_1, \dots, z_n] &\longrightarrow \mathbb{K}[z_1, \dots, z_n]^G \\ \phi &\longmapsto \frac{1}{|G|} \sum_{g \in G} g\phi. \end{aligned}$$

Prove the Reynolds identity

$$R(\phi\psi) = \phi R(\psi) \quad \forall \phi \in \mathbb{K}[z_1, \dots, z_n]^G.$$

2. Let  $I \subset \mathbb{K}[z_1, \dots, z_n]$  be the ideal generated by homogeneous  $\phi \in \mathbb{K}[z_1, \dots, z_n]^G$  of strictly positive degree (i.e. non-constant). By Hilbert's basis theorem there exists a finite basis  $\{\phi_1, \dots, \phi_d\}$  of  $I$ ; we may assume that each  $\phi_i$  is homogeneous and  $G$ -invariant. Prove that  $\mathbb{K}[z_1, \dots, z_n]^G$  is generated as  $\mathbb{K}$ -algebra by  $\phi_1, \dots, \phi_d$ . Since  $\mathbb{K}[z_1, \dots, z_n]^G$  is an integral domain with no nilpotents it follows that there exist an affine variety  $X$  (well-defined up to isomorphism) such that  $\mathbb{K}[X] \xrightarrow{\sim} \mathbb{K}[z_1, \dots, z_n]^G$ . One sets  $\mathbb{A}^n/G =: X$ .

3. Let  $\iota: \mathbb{K}[z_1, \dots, z_n]^G \hookrightarrow \mathbb{K}[z_1, \dots, z_n]$  be the inclusion map. By Proposition 1.8.3, there exist a unique regular map

$$\mathbb{A}^n \xrightarrow{\pi} X = \mathbb{A}^n/G. \tag{1.9.16}$$

such that  $\iota = \pi^*$ . Prove that

$$\pi(p) = \pi(q) \quad \text{if and only if} \quad q = gp \text{ for some } g \in G,$$

and that  $\pi$  is surjective. [Hint: Let  $J \subset \mathbb{K}[z_1, \dots, z_n]^G$  be an ideal. Show that  $J^e \cap \mathbb{K}[z_1, \dots, z_n]^G = J$  where  $J^e$  is the extension relative to the inclusion  $\iota$ .]

**Exercise 1.9.15.** Keep notation and hypotheses as in Exercise 1.9.14. Describe explicitly  $\mathbb{A}^n/G$  and the quotient map  $\pi: \mathbb{A}^n \rightarrow \mathbb{A}^n/G$  for the following groups  $G < \mathrm{GL}_n(\mathbb{K})$ :

1.  $n = 2$ ,  $G = \{\pm 1_2\}$ .
2.  $n = 2$ ,  $G = \left\langle \begin{pmatrix} \omega_k & 0 \\ 0 & \omega_k^{-1} \end{pmatrix} \right\rangle$  where  $\omega_k$  is a primitive  $k$ -th root of 1.
3.  $G = \mathcal{S}_n$ , the group of permutation of  $n$  elements viewed in the obvious way as a subgroup of  $\mathrm{GL}_n(\mathbb{K})$  (group of permutations of coordinates).

## Chapter 2

# Algebraic varieties

### 2.1 Introduction

The definition of quasi projective variety that we have given sounds very classical when compared to the definition of smooth manifold that one learns in a first course in Differential Geometry. In the present chapter we provide a definition of algebraic variety along the lines of the definition of smooth manifold. Quasi projective varieties are examples of algebraic varieties. \*\*\*\*\*

### 2.2 Algebraic prevarieties

#### Definition of algebraic prevariety

**Definition 2.2.1.** Let  $X$  be a topological space. An *algebraic atlas* of  $X$  defined over  $\mathbb{K}$  consists of an open covering  $\mathcal{A} = \{A_i\}_{i \in I}$  of  $X$ , and for each  $i \in I$  an affine variety  $V_i$  defined over  $\mathbb{K}$  (with the Zariski topology) together with a homeomorphism  $\varphi_i: V_i \rightarrow A_i$  (an *affine chart*), such that for each  $i, j \in I$  the transition map

$$\begin{array}{ccc} V_i \cap \varphi_i^{-1}(A_i \cap A_j) & \xrightarrow{\varphi_{j,i}} & V_j \cap \varphi_j^{-1}(A_j \cap A_i) \\ p & \mapsto & \varphi_j^{-1}(\varphi_i(p)) \end{array} \quad (2.2.1)$$

is a regular map of quasi projective varieties.

*Example 2.2.2.* Let  $X$  be a quasi projective variety. The collection  $\mathcal{A} := \{A_i\}_{i \in I}$  of open affine subsets of  $X$  is a basis for the Zariski topology of  $X$ , see Proposition 1.5.15. Choosing for every  $i \in I$  the identity affine chart  $\text{Id}_{A_i}: A_i \xrightarrow{\sim} A_i$  we get the *canonical algebraic atlas* of  $X$ .

Let  $(X, \mathcal{A})$  and  $(Y, \mathcal{B})$  be topological spaces with algebraic atlases over  $\mathbb{K}$ . Thus  $\mathcal{A} = \{A_i\}_{i \in I}$  and  $\mathcal{B} = \{B_j\}_{j \in J}$  are open coverings of  $X$  and  $Y$  respectively, and we are given homeomorphisms  $\varphi_i: V_i \xrightarrow{\sim} A_i$  and  $\psi_j: W_j \xrightarrow{\sim} B_j$  for all  $i \in I$  and  $j \in J$ , where  $V_i$  and  $W_j$  are affine varieties.

**Definition 2.2.3.** A *regular map*  $(X, \mathcal{A}) \rightarrow (Y, \mathcal{B})$  of topological spaces with algebraic atlases defined over  $\mathbb{K}$  is a continuous map  $f: X \rightarrow Y$  such that for all  $i \in I$  and  $j \in J$  the composition

$$\varphi_i^{-1}(A_i \cap f^{-1}B_j) \xrightarrow{\varphi_{i,\dots}} A_i \cap f^{-1}B_j \xrightarrow{f,\dots} B_j \xrightarrow{\psi_j^{-1}} W_j \quad (2.2.2)$$

is a regular map of (quasi projective) varieties. As a matter of notation we denote the map by  $f: (X, \mathcal{A}) \rightarrow (Y, \mathcal{B})$  or simply by  $f: X \rightarrow Y$ .

*Example 2.2.4.* Let  $X, Y$  be quasi projective varieties and let  $\mathcal{A}, \mathcal{B}$  be their canonical atlases, see Example 2.2.2. If  $f: X \rightarrow Y$  is a regular map, then it is a regular map of topological spaces with atlases.

Note that the composition of regular maps between topological spaces with algebraic atlases is regular, and the identity map  $(X, \mathcal{A}) \rightarrow (X, \mathcal{A})$  is regular.

**Definition 2.2.5.** Let  $X$  be a topological space. An algebraic atlas  $\mathcal{A}$  on  $X$  is *equivalent* to an algebraic atlas  $\mathcal{B}$  on  $X$  (both atlases defined over  $\mathbb{K}$ ) if the identity maps  $\text{Id}_X := (X, \mathcal{A}) \rightarrow (X, \mathcal{B})$  and  $\text{Id}_X := (X, \mathcal{B}) \rightarrow (X, \mathcal{A})$  are both regular.

Note that  $\mathcal{A}$  is equivalent to itself,  $\mathcal{A}$  equivalent to  $\mathcal{B}$  implies that  $\mathcal{B}$  equivalent to  $\mathcal{A}$ , and that if  $\mathcal{A}$  is equivalent to  $\mathcal{B}$  and  $\mathcal{B}$  is equivalent to  $\mathcal{C}$ , then  $\mathcal{A}$  is equivalent to  $\mathcal{C}$ . This justifies the use of the word “equivalent”.

**Definition 2.2.6.** An *algebraic prevariety defined over  $\mathbb{K}$*  (or simply a prevariety) is a couple  $(X, [\mathcal{A}])$  where  $X$  is a topological space and  $[\mathcal{A}]$  is an equivalence class of algebraic atlases. It is of *finite type* if there exists a representative of the equivalence class of  $\mathcal{A}$  with a finite set of indices. Let  $(X, [\mathcal{A}])$  and  $(Y, [\mathcal{B}])$  be algebraic prevarieties over  $\mathbb{K}$ ; a map  $f: X \rightarrow Y$  is *regular* if it is regular as map  $(X, \mathcal{A}) \rightarrow (Y, \mathcal{B})$  (this makes sense because if it is regular for one choice of representative atlases then it is regular for any choice).

Whenever the equivalence class of finite algebraic atlases  $[\mathcal{A}]$  is understood (or when we are too lazy to write it out) we denote  $(X, [\mathcal{A}])$  by  $X$ . The topology of an algebraic prevariety  $(X, [\mathcal{A}])$  is called (for obvious reasons) the Zariski topology of  $X$ .

*Remark 2.2.7.* A quasi projective variety with the equivalence class of its canonical atlas is a prevariety. In fact it is a prevariety of finite type because the Zariski topology is quasi-compact, see Corollary 1.3.9. Let  $X, Y$  be quasi projective varieties viewed as prevarieties (via their canonical atlases). A map  $f: X \rightarrow Y$  is regular (as map of prevarieties) if and only if it is a regular map of quasi projective varieties.

*Example 2.2.8.* A finite algebraic atlas for  $\mathbb{P}^n$  is as follows. Let  $A_i := \mathbb{P}_{Z_i}^n \cong \mathbb{A}^n$  for  $i \in \{0, \dots, n\}$ . Let  $z_0(i), \dots, z_{i-1}(i), z_{i+1}(i), \dots, z_n(i)$  (there is no  $z_i(i)$ ) be the affine coordinates on  $A_i$  given by  $z_s(i) := Z_s/Z_i$ . We can think of the coordinates  $z_s(i)$  as giving the map  $\varphi_i: \mathbb{A}^n \rightarrow A_i$ . Thus  $\varphi_i^{-1}(A_i \cap A_j) = \mathbb{A}^n \setminus V(z_j(i))$  and  $\varphi_j^{-1}(A_j \cap A_i) = \mathbb{A}^n \setminus V(z_i(j))$ . The transition map  $\varphi_{j,i}$  is determined by the formulae

$$\varphi_{j,i}^*(z_s(j)) := \begin{cases} z_j(i)^{-1} \cdot z_s(i) & \text{if } s \neq i \\ z_j^{-1}(i) & \text{if } s = i \end{cases} \quad (2.2.3)$$

*Example 2.2.9.* Let  $X$  be a prevariety. An open subset  $U \subset X$  can be given the structure of a prevariety so that the inclusion  $U \hookrightarrow X$  is regular. In fact let  $\{A_i\}_{i \in I}$  be an algebraic atlas, with affine charts  $\varphi_i: V_i \rightarrow A_i$ . For  $i \in I$  let  $W_i := \varphi_i^{-1}(A_i \cap U)$ . Then  $W_i$  is an open subset of  $V_i$ , and it is the union of its open affine subsets  $U_{i,j}$  where  $j \in J(i)$  for an index set  $J(i)$  which depends on  $i \in I$ . As algebraic atlas of  $U$  we take the collection  $\{\varphi_i(U_{i,j})\}_{i \in I, j \in J(i)}$  with affine charts  $\varphi_i|_{U_{i,j}}: U_{i,j} \rightarrow \varphi_i(U_{i,j})$ . Similarly, a closed subset  $Y \subset X$  can be given the structure of a prevariety so that the inclusion  $Y \hookrightarrow X$  is regular. We leave details to the reader. Lastly, if  $Y \subset X$  is a locally closed subset, say  $Y = U \cap W$  where  $U \subset X$  is open and  $W \subset X$  is closed, then  $Y$  is a closed subset of the prevariety  $U$ , and hence it inherits a structure of prevariety.

Prevarieties of finite type have an irreducible decomposition. First we prove the following result.

**Lemma 2.2.10.** *Let  $X$  be a prevariety of finite type, and let*

$$X \supset X_0 \supset X_1 \supset \dots \supset X_n \supset X_{n+1} \dots \quad (2.2.4)$$

*be a descending chain of closed subsets indexed by  $\mathbb{N}$ . Then the chain is stationary, i.e. there exists  $m \in \mathbb{N}$  such that  $X_n = X_{n+1}$  for all  $n \geq m$ .*

*Proof.* Let  $\{A_i\}_{i \in I}$  be a finite algebraic atlas, with affine charts  $\varphi_i: V_i \rightarrow A_i$ . For each  $i \in I$  the descending chain of closed subsets

$$V_i \supset \varphi_i^1(X_0) \supset \varphi_i^1(X_1) \supset \dots \supset \varphi_i^1(X_n) \supset \varphi_i^1(X_{n+1}) \dots \quad (2.2.5)$$

is stationary by Proposition 1.3.7. Thus there exists  $m_i \in \mathbb{N}$  such that  $X_n = X_{n+1}$  for all  $n \geq m_i$ . The proposition holds with  $m := \max\{m_i\}_{i \in I}$  (which exists because  $I$  is finite).  $\square$

**Proposition 2.2.11.** *If  $X$  is a prevariety of finite type it has an irreducible decomposition.*

*Proof.* Since Lemma 2.2.10 holds, one can repeat word-by-word the proof of Theorem 1.3.6.  $\square$

### Prevarieties defined over a subfield

Let  $F \subset \mathbb{K}$  be a subfield. Then one can repeat all the definitions above restricting to affine varieties and regular maps defined over  $F$  in order to define prevarieties defined over  $F$ . An algebraic atlas  $\mathcal{A} = \{A_i\}_{i \in I}$  on a topological space  $X$  with affine charts  $\varphi_i: V_i \rightarrow A_i$  is *defined over  $F$*  if

1. for all  $i \in I$  the affine variety  $V_i$  is defined over  $F$ ,
2. for all  $i, j \in I$  the quasi projective variety  $V_i \cap \varphi_i^{-1}(A_i \cap A_j)$  is defined over  $F$  and the transition map in (2.2.1) is regular.

Let  $(X, [\mathcal{A}])$  and  $(Y, [\mathcal{B}])$  be topological spaces  $X$  with algebraic atlases defined over  $F$ . A regular map  $f: (X, [\mathcal{A}]) \rightarrow (Y, [\mathcal{B}])$  is *defined over  $F$*  if the maps in (2.2.2) are defined over  $F$  for every  $i, j$ . This said it is clear how to mimick the definitions that we have given in order to define what are prevarieties defined over  $F$  and what are regular maps defined over  $F$ . Note that if  $(X, [\mathcal{A}])$  is a prevariety defined over  $F$  then  $X(F)$  makes sense, it consists of all the points  $\varphi_i(a)$  where  $a \in V_i(F)$ . This makes sense because if  $\varphi_i(a) \in A_j$  then  $\varphi_i(a) = \varphi_j(\varphi_j^{-1}(\varphi_i(a)))$  and since the map appearing in (2.2.1) is defined over  $F$  we have  $\varphi_j^{-1}(\varphi_i(a)) \in V_j(F)$ . Moreover if  $f: (X, [\mathcal{A}]) \rightarrow (Y, [\mathcal{B}])$  is a regular map defined over  $F$  then  $f(X(F)) \subset Y(F)$ .

### Gluing affine varieties

A method for producing a topological space with an algebraic atlas is to glue affine varieties along open subsets via regular maps. The simplest case is the following: let  $V, W$  be affine varieties, with isomorphic open subsets  $A \subset V$  and  $B \subset W$ , and let  $f: A \xrightarrow{\sim} B$  be an isomorphism. Let  $\sim$  be the equivalence relation on  $V \sqcup W$  generated by letting  $p \sim f(p)$  for  $p \in A \subset V$  (and  $f(p) \in B \subset W$ ). Let

$$X := V \sqcup W / \sim$$

be the quotient topological space. Let  $\pi: (V \sqcup W) \rightarrow X$  be the quotient map. The associated algebraic atlas of  $X$  is given by the open covering  $\{\pi(V), \pi(W)\}$  and the homeomorphisms  $V \xrightarrow{\sim} \pi(V)$ ,  $W \xrightarrow{\sim} \pi(W)$  obtained by restricting  $\pi$ .

*Example 2.2.12.* Let  $V = W = \mathbb{A}^1$ ,  $A = B = \mathbb{A}^1 \setminus \{0\}$ , and let

$$\begin{array}{ccc} A \supset \mathbb{A}^1 \setminus \{0\} & \xrightarrow{f} & \mathbb{A}^1 \setminus \{0\} \subset B \\ z & \mapsto & z^{-1} \end{array} \quad (2.2.6)$$

and

$$\begin{array}{ccc} A \supset \mathbb{A}^1 \setminus \{0\} & \xrightarrow{g} & \mathbb{A}^1 \setminus \{0\} \subset B \\ z & \mapsto & z \end{array} \quad (2.2.7)$$

Let  $X$  be the quotient topological space for the identification in (2.2.6), and let  $\mathcal{A}$  be the corresponding atlas. The prevariety  $(X, [\mathcal{A}])$  is isomorphic to  $\mathbb{P}^1$  with its canonical algebraic atlas. In fact let  $\tilde{\varphi}: V \sqcup W \rightarrow \mathbb{P}^1$  be the map defined by

$$\tilde{\varphi}(z) := \begin{cases} [1, z] & \text{if } z \in V, \\ [z, 1] & \text{if } z \in W. \end{cases} \quad (2.2.8)$$

Then  $\tilde{\varphi}$  descends to a regular map  $\varphi := (X, \mathcal{A}) \rightarrow \mathbb{P}^1$  which is an isomorphism. We will come back later to the prevariety corresponding to the identification in (2.2.7).

A more general version of the gluing construction is as follows. Suppose that we are given

- a family of affine varieties  $\{V_i\}_{i \in I}$ ,
- for all  $i, j \in I$  open subsets  $A_{i,j} \subset V_i$  and  $B_{i,j} \subset V_j$  and a (gluing) regular map  $\varphi_{j,i}: A_{i,j} \rightarrow B_{i,j}$ ,

subject to the following conditions:

**Hypothesis 2.2.13.** 1. For all  $i \in I$  we have  $A_{i,i} = B_{i,i} = V_i$  and  $\varphi_{i,i} = \text{Id}_{V_i}$ .

2. For all  $i, j \in I$  we have  $A_{j,i} = B_{i,j}$  (and of course  $B_{j,i} = A_{i,j}$ )  $\varphi_{i,j} = \varphi_{j,i}^{-1}$ .

3. For all  $i, j, k \in I$  and  $p \in A_{i,j}$  such that  $\varphi_{j,i}(p) \in A_{j,k}$  we have

$$\varphi_{k,j}(\varphi_{j,i}(p)) = \varphi_{k,i}(p). \quad (2.2.9)$$

*Gluing construction 2.2.14.* Let  $\sim$  be the relation on  $\bigsqcup_{i \in I} V_i$  defined by letting  $p \sim \varphi_{j,i}(p)$  for  $p \in A_{i,j} \subset V_i$  and arbitrary  $i, j \in I$ . Then  $\sim$  is an equivalence relation. Let

$$X := \bigsqcup_{i \in I} V_i / \sim$$

be the quotient topological space. Let  $\pi: \bigsqcup_{i \in I} V_i \rightarrow X$  be the quotient map. The associated algebraic atlas of  $X$  is given by the open covering  $\{\pi(V_i)\}_{i \in I}$  and the homeomorphisms  $V_i \xrightarrow{\sim} \pi(V_i)$  obtained by restricting  $\pi$ .

*Example 2.2.15.* Let  $I := \{0, 1, \dots, n\}$  and let  $V_i = \mathbb{A}^n$  for all  $i \in I$ . Let  $(z_0(i), \dots, z_{i-1}(i), z_{i+1}(i), \dots, z_n(i))$  be affine coordinates on  $V_i$  (note that there is no coordinate  $z_i(i)$ ). Let  $A_{i,j} := \mathbb{A}^n \setminus V(z_j(i))$  and  $B_{i,j} := \mathbb{A}^n \setminus V(z_i(j))$ . We define  $\varphi_{j,i}: A_{i,j} \rightarrow B_{i,j}$  by letting

$$\varphi_{j,i}^*(z_s(j)) := \begin{cases} z_j(i)^{-1} \cdot z_s(i) & \text{if } s \neq i \\ z_j^{-1}(i) & \text{if } s = i \end{cases} \quad (2.2.10)$$

One checks that Items (1), (2) and (3) above hold. The corresponding prevariety  $(X, [\mathcal{A}])$  is isomorphic to  $\mathbb{P}^n$ , see Example 2.2.8. Explicitly, let  $\tilde{\varphi}: V_0 \sqcup \dots \sqcup V_n \rightarrow \mathbb{P}^n$  be the map defined by setting

$$\begin{array}{ccc} V_i & \longrightarrow & \mathbb{P}^n \\ (z_0(i), \dots, z_{i-1}(i), z_{i+1}(i), \dots, z_n(i)) & \mapsto & [z_0(i), \dots, z_{i-1}(i), 1, z_{i+1}(i), \dots, z_n(i)] \end{array} \quad (2.2.11)$$

Then  $\tilde{\varphi}$  descends to a regular map  $\varphi := (X, \mathcal{A}) \rightarrow \mathbb{P}^n$  which is an isomorphism.

*Example 2.2.16.* Let  $(Y, [\mathcal{A}])$  be a prevariety, with affine charts  $\psi_i := V_i \rightarrow A_i$ . For  $i, j \in I$  let  $A_{i,j} := \psi_i^{-1}(A_i \cap A_j)$  and  $B_{i,j} := \psi_j^{-1}(A_j \cap A_i)$ . Let

$$\begin{array}{ccc} A_{i,j} & \xrightarrow{\varphi_{j,i}} & B_{i,j} \\ p & \mapsto & \varphi_j^{-1}(\varphi_i(p)) \end{array} \quad (2.2.12)$$

Then Hypothesis 2.2.13 holds, hence there is a corresponding prevariety  $(X, [\mathcal{B}])$ , where  $\mathcal{B}$  is the algebraic atlas  $\{\pi(V_i)\}_{i \in I}$ . Clearly  $(X, [\mathcal{B}])$  is isomorphic to  $(Y, \mathcal{A})$  - this generalizes Example 2.2.15.

As shown by the example above, the gluing construction is at the heart of the definition of prevariety. In fact they are two different point of views of the same objects. In the definition of a prevariety we are given a topological space and a collection of affine charts, in the gluing construction we are given a collection of affine varieties and gluing data  $\varphi_{j,i}$  and we define a topological space.



### 2.3 Products and algebraic varieties

Let  $X$  be a prevariety. The Zarisky Topology of  $X$  is not Hausdorff unless  $X$  is finite. Nonetheless  $X$  might share key properties of Hausdorff topological spaces. In fact suppose that  $X$  is an affine variety. Thus we may assume that  $X \subset \mathbb{A}^n$  is closed. The square  $X \times X \subset \mathbb{A}^n \times \mathbb{A}^n = \mathbb{A}^{2n}$  is closed, so it is an affine variety. Moreover the diagonal  $\Delta_X \subset X \times X$  is closed in the Zariski topology. In fact let  $(x_1, \dots, x_n, y_1, \dots, y_n)$  be the obvious affine coordinates on  $\mathbb{A}^n \times \mathbb{A}^n$ : then  $\Delta_X$  is the intersection of  $X \times X$  and the closed subset  $V(x_1 - y_1, x_2 - y_2, \dots, x_n - y_n)$ . Recall that a topological space  $X$  is Hausdorff if and only if the diagonal in  $X \times X$  (with the product topology) is closed. So apparently we have a contradiction: if  $X$  is an affine variety which is not finite then it is not Hausdorff but its diagonal is closed in  $X \times X$ . In fact this is not a contradiction because if  $X$  is not finite the Zariski topology on  $X \times X$  is much finer than the product topology. The conclusion is that the right version of Hausdorffness for an algebraic prevariety  $X$  is that the diagonal be closed in  $X \times X$ . Thus our first step is to define the product of prevarieties.

#### Products in a category

We start by recalling the definition of product of two objects in a category.

**Definition 2.3.1.** Let  $\mathcal{C}$  be a category, and let  $X, Y \in \text{Ob}(\mathcal{C})$  be objects of  $\mathcal{C}$ . A *product of  $X$  and  $Y$*  consists of an object  $Z \in \text{Ob}(\mathcal{C})$  and morphisms  $p_X: Z \rightarrow X$  and  $p_Y: Z \rightarrow Y$  (the projections) which have the following universal property. Assume that  $W \in \text{Ob}(\mathcal{C})$  and that  $f: W \rightarrow X, g: W \rightarrow Y$  are morphisms. Then there exists a unique morphism  $h: W \rightarrow Z$  such that the following is a commutative diagram

$$\begin{array}{ccc}
 W & & Y \\
 & \searrow g & \\
 & \exists! h & \\
 & \searrow & Z \xrightarrow{p_Y} Y \\
 & \searrow f & \downarrow p_X \\
 & & X
 \end{array}
 \tag{2.3.1}$$

Suppose that a product  $W$  of  $X$  and  $Y$  exists. If  $W'$  is another product of  $X$  and  $Y$  (with projections  $p'_X: W' \rightarrow X$  and  $p'_Y: W' \rightarrow Y$ ), then there exists a unique morphism  $h: W \rightarrow W'$  commuting with the projections, i.e.  $p'_X \circ h = p_X$  and  $p'_Y \circ h = p_Y$ . Of course we also have the corresponding morphism  $h': W' \rightarrow W$ . By the unicity requirement in the definition of product the compositions  $h' \circ h$  and  $h \circ h'$  are equal to the identities of  $W$  and  $W'$ . Thus we have a well defined isomorphism between any two products of  $X$  and  $Y$  (assuming a product exists). Since the product is well defined up to (unique) isomorphism it makes sense to talk of “the” product of  $X$  and  $Y$ . One denotes it by  $X \times Y$ . We denote by  $(f, g)$  the unique morphism  $h$  appearing in (2.3.1).

*Example 2.3.2.* Let **Sets** be the category of sets (one has to be careful with definitions or one runs into Russell’s paradox, but we ignore this point here). If  $X, Y \in \text{Ob}(\mathbf{Sets})$  i.e.  $X, Y$  are sets, then the Cartesian product  $X \times Y$  with projections  $p_X(x, y) := x$  and  $p_Y(x, y) := y$  is the product of  $X$  and  $Y$  in the category **Sets**.

*Example 2.3.3.* Let **Grps** be the category of groups. If  $G, H \in \text{Ob}(\mathbf{Grps})$  i.e.  $G, H$  are groups, then the direct product  $G \times H$  with projections  $p_G(g, h) := g$  and  $p_H(g, h) := h$  is the product of  $G$  and  $H$  in the category **Grps**.

*Example 2.3.4.* Let  $S$  be a set, and let **Sets**/ $S$  be the category whose objects are maps  $f: X \rightarrow S$  from a set  $X$  to  $S$ , and morphisms from a map  $f: X \rightarrow S$  to a map  $g: Y \rightarrow S$  are morphisms  $\varphi: X \rightarrow Y$

which commute with  $f$  and  $g$ , i.e. a commutative diagram

$$\begin{array}{ccc} X & \xrightarrow{\varphi} & Y \\ & \searrow f & \swarrow g \\ & & S \end{array} \quad (2.3.2)$$

The product of  $f: X \rightarrow S$  and  $g: Y \rightarrow S$  in the category  $\mathbf{Sets}/S$  is given by the object

$$\begin{array}{ccc} X \times_S Y := \{(x, y) \in X \times Y \mid f(x) = g(y)\} & \longrightarrow & S \\ (x, y) & \mapsto & f(x) (= g(y)) \end{array} \quad (2.3.3)$$

(the *fiber product* of  $X$  and  $Y$  over  $S$ ) with projections given by the restrictions of the projections  $X \times Y \rightarrow X$  and  $X \times Y \rightarrow Y$ .

### Products of affine varieties

Let  $X, Y$  be affine varieties. Thus, we may assume that  $X \subset \mathbb{A}^m$  and  $Y \subset \mathbb{A}^n$  are closed subsets. Then  $X \times Y \subset \mathbb{A}^m \times \mathbb{A}^n \cong \mathbb{A}^{m+n}$  is a closed subset, and the projections  $p_X: X \times Y \rightarrow X$  and  $p_Y: X \times Y \rightarrow Y$  given by the two projections are regular.

**Proposition 2.3.5.** *Keeping notation as above,  $X \times Y$  with projections  $p_X, p_Y$  is the product of  $X$  and  $Y$  in the category of prevarieties.*

*Proof.* Let  $W$  be a prevariety and let  $f: W \rightarrow X, g: W \rightarrow Y$  be regular maps. We must prove that there exists a regular map  $h: W \rightarrow X \times Y$  such that  $p_X \circ h = f, p_Y \circ h = g$ , and that  $h$  is unique. Since prevarieties are sets (with extra structure) and regular maps between prevarieties are maps between the underlying sets (satisfying suitable conditions), if  $h$  exists it is necessarily given by

$$\begin{array}{ccc} W & \xrightarrow{(f,g)} & X \times Y \\ p & \mapsto & (f(p), g(p)) \end{array} \quad (2.3.4)$$

Thus all we need to prove is that  $(f, g)$  is regular. As we showed (see Example 2.2.16) any prevariety is obtained by the gluing construction in 2.2.14. Thus  $W$  is obtained by gluing affine varieties  $\{V_i\}_{i \in I}$  as in 2.2.14. To simplify notation denote  $\pi(V_i) \subset W$  by  $V_i$ . It suffices to show that the restriction of  $(f, g)$  to  $V_i$  is regular. Since  $f$  and  $g$  are regular both the restrictions of  $f$  and  $g$  to  $V_i$  are regular. It follows at once that the restriction of  $(f, g)$  to  $V_i$  is regular.  $\square$

The  $\mathbb{K}$  algebra of regular functions of  $X \times Y$  is constructed from  $\mathbb{K}[X]$  and  $\mathbb{K}[Y]$  as follows. Let  $\pi_X: X \times Y \rightarrow X$  and  $\pi_Y: X \times Y \rightarrow Y$  be the projections. The  $\mathbb{K}$ -bilinear map

$$\begin{array}{ccc} \mathbb{K}[X] \times \mathbb{K}[Y] & \longrightarrow & \mathbb{K}[X \times Y] \\ (f, g) & \mapsto & \pi_X^*(f) \cdot \pi_Y^*(g) \end{array} \quad (2.3.5)$$

induces a linear map

$$\mathbb{K}[X] \otimes_{\mathbb{K}} \mathbb{K}[Y] \longrightarrow \mathbb{K}[X \times Y]. \quad (2.3.6)$$

**Proposition 2.3.6.** *The map in (2.3.6) is an isomorphism.*

*Proof.* We may assume that  $X \subset \mathbb{A}^m$  and  $Y \subset \mathbb{A}^n$  are closed subsets. Since  $X \times Y \subset \mathbb{A}^{m+n}$  is closed the map in (2.3.6) is surjective by Theorem 1.6.2. It remains to prove injectivity, i.e. the following: if  $A \subset \mathbb{K}[X]$  and  $B \subset \mathbb{K}[Y]$  are finite-dimensional complex vector subspaces, then the map  $A \otimes B \rightarrow \mathbb{K}[X \times Y]$  obtained by restriction of (2.3.6) is injective. Let  $\{f_1, \dots, f_a\}, \{g_1, \dots, g_b\}$  be bases of  $A$  and  $B$ . By considering the maps

$$\begin{array}{ccc} X & \longrightarrow & \mathbb{K}^a & & Y & \longrightarrow & \mathbb{K}^b \\ z & \mapsto & (f_1(z), \dots, f_a(z)) & & z & \mapsto & (g_1(z), \dots, g_b(z)) \end{array} \quad (2.3.7)$$

we get that there exist  $p_1, \dots, p_a \in X$  and  $q_1, \dots, q_b \in Y$  such that the square matrices  $(f_i(p_j))$  and  $(g_i(q_j))$  are non-singular. By change of bases, we may assume that  $f_i(p_j) = \delta_{ij}$  and  $g_k(q_h) = \delta_{kh}$ . Computing the values of  $\pi_X^*(f_i) \cdot \pi_Y^*(g_j)$  on  $(p_s, q_t)$  for  $1 \leq i, s \leq a$  and  $1 \leq j, t \leq b$  we get that the functions  $\dots, \pi_X^*(f_i) \cdot \pi_Y^*(g_j), \dots$  are linearly independent. Thus  $A \otimes B \rightarrow \mathbb{K}[W \times Z]$  is injective.  $\square$

### Products of prevarieties

**Proposition 2.3.7.** *Let  $X, Y$  be prevarieties. There exists a product of  $X$  and  $Y$  in the category of prevarieties.*

*Proof.* By Example 2.2.16  $X$  is obtained by gluing affine varieties  $\{V_i\}_{i \in I}$  as in 2.2.14, and  $Y$  is obtained by gluing affine varieties  $\{W_j\}_{j \in J}$ . More precisely for each  $i_1, i_2 \in I$  we have regular maps  $\varphi_{i_2, i_1}^X: A_{i_1, i_2}^X \rightarrow B_{i_1, i_2}^X$ , where  $A_{i_1, i_2}^X \subset V_{i_1}$  and  $B_{i_1, i_2}^X \subset V_{i_2}$  are open subsets, and they are the gluings defining  $X$ . Analogously, for each  $j_1, j_2 \in J$  we have regular maps  $\varphi_{j_2, j_1}^Y: A_{j_1, j_2}^Y \rightarrow B_{j_1, j_2}^Y$ , where  $A_{j_1, j_2}^Y \subset W_{j_1}$  and  $B_{j_1, j_2}^Y \subset W_{j_2}$  are open subsets, and they are the gluings defining  $Y$ . Then we can glue the collection of affine varieties  $\{V_i \times W_j\}_{(i, j) \in I \times J}$  as follows. For  $(i_1, j_1), (i_2, j_2) \in I \times J$  let

$$A_{(i_1, j_1), (i_2, j_2)} := A_{i_1, i_2}^X \times A_{j_1, j_2}^Y \subset V_{i_1} \times W_{j_1}, \quad B_{(i_1, j_1), (i_2, j_2)} := B_{i_1, i_2}^X \times B_{j_1, j_2}^Y \subset V_{i_2} \times W_{j_2} \quad (2.3.8)$$

These are open subsets of  $V_{i_1} \times W_{j_1}$  and  $V_{i_2} \times W_{j_2}$  respectively. We let

$$A_{(i_1, j_1), (i_2, j_2)} \xrightarrow{\varphi_{(i_1, j_1), (i_2, j_2)}} B_{(i_1, j_1), (i_2, j_2)} \\ (p, q) \mapsto (\varphi_{i_2, i_1}^X(p), \varphi_{j_2, j_1}^Y(q)) \quad (2.3.9)$$

This collection of affine varieties and gluing maps satisfy the conditions in Hypothesis 2.2.13. Let  $Z$  be the prevariety obtained by gluing the  $\{V_i \times W_j\}_{(i, j) \in I \times J}$ 's as specified above. We have obvious maps  $p_X: Z \rightarrow X$  and  $p_Y: Z \rightarrow Y$ . In fact let  $z \in Z$ . Then  $z = (p, q) \in V_i \times W_j$  for some  $(i, j) \in I \times J$  (by no means unique). Here, in order to simplify notation, we denote  $\pi^X(V_i) \subset X$  and  $\pi^Y(W_j) \subset Y$  by  $V_i$  and  $W_j$  respectively. Then we let  $p_X(p, q) := p$  and  $p_Y(p, q) := q$ . As is easily checked the maps  $p_X, p_Y$  are regular. We claim that  $Z$  with the regular maps  $p_X$  and  $p_Y$  is the categorical product of  $X$  and  $Y$ . First note that the map of sets  $(p_X, p_Y): Z \rightarrow X \times Y$  is bijective. Hence, given regular maps  $f: U \rightarrow X$  and  $g: U \rightarrow Y$ , there is a unique map  $h: U \rightarrow Z$  of sets commuting with the projections. In fact if  $u \in U$  we let  $h(u)$  be the unique  $z \in Z$  such that  $p_X(z) = f(u)$  and  $p_Y(z) = g(u)$ . Arguing as in the proof of Proposition 2.3.5 one shows that  $h$  is a regular map.  $\square$

*Remark 2.3.8.* We stress that the categorical product of prevarieties  $X, Y$  is canonically identified, as a set, with the Cartesian product of  $X$  and  $Y$ .

*Remark 2.3.9.* Let  $X, Y$  be prevarieties, and let  $X_0 \subset X, Y_0 \subset Y$  be locally closed subsets. Then  $X_0 \times Y_0 \subset X \times Y$  is a locally closed subset. The restrictions of the projections  $X \times Y \rightarrow X$  and  $X \times Y \rightarrow Y$  to  $X_0 \times Y_0$  define regular maps  $p_{X_0}: X_0 \times Y_0 \rightarrow X_0$  and  $p_{Y_0}: X_0 \times Y_0 \rightarrow Y_0$ . As is easily checked,  $X_0 \times Y_0$  with the regular maps  $p_{X_0}$  and  $p_{Y_0}$  is the product of  $X_0$  and  $Y_0$ .

*Remark 2.3.10.* If  $F \subset \mathbb{K}$  is a subfield and  $X, Y$  are prevarieties over  $F$ , then  $X \times Y$  is defined over  $F$ . We leave the reader check this fact.

### Separated prevarieties

Let  $X$  be a prevariety. The *diagonal*  $\Delta_X \subset X \times X$  is defined to be

$$\Delta_X := \{(x, x) \mid x \in X\}. \quad (2.3.10)$$

This makes sense because as a set  $X \times X$  is identified with the Cartesian square of  $X$ .

*Example 2.3.11.* Let  $X$  be an affine variety. Thus we may assume that  $X \subset \mathbb{A}^n$  is closed. Then  $X \times X \subset \mathbb{A}^{2n}$  is closed. Letting  $(x_1, \dots, x_n, y_1, \dots, y_n)$  be the standard affine coordinates on  $\mathbb{A}^{2n}$ , we have

$$\Delta_X \cap (X \times X) = V(x_1 - y_1, x_2 - y_2, \dots, x_n - y_n) \cap (X \times X). \quad (2.3.11)$$

hence the diagonal of an affine variety is closed.

*Remark 2.3.12.* Let  $X$  be an algebraic prevariety. The diagonal  $\Delta_X$  is a locally closed subset of  $X \times X$ . In fact by Example 2.2.16  $X$  is obtained by gluing affine varieties  $\{V_i\}_{i \in I}$  as in 2.2.14. The open subsets  $V_i \times V_j$  for  $(i, j) \in I$  cover  $X$  (to simplify notation we denote  $\pi(V_i)$ ,  $\pi(A_{ij})$  and  $\pi(B_{ij})$  by  $V_i$ ,  $A_{ij}$  and  $B_{ij}$  respectively). Thus it suffices to show that the intersection of  $\Delta_X$  with each open subset  $V_i \times V_j$  is locally closed in  $V_i \times V_j$ . We have

$$\Delta_X \cap (V_i \times V_j) = \{(x, y) \in A_{ij} \times B_{ij} \mid y = \varphi_{ji}(x)\}. \quad (2.3.12)$$

Arguing as in Example 2.3.11 we get that  $\Delta_X \cap (V_i \times V_j)$  is a closed subset of the open set  $A_{ij} \times B_{ij}$ , and hence  $\Delta_X$  is a closed subset of the open subset of  $X \times X$  given by the union of all the  $A_{ij} \times B_{ij}$ 's.

*Example 2.3.13.* Let  $(Y, [\mathcal{B}])$  be the prevariety defined by the second atlas (given by the regular map  $g$ ) in Example 2.2.12. Then the diagonal is not closed in  $Y \times Y$ . In fact denote by  $V, W$  the open subsets  $\pi(V), \pi(W)$  respectively. Then  $V \times W \cong \mathbb{A}^2$  and

$$\Delta_Y \cap (V \times W) = \{(z, z) \in \mathbb{A}^2 \mid z \neq 0\}, \quad (2.3.13)$$

which is not closed.

**Definition 2.3.14.** An algebraic prevariety  $X$  is *separated* if the diagonal  $\Delta_X \subset X \times X$  is closed.

*Example 2.3.15.* By Example 2.3.11 an affine variety  $X$  with its canonical structure of prevariety is separated.

*Remark 2.3.16.* Let  $X$  be a prevariety. We may assume that  $X$  is obtained by gluing affine varieties  $\{V_i\}_{i \in I}$  as in 2.2.14. As usual we denote  $\pi(V_i)$ ,  $\pi(A_{ij})$  and  $\pi(B_{ij})$  by  $V_i$ ,  $A_{ij}$  and  $B_{ij}$  respectively. Since  $\{V_i \times V_j\}_{(i,j) \in I^2}$  is an open covering of  $X \times X$  the diagonal  $\Delta_X$  is closed in  $X \times X$  if and only if  $\Delta_X \cap (V_i \times V_j)$  is closed for all  $(i, j) \in I^2$ . Since  $\Delta_X \cap (V_i \times V_i)$  is closed, see Example 2.3.15, it suffices to check that  $\Delta_X \cap (V_i \times V_j)$  is closed for all couples  $i \neq j$ . We can halve the verifications needed because  $\Delta_X \cap (V_i \times V_j)$  is closed if and only if  $\Delta_X \cap (V_j \times V_i)$  is closed. Moreover, since  $\Delta_X \cap (A_{ij} \times B_{ij})$  is closed (see Remark 2.3.12), in order to show that  $\Delta_X \cap (V_i \times V_j)$  is closed it suffices to show that there exists a subset  $C \subset A_{ij} \times B_{ij}$  containing  $\Delta_X \cap (V_i \times V_j)$  which is closed in  $V_i \times V_j$ .

*Example 2.3.17.* Let  $(X, [\mathcal{A}])$  be the prevariety defined by the first atlas (given by the regular map  $f$ ) in Example 2.2.12. Then  $(X, [\mathcal{A}])$  is separated. In fact denote by  $V, W$  the open subsets  $\pi(V), \pi(W)$  respectively. Then  $V \times W \cong \mathbb{A}^2$  and

$$\Delta_Y \cap (V \times W) = V(wz - 1). \quad (2.3.14)$$

Since  $(X, [\mathcal{A}])$  is isomorphic to  $\mathbb{P}^1$ , we get that  $\mathbb{P}^1$  is separated.

*Example 2.3.18.* Let  $(Y, [\mathcal{B}])$  be the prevariety defined by the second atlas (given by the regular map  $g$ ) in Example 2.2.12. The diagonal  $\Delta_Y$  is not closed in  $Y \times Y$ , see Example 2.3.13. Hence  $(Y, [\mathcal{B}])$  is not separated.

The following result shows that separated prevarieties enjoy a key property of Hausdorff topological spaces.

**Proposition 2.3.19.** *Let  $X, Y$  be prevarieties, and assume that  $Y$  is separated. If  $f, g: X \rightarrow Y$  are regular maps, then the subset of  $X$  defined by*

$$\{x \in X \mid f(x) = g(x)\} \quad (2.3.15)$$

*is closed in  $X$ .*

*Proof.* By the universal property of the product  $Y \times Y$  (we let  $p_1, p_2$  be the projections to  $Y$ ) we have the regular map  $(f, g): X \rightarrow Y \times Y$  such that  $p_1 \circ (f, g) = f$  and  $p_2 \circ (f, g) = g$ . Let  $W$  be the subset of  $X$  appearing in (2.3.15). Then  $W = (f, g)^{-1}(\Delta_Y)$ . Since  $Y$  is separated  $\Delta_Y$  is closed and hence  $W$  is closed. □

### Algebraic varieties

**Definition 2.3.20.** An algebraic prevariety is an *algebraic variety* if it is of finite type and separated.

An affine variety is an algebraic variety by Remark 2.2.7 and Example 2.3.15. Also  $\mathbb{P}^1$  is an algebraic variety by Remark 2.2.7 and Example 2.3.17. More generally, a quasi projective variety is an algebraic variety.

**Proposition 2.3.21.** *A quasi projective variety (with its canonical structure of prevariety) is an algebraic variety.*

*Proof.* We have already noticed that a quasi projective variety is of finite type, see Remark 2.2.7. It remains to show that it is separated. First we consider  $\mathbb{P}^n$  (the key case). Following Example 2.2.15,  $\mathbb{P}^n$  is obtained by gluing  $(n + 1)$  copies  $\{V_0, \dots, V_n\}$  of affine space  $\mathbb{A}^n$ . As usual we use the same symbol  $V_i$  to denote  $\pi(V_i)$ . It suffices to check that  $\Delta_{\mathbb{P}^n} \cap (V_i \times V_j)$  is closed in  $(V_i \times V_j)$  for all  $i \neq j$ . By the formulae for the gluing maps in (2.2.10) we get that  $\Delta_{\mathbb{P}^n} \cap (V_i \times V_j)$  is contained in the closed subset  $V(x_j(i) \cdot x_i(j) - 1) \subset (V_i \times V_j)$ . Since this closed subset is contained in  $A_{ij} \times B_{ij}$  we are done, see the last sentence of Remark 2.3.16. Now let  $X \subset \mathbb{P}^n$  be a locally closed subset. Then  $X \times X$  is a locally closed subset of  $\mathbb{P}^n \times \mathbb{P}^n$  and  $\Delta_{\mathbb{P}^n} \cap (X \times X) = \Delta_X$ . Since  $\Delta_{\mathbb{P}^n}$  is closed in  $\mathbb{P}^n \times \mathbb{P}^n$ , it follows that  $\Delta_X$  is closed in  $X \times X$ . □

Next we consider constructions which, starting from an algebraic variety (or algebraic varieties) produce another algebraic variety.

Let  $X$  be an algebraic prevariety, with algebraic atlas  $\mathcal{A} = \{A_i\}_{i \in I}$  and affine charts  $\varphi_i: V_i \rightarrow A_i$ . If  $U \subset X$  is an open subset then we define an algebraic atlas on  $U$  as follows. For  $i \in I$  the open subset  $\varphi_i^{-1}(V_i \cap U) \subset V_i$  is the union of its open affine subsets. The restriction of  $\varphi_i$  to an open affine subset  $W_{i,k} \subset \varphi_i^{-1}(V_i \cap U)$  defines a homeomorphism  $\psi_{i,k}: W_{i,k} \rightarrow \varphi_i(W_{i,k})$ , and  $\varphi_i(W_{i,k})$  is open in  $U$ . Thus  $U$  is covered by the open subsets  $\varphi_i(W_{i,k})$  and the maps  $\psi_{i,k}$  give affine charts. If the algebraic atlas  $\mathcal{A}$  is replaced by an equivalent one we get an equivalent atlas of  $U$ . Thus we have equipped  $U$  with a canonical structure of prevariety. Note that the inclusion map  $U \hookrightarrow X$  is regular.

**Proposition 2.3.22.** *Let  $X$  be an algebraic variety. If  $U \subset X$  is an open subset with its canonical structure of prevariety (see Example 2.2.9), then  $U$  is a variety.*

*Proof.* We must prove that  $U$  is of finite type and separated. Since  $X$  is of finite type there exists a finite algebraic atlas  $\mathcal{A} = \{A_1, \dots, A_n\}$  with affine charts  $\varphi_i: V_i \rightarrow A_i$ . The open subset  $\varphi_i^{-1}(V_i \cap U)$  is the union of its affine open subsets, and since the Zariski topology of a quasi projective variety is quasi compact it is the union of a finite family of open subsets. From this it follows that  $U$  has a finite algebraic atlas, and hence is of finite type. The subset  $(U \times U) \subset (X \times X)$  is the (categorical) square of  $U$ , this follows from the universal property of the product  $X \times X$  and the fact that the inclusion  $U \hookrightarrow X$  is regular. Since the diagonal  $\Delta_U$  is equal to the intersection  $(U \times U) \cap \Delta_X$  in  $X \times X$  and  $\Delta_X$  is closed in  $X \times X$ , it follows that  $\Delta_U$  is closed in  $U \times U$ , and hence  $U$  is separated. □

Arguing as above one proves the following result.

**Proposition 2.3.23.** *Let  $X$  be an algebraic variety. If  $Y \subset X$  is a locally closed subset with its canonical structure of prevariety (see Example 2.2.9), then  $Y$  is a variety.*

**Proposition 2.3.24.** *If  $X, Y$  are algebraic varieties, then the product  $X \times Y$  is an algebraic variety.*

*Proof.* By hypothesis there exists finite algebraic atlases  $\mathcal{A} = \{A_i\}_{i \in I}$ ,  $\mathcal{B} = \{B_j\}_{j \in J}$  of  $X$  and  $Y$  respectively, with affine charts  $\varphi_i: V_i \rightarrow A_i$  and  $\psi_j: W_j \rightarrow B_j$ . Then  $\mathcal{A} \times \mathcal{B} := \{A_i \times B_j\}_{(i,j) \in I \times J}$ , with affine charts  $\varphi_i \times \psi_j: V_i \times W_j \rightarrow A_i \times B_j$  is a finite algebraic atlas of  $X \times Y$ . Thus  $X \times Y$  is of finite type. The projection maps  $f: (X \times Y) \times (X \times Y) \rightarrow X \times X$  and to  $Y \times Y$

$$\begin{aligned} (X \times Y) \times (X \times Y) &\xrightarrow{f} X \times X & (X \times Y) \times (X \times Y) &\xrightarrow{g} Y \times Y \\ ((x_1, y_1), (x_2, y_2)) &\mapsto (x_1, x_2) & ((x_1, y_1), (x_2, y_2)) &\mapsto (y_1, y_2) \end{aligned} \quad (2.3.16)$$

are regular (by the universal property of products) and hence continuous. Thus  $\Delta_{X \times Y} = f^{-1}(\Delta_X) \cap g^{-1}(\Delta_Y)$  is closed. This proves that  $X \times Y$  is separated.  $\square$

### Products of quasi projective varieties

In the present subsection we prove the following result.

**Proposition 2.3.25.** *If  $X$  and  $Y$  are quasi projective varieties, then  $X \times Y$  is a quasi projective variety.*

Before proving Proposition 2.3.25 we go through a few preliminary results. A polynomial  $F(W; Z) \in \mathbb{K}[W_0, \dots, W_m, Z_0, \dots, Z_n]$  is *bihomogeneous of degree  $(d, e)$*  if  $F(\lambda W; \mu Z) = \lambda^d \mu^e F(W; Z)$  for all  $\lambda, \mu \in \mathbb{K}$ . Let  $F_i \in \mathbb{K}[W_0, \dots, W_m, Z_0, \dots, Z_n]$  for  $i \in \{1, \dots, r\}$  be a bihomogeneous polynomial of degree  $(d_i, e_i)$ . Then it makes sense to let

$$V(F_1, \dots, F_r) := \{([W], [Z]) \in \mathbb{P}^m \times \mathbb{P}^n \mid F_1(W; Z) = \dots = F_r(W; Z) = 0\}. \quad (2.3.17)$$

**Claim 2.3.26.** *A subset  $X \subset \mathbb{P}^m \times \mathbb{P}^n$  is closed if and only if there exist bihomogeneous polynomials  $F_1, \dots, F_r \in \mathbb{K}[W_0, \dots, W_m, Z_0, \dots, Z_n]$  such that  $X = V(F_1, \dots, F_r)$ .*

*Proof.* We have

$$\mathbb{P}^m \times \mathbb{P}^n = \bigcup_{\substack{0 \leq i \leq m \\ 0 \leq j \leq n}} \mathbb{P}_{W_i}^m \times \mathbb{P}_{Z_j}^n \quad (2.3.18)$$

and each of the open subsets  $\mathbb{P}_{W_i}^m \times \mathbb{P}_{Z_j}^n$  is isomorphic to  $\mathbb{A}^{m+n}$ . If  $F_1, \dots, F_r$  are as above, then  $V(F_1, \dots, F_r) \cap \mathbb{P}_{W_i}^m \times \mathbb{P}_{Z_j}^n$  is clearly closed. It follows that  $V(F_1, \dots, F_r)$  is closed. Now suppose that  $X \subset \mathbb{P}^m \times \mathbb{P}^n$  is closed. Then  $X \cap \mathbb{P}_{W_i}^m \times \mathbb{P}_{Z_j}^n$  is closed for every  $i, j$ , and hence there exists  $f_1^{i,j}, \dots, f_s^{i,j} \in \mathbb{K}[\frac{W_0}{W_i}, \dots, \frac{W_m}{W_i}, \frac{Z_0}{Z_j}, \dots, \frac{Z_n}{Z_j}]$  such that

$$X \cap \mathbb{P}_{W_i}^m \times \mathbb{P}_{Z_j}^n = V(f_1^{i,j}, \dots, f_s^{i,j}). \quad (2.3.19)$$

If  $d \gg 0$  and  $e \gg 0$  are natural numbers then  $F_l^{i,j} := W_i^d \cdot Z_j^e \cdot f_l^{i,j} \left( \frac{W_0}{W_i}, \dots, \frac{W_m}{W_i}, \frac{Z_0}{Z_j}, \dots, \frac{Z_n}{Z_j} \right)$  is bihomogeneous of degree  $(d, e)$  and  $V(F_l^{i,j}) = \overline{V(f_l^{i,j})} \cup V(W_i) \cup V(Z_j)$ . Thus  $X$  is the zero locus of all the bihomogeneous polynomials  $F_l^{i,j}$ 's.  $\square$

*Remark 2.3.27.* In the statement of Claim 2.3.26 we may require, if we wish, that all the  $F_i$ 's are bihomogeneous of degrees  $(d_i, d_i)$ , i.e. of the same degrees in both variables.

Next we introduce Segre varieties and Segre maps. Let  $M_{m+1, n+1}(\mathbb{K})$  be the  $\mathbb{K}$  vector space of complex  $(m+1) \times (n+1)$  matrices. Row and column indices for matrices in  $M_{m+1, n+1}(\mathbb{K})$  start from 0. Thus we denote them by

$$T = \begin{pmatrix} T_{00} & T_{01} & \dots & T_{0n} \\ T_{10} & T_{11} & \dots & T_{1n} \\ \dots & \dots & \dots & \dots \\ T_{m0} & T_{m1} & \dots & T_{mn} \end{pmatrix} \quad (2.3.20)$$

Let

$$\Sigma_{m,n} := \{[T] \in \mathbb{P}(M_{m+1,n+1}(\mathbb{K})) \mid \text{rk } T \leq 1\}.$$

Then  $\Sigma_{m,n}$  is a projective variety in  $\mathbb{P}(M_{m+1,n+1}(\mathbb{K})) = \mathbb{P}^{mn+m+n}$ . In fact  $[T] \in \mathbb{P}(M_{m+1,n+1}(\mathbb{K}))$  belongs to  $\Sigma_{m,n}$  if and only if the determinants of all  $2 \times 2$  minors of  $T$  vanish. This is the *Segre variety* in  $\mathbb{P}(M_{m+1,n+1}(\mathbb{K}))$ .

If  $[W] \in \mathbb{P}^m$  and  $[Z] \in \mathbb{P}^n$ , viewed as column matrices, then  $W \cdot Z^t \in M_{m+1,n+1}(\mathbb{K})$  and the rank of  $W \cdot Z^t$  is 1. If we rescale  $W$  or  $Z$  then  $W \cdot Z^t$  gets rescaled. Thus we have a well defined *Segre map*

$$\begin{array}{ccc} \mathbb{P}^m \times \mathbb{P}^n & \xrightarrow{\sigma_{m,n}} & \Sigma_{m,n} \\ ([W], [Z]) & \mapsto & [W \cdot Z^t] \end{array} \quad (2.3.21)$$

Explicitly

$$\sigma_{m,n}([W], [Z]) = \left[ \begin{array}{cccc} W_0 \cdot Z_0 & W_0 \cdot Z_1 & \dots & W_0 \cdot Z_n \\ W_1 \cdot Z_0 & W_1 \cdot Z_1 & \dots & W_1 \cdot Z_n \\ \dots & \dots & \dots & \dots \\ W_m \cdot Z_0 & W_m \cdot Z_1 & \dots & W_m \cdot Z_n \end{array} \right] \quad (2.3.22)$$

**Proposition 2.3.28.** *The Segre map in (2.3.21) is an isomorphism of algebraic varieties.*

*Proof.* First we prove that the Segre map is bijective. Let  $[T] \in \Sigma_{m,n}$ . Then  $T$  has rank 1 because  $T \neq 0$ . Hence the associated linear map  $L_T: \mathbb{K}^{n+1} \rightarrow \mathbb{K}^{m+1}$  (given by  $L_T(U) := T \cdot U$ , where  $U$  is a column matrix) can be factored as  $L_T = L_W \circ L_{Z^t}$  where  $L_{Z^t}: \mathbb{K}^{n+1} \rightarrow \mathbb{K}$  is surjective and  $L_W: \mathbb{K} \rightarrow \mathbb{K}^{m+1}$  is injective. This gives that  $T = W \cdot Z^t$ . We also get that  $\ker(L_T) = \ker(L_{Z^t})$  and  $\text{im}(L_T) = \text{im}(L_W)$ . Thus  $W$  and  $Z$  are determined by  $[T]$  up to a scalar factor, and hence  $\sigma_{m,n}$  is injective.

Next we prove that the Segre map is a homeomorphism. Let  $C \subset \Sigma_{m,n}$  be a closed subset, i.e.  $C = \Sigma_{m,n} \cap V(P_1, \dots, P_r)$  where  $P_i \in \mathbb{K}[T_{00}, T_{01}, \dots, T_{mn}]_{d_i}$ . Then

$$\sigma_{m,n}^{-1}(C) = V(P_1(W_0 \cdot Z_0, W_0 \cdot Z_1, \dots, W_m \cdot Z_n), \dots, P_r(W_0 \cdot Z_0, W_0 \cdot Z_1, \dots, W_m \cdot Z_n)). \quad (2.3.23)$$

Since  $P_i(W_0 \cdot Z_0, W_0 \cdot Z_1, \dots, W_m \cdot Z_n)$  for  $i \in \{1, \dots, r\}$  is a bihomogeneous polynomial (of degree  $(d_i, d_i)$ ), it follows that  $\sigma_{m,n}^{-1}(C)$  is closed in  $\mathbb{P}^m \times \mathbb{P}^n$ , see Claim 2.3.26. This shows that  $\sigma_{m,n}$  is continuous. Now suppose that  $D \subset \mathbb{P}^m \times \mathbb{P}^n$  is closed. By Claim 2.3.26 there exist bihomogeneous polynomials  $F_1, \dots, F_r \in \mathbb{K}[W_0, \dots, W_m, Z_0, \dots, Z_n]$  such that  $X = V(F_1, \dots, F_r)$ . As noticed in Remark 2.3.27 we may assume that  $F_i$  is bihomogeneous polynomial of degree  $(d_i, d_i)$  for each  $i \in \{1, \dots, r\}$ , and hence there exists  $P_i \in \mathbb{K}[T_{00}, T_{01}, \dots, T_{mn}]_{d_i}$  such that

$$P_i(W_0 \cdot Z_0, W_0 \cdot Z_1, \dots, W_m \cdot Z_n) = F_i(W_0, \dots, W_m; Z_0, \dots, Z_n). \quad (2.3.24)$$

This implies that  $\sigma_{m,n}(D) = \Sigma_{m,n} \cap V(P_1, \dots, P_r)$  and hence  $\sigma_{m,n}(D)$  is closed. Thus also the inverse of the Segre map is continuous and hence  $\sigma_{m,n}$  is a homeomorphism.

It remains to show that the Segre map is an isomorphism of algebraic varieties. Recall that we have the open covering in (2.3.18). Now  $\sigma_{m,n}$  maps the affine space  $\mathbb{P}_{W_i}^m \times \mathbb{P}_{Z_j}^n$  to the open set  $\Sigma_{m,n} \setminus V(T_{ij})$  and, as is easily checked the map

$$\mathbb{P}_{W_i}^m \times \mathbb{P}_{Z_j}^n \longrightarrow \Sigma_{m,n} \setminus V(T_{ij}) \quad (2.3.25)$$

is an isomorphism (of affine spaces). It follows that  $\sigma_{m,n}$  is an isomorphism of algebraic varieties.  $\square$

*Proof of Proposition 2.3.25.* We may assume that  $X \subset \mathbb{P}^m$  and  $Y \subset \mathbb{P}^n$  are locally closed. Then  $X \times Y \subset \mathbb{P}^m \times \mathbb{P}^n$  is locally closed, and it is the product of  $X$  and  $Y$ , see Remark 2.3.9. Since the Segre map  $\sigma_{m,n}: \mathbb{P}^m \times \mathbb{P}^n \rightarrow \Sigma_{m,n}$  is an isomorphism, it restricts to an isomorphism between  $X \times Y$  and the locally closed subset  $\sigma_{m,n}(X \times Y) \subset \Sigma_{m,n}$ . Since  $\Sigma_{m,n}$  is a projective variety,  $\sigma_{m,n}(X \times Y)$  is a quasi projective variety.  $\square$

## 2.4 Complete varieties

In the present section we introduce the notion of complete varieties, which are the analogues of compact topological space in the category of prevarieties. The prime example of complete varieties are projective varieties. We note that a complex quasi projective variety is complete if and only if, equipped with the Euclidean topology, it is compact. Since every quasi projective variety is quasi compact (and also every prevariety of finite type), one defines “compactness” for algebraic varieties by relying on a different characterization of compact topological spaces.

Let  $M$  be a topological space. Then  $M$  is quasi compact, i.e. every open covering has a finite subcovering, if and only if  $M$  is universally closed, i.e. for any topological space  $T$ , the projection map  $T \times M \rightarrow T$  is closed, i.e. it maps closed sets to closed sets. (See tag/005M in [Sta].) A quasi projective variety  $X$  is quasi compact, but it is not generally true that, for a variety  $T$ , the projection  $T \times X \rightarrow T$  is closed. In fact, let  $X \subset \mathbb{P}^n$  be locally closed; then  $\Delta_X$ , the diagonal of  $X$ , is closed in  $X \times \mathbb{P}^n$ , because it is the intersection of  $X \times X \subset \mathbb{P}^n \times \mathbb{P}^n$  with the diagonal  $\Delta_{\mathbb{P}^n} \subset \mathbb{P}^n \times \mathbb{P}^n$ , which is closed. The projection  $X \times \mathbb{P}^n \rightarrow \mathbb{P}^n$  maps  $X$  to  $X$ , hence if  $X$  is not closed in  $\mathbb{P}^n$ , then  $X$  is not universally closed. This does not contradict the result in topology quoted above, because the Zariski topology of the product of quasi projective varieties is not the product topology.

**Definition 2.4.1.** A prevariety  $X$  is *complete* (or *proper over  $\mathbb{K}$* ) if  $X$  is an algebraic variety and it is universally closed, i.e. for any prevariety  $T$ , the projection map  $T \times X \rightarrow T$  is closed.

As noticed above, if  $X \subset \mathbb{P}^n$  is not closed (e.g.  $\mathbb{P}_{Z_i}^n$  if  $n > 0$ ), then it is not universally closed, and hence  $X$  is not complete.

The following is a key result.

**Theorem 2.4.2** (Main Theorem of elimination theory). *Projective varieties are complete.*

*Proof.* Let  $X$  be projective variety. Since  $X$  is an algebraic variety we must prove that  $X$  is universally closed.

By hypothesis we may assume that  $X \subset \mathbb{P}^n$  is closed. We claim that it suffices to prove that  $\mathbb{P}^n$  is universally closed. In fact assume that  $\mathbb{P}^n$  is universally closed. Let  $T$  be a prevariety and let  $\pi_T^X: T \times X \rightarrow T$  be the projection. Let  $C \subset T \times X$  be closed. Since  $T \times X \subset T \times \mathbb{P}^n$  is closed,  $C$  is closed also in  $T \times \mathbb{P}^n$ . Let  $\pi_T^{\mathbb{P}^n}: T \times \mathbb{P}^n \rightarrow T$  be the projection. Then  $\pi_T^X(C) = \pi_T^{\mathbb{P}^n}(C)$ , and hence it is closed because by assumption  $\mathbb{P}^n$  is universally closed. Since  $T$  is covered by open affine subsets, we may assume that  $T$  is affine, i.e.  $T$  is (isomorphic to) a closed subset of  $\mathbb{A}^m$  for some  $m$ . Lastly, we may as well assume that  $T = \mathbb{A}^m$ .

To sum up: it suffices to prove that if  $C \subset \mathbb{A}^m \times \mathbb{P}^n$  is closed, then  $\pi(C)$  is closed in  $\mathbb{A}^m$ , where  $\pi: \mathbb{A}^m \times \mathbb{P}^n \rightarrow \mathbb{A}^m$  is the projection. We will show that  $(\mathbb{A}^m \setminus \pi(C))$  is open. By Claim 2.3.26 there exist  $F_i \in \mathbb{K}[t_1, \dots, t_m, Z_0, \dots, Z_n]$  for  $i = 1, \dots, r$ , *homogeneous* as polynomial in  $Z_0, \dots, Z_n$  such that

$$C = \{(t, [Z]) \mid 0 = F_1(t, Z) = \dots = F_r(t, Z)\}.$$

Suppose that  $F_i \in \mathbb{K}[t_1, \dots, t_m][Z_0, \dots, Z_n]_{d_i}$  i.e.  $F_i$  is homogeneous of degree  $d_i$  in  $Z_0, \dots, Z_n$ . Let  $\bar{t} \in (\mathbb{A}^m \setminus \pi(C))$ . By Hilbert’s Nullstellensatz, there exists  $N \geq 0$  such that

$$(F_1(\bar{t}, Z), \dots, F_r(\bar{t}, Z)) \supset \mathbb{K}[Z_0, \dots, Z_n]_N. \quad (2.4.1)$$

We may assume that  $N \geq d_i$  for  $1 \leq i \leq r$ . For  $t \in \mathbb{A}^m$  let

$$\begin{array}{ccc} \mathbb{K}[Z_0, \dots, Z_n]_{N-d_1} \times \dots \times [Z_0, \dots, Z_n]_{N-d_r} & \xrightarrow{\Phi(t)} & \mathbb{K}[Z_0, \dots, Z_n]_N \\ (G_1, \dots, G_r) & \mapsto & \sum_{i=1}^r G_i \cdot F_i \end{array}$$

Thus  $\Phi(t)$  is a linear map: choose bases of domain and codomain and let  $M(t)$  be the matrix associated to  $\Phi(t)$ . Clearly the entries of  $M(t)$  are elements of  $\mathbb{K}[t_1, \dots, t_m]$ . By hypothesis  $\Phi(\bar{t})$  is surjective and hence there exists a maximal minor of  $M(t)$ , say  $M_{I,J}(t)$ , such that  $\det M_{I,J}(\bar{t}) \neq 0$ . The open  $(\mathbb{A}^m \setminus V(\det M_{I,J}))$  is contained in  $(T \setminus \pi(C))$ . This finishes the proof of Theorem 2.4.2.  $\square$



Next give a few general results on complete varieties.

**Proposition 2.4.3.** *Let  $X, Y$  be complete (algebraic) varieties.*

1. *If  $W \subset X$  is closed then (with its canonical structure of variety, see Proposition 2.3.23)  $W$  is complete.*
2. *The product  $X \times Y$  is complete.*

*Proof.* (1): We must check that  $W$  is universally closed. One argues as in the second paragraph of the proof of Theorem 2.4.2. (2): By Proposition 2.3.24  $X \times Y$  is an algebraic variety. Hence it remains to check that  $X \times Y$  is universally closed. Let  $T$  be a prevariety, and let  $C \subset T \times (X \times Y)$  be closed. Factoring the projection  $\pi: T \times (X \times Y) \rightarrow T$  as the composition of  $f: T \times (X \times Y) \rightarrow T \times Y$  followed by  $g: T \times Y \rightarrow T$ , we get that  $f(C) \subset T \times Y$  is closed because  $X$  is universally closed, and  $g(f(C))$ , i.e.  $\pi(C)$ , is closed because  $Y$  is universally closed.  $\square$

If  $f: X \rightarrow Y$  is a regular map between prevarieties, the *graph of  $f$*  is the subset  $\Gamma_f$  of  $X \times Y$  defined by

$$\Gamma_f := \{(x, f(x)) \mid x \in X\}. \quad (2.4.2)$$

**Lemma 2.4.4.** *Let  $f: X \rightarrow Y$  be a regular map between prevarieties, and suppose that  $Y$  is separated. Then the graph of  $f$  is closed in  $X \times Y$ .*

*Proof.* The map

$$\begin{array}{ccc} X \times Y & \xrightarrow{f \times \text{Id}_Y} & Y \times Y \\ (x, y) & \mapsto & (f(x), y) \end{array} \quad (2.4.3)$$

is regular, and hence continuous. Since  $\Gamma_f = (f \times \text{Id}_X)^{-1}(\Delta_Y)$  and  $\Delta_Y$  is closed in  $Y \times Y$  (because  $Y$  is separated) it follows that  $\Gamma_f$  is closed.  $\square$

**Proposition 2.4.5.** *Let  $X, Y$  be algebraic varieties, with  $X$  complete and  $Y$  separated. If  $f: X \rightarrow Y$  is a regular map then it is closed.*

*Proof.* Since, by Proposition 2.4.3, closed subsets of  $X$  are complete varieties, it suffices to prove that  $f(X)$  is closed in  $Y$ . Let  $\pi: X \times Y \rightarrow Y$  be the projection map. By Lemma 2.4.4  $\Gamma_f$  is closed in  $X \times Y$ , and hence  $\pi(\Gamma_f)$  is closed in  $Y$  because  $X$  is complete. Since  $f(X) = \pi(\Gamma_f)$  we are done.  $\square$

**Corollary 2.4.6.** *Let  $X$  be a complete variety and let  $Y \subset X$  be a locally closed subset (with its canonical structure of variety, see Proposition 2.3.23). Then  $Y$  is complete if and only if  $Y$  is closed.*

*Proof.* If  $Y$  is closed then it is complete by Proposition 2.4.3. Conversely, suppose that  $Y$  is complete. Since the inclusion map  $i: Y \hookrightarrow X$  is regular and  $X$  is separated,  $Y = i(Y)$  is closed in  $X$  by Proposition 2.4.5.  $\square$

*Remark 2.4.7.* In particular let  $Y \subset \mathbb{P}^n$  be locally closed and projective. Then, since projective varieties are complete (by the Main Theorem of Elimination Theory), it follows that  $Y$  is closed in  $\mathbb{P}^n$ . By way of contrast, notice that it is *not* true that a locally-closed subset of  $\mathbb{A}^n$  is affine if and only if it is closed. In fact the complement of a hypersurface  $V(f) \subset \mathbb{A}^n$  is affine but not closed.

**Corollary 2.4.8.** *Let  $X$  be a complete variety. A regular map  $f: X \rightarrow \mathbb{K}$  is locally constant. If  $X$  is irreducible (recall Proposition 2.2.11) then  $f$  is constant.*

*Proof.* Composing  $f$  with the inclusion  $j: \mathbb{K} \hookrightarrow \mathbb{P}^1$

$$\begin{array}{ccc} \mathbb{K} & \xrightarrow{j} & \mathbb{P}^1 \\ z & \mapsto & [1, z] \end{array} \quad (2.4.4)$$

we get the regular map  $j \circ f: X \rightarrow \mathbb{P}^1$ . By Proposition 2.4.5  $j \circ f(X)$  is closed, i.e.  $j \circ f(X) = V(I)$  for some homogeneous ideal  $I \subset \mathbb{K}[Z_0, Z_1]$ . Since  $[0, 1] \notin j \circ f(X)$  there exists a non zero polynomial  $P \in I$  and hence  $j \circ f(X) = f(X)$  is contained in the finite set  $V(P)$ . The second statement follows at once from the first.  $\square$

## 2.5 Algebraic vector bundles

A very important notion in Topology and in Differential Geometry is that of continuous and  $C^\infty$  vector bundle respectively. One defines an analogous notion in the context of algebraic varieties.

**Definition 2.5.1.** Let  $X$  be an algebraic variety defined over  $\mathbb{K}$ . A rank  $r$  algebraic vector bundle over  $X$  (we call it a *line bundle* if  $r = 1$ ) consists of the following data:

1. A regular map  $\pi: E \rightarrow X$  of algebraic varieties.
2. For each  $x \in X$  a structure of  $\mathbb{K}$  vector space of dimension  $r$  on the fiber  $E(x) := \pi^{-1}(x)$ .

These data are subject to the following conditions.

- (a) There exist an open cover  $X = \bigcup_{\alpha \in A} U_\alpha$  and for each  $\alpha \in A$  an isomorphism of varieties  $\varphi_\alpha: \pi^{-1}(U_\alpha) \xrightarrow{\sim} U_\alpha \times \mathbb{K}^r$  such that the diagram

$$\begin{array}{ccc} \pi^{-1}(U_\alpha) & \xrightarrow{\varphi_\alpha} & U_\alpha \times \mathbb{K}^r \\ & \searrow \pi|_{U_\alpha} & \swarrow \text{pr}_\alpha \\ & & U_\alpha \end{array} \quad (2.5.1)$$

where  $\text{pr}_\alpha$  is the projection.

- (b) For each  $\alpha \in A$  and  $x \in U_\alpha$  the restriction of  $\varphi_\alpha$  to  $E(x)$ , which is an isomorphism

$$\varphi_\alpha(x): E(x) \xrightarrow{\sim} \{x\} \times \mathbb{K}^r = \mathbb{K}^r \quad (2.5.2)$$

by Item (a), is a  $\mathbb{K}$ -linear map (and hence an isomorphism of  $\mathbb{K}$  vector spaces)

*Example 2.5.2.* Let  $X$  be an algebraic variety. Then  $E := X \times \mathbb{K}^r$  with  $\pi: E \rightarrow X$  given by the projection map is clearly a rank  $r$  vector bundle on  $X$ .

**Definition 2.5.3.** Let  $X$  be an algebraic variety (defined over  $\mathbb{K}$ ), and let  $\pi: E \rightarrow X$ ,  $\rho: F \rightarrow X$  be vector bundles on  $X$ . A *morphism of vector bundles*  $E \rightarrow F$  consists of a regular map of algebraic varieties  $\varphi: E \rightarrow F$  such that the diagram

$$\begin{array}{ccc} E & \xrightarrow{\varphi} & F \\ & \searrow \pi & \swarrow \rho \\ & & X \end{array} \quad (2.5.3)$$

is commutative, and such that for every  $x \in X$  the restriction of  $\varphi$  to  $E(x)$  is a linear map  $\varphi(x): E(x) \rightarrow F(x)$  of  $\mathbb{K}$  vector spaces.

The identity map  $\text{im}_E: E \rightarrow E$  of a vector bundle is a morphism of vector bundles, and the composition of morphisms of vector bundles on  $X$  is a morphism of vector bundles on  $X$ . Hence vector bundles on  $X$  form a category. In particular we have the notion of *isomorphic vector bundles on  $X$* .

**Definition 2.5.4.** Let  $X$  be an algebraic variety. A vector bundle  $E$  of rank  $r$  on  $X$  is *trivial* if it is isomorphic to the vector bundle in Example 2.5.2.

Next we define a fundamental line bundle on projective space. Let  $V$  be a finite dimensional  $\mathbb{K}$  vector space. We view points of  $\mathbb{P}(V)$  as 1 dimensional (vector) subspaces  $\ell \subset V$ . Let  $L \subset \mathbb{P}(V) \times V$  be defined by

$$L := \{(\ell, v) \in \mathbb{P}(V) \times V \mid v \in \ell\}. \tag{2.5.4}$$

We claim that  $L$  is a closed subset of  $\mathbb{P}(V) \times V$ , and hence an algebraic variety. In fact choose a basis of  $V$  so that  $V$  and  $\mathbb{P}(V)$  are identified with  $\mathbb{K}^{n+1}$  and  $\mathbb{P}^n$  respectively. Then

$$L = \{([Z_0, \dots, Z_n], (W_0, \dots, W_n)) \in \mathbb{P}^n \times \mathbb{K}^{n+1} \mid \text{rk} \begin{pmatrix} Z_0 & \dots & Z_n \\ W_0 & \dots & W_n \end{pmatrix} \leq 1\}. \tag{2.5.5}$$

This shows that  $L$  is closed in  $\mathbb{P}(V) \times V$ . Let  $\pi: L \rightarrow \mathbb{P}(V)$  be the restriction of the projection  $\mathbb{P}(V) \times V \rightarrow \mathbb{P}(V)$ . If  $\ell \in \mathbb{P}(V)$  then  $L(\ell) = \pi^{-1}(\ell) = \ell$  and this gives the structure of 1 dimensional  $\mathbb{K}$  vector space to  $L(\ell)$ . Let  $i \in \{0, \dots, n\}$ . We define  $\varphi_i: \pi^{-1}(\mathbb{P}_{Z_i}^n) \rightarrow \mathbb{P}_{Z_i}^n \times \mathbb{K}$  as follows:

$$\begin{array}{ccc} \pi^{-1}(\mathbb{P}_{Z_i}^n) & \xrightarrow{\varphi_i} & \mathbb{P}_{Z_i}^n \times \mathbb{K} \\ ([Z], Y) & \mapsto & ([Z], Y_i) \end{array} \tag{2.5.6}$$

This shows that  $L \rightarrow \mathbb{P}(V)$  is a line bundle. It is called the *tautological line bundle on  $\mathbb{P}(V)$* . If  $n > 0$  then  $L$  is not trivial. Before showing this we introduce sections of a vector bundle.

**Definition 2.5.5.** Let  $X$  be an algebraic variety, and let  $\pi: E \rightarrow X$  be a vector bundle on  $X$ . A *section of  $E$*  is a map  $\sigma: X \rightarrow E$  such that  $\pi \circ \sigma = \text{Id}_X$ . The section  $\sigma$  is *regular* if it is regular as map of algebraic varieties.

*Example 2.5.6.* Let  $X$  be an algebraic variety, and let  $E = X \times \mathbb{K}^r$  with  $\pi: E \rightarrow X$  the projection. A regular section  $\sigma: X \rightarrow E$  is equivalent to regular maps  $f_i: X \rightarrow \mathbb{K}$  for  $i \in \{1, \dots, r\}$ . In fact given such  $f_i$  we let  $\sigma(x) := (x, (f_1(x), \dots, f_r(x)))$ . Viceversa\*\*\*

The set of sections of  $E$  is denoted by

$$\Gamma(X, E) := \{\sigma: X \rightarrow E \mid \sigma \text{ is a section of } E\}. \tag{2.5.7}$$

The sum of regular sections is a regular section, and the product of an element of  $\mathbb{K}$  by a regular section is a regular section. With these operations  $\Gamma(X, E)$  acquires the structure of a  $\mathbb{K}$  vector space. Of course the zero of the vector space is the zero section \*\*\*\*

**Proposition 2.5.7.** *Let  $L$  be tautological line bundle on  $\mathbb{P}(V)$ . If the projective space  $\mathbb{P}(V)$  has dimension at least 1 then  $\Gamma(\mathbb{P}(V), L) = \{0\}$ .*

*Proof.* Let  $\sigma \in \Gamma(\mathbb{P}(V), L)$ . The composition

$$\mathbb{P}(V) \xrightarrow{\sigma} L \hookrightarrow \mathbb{P}(V) \times V \longrightarrow V \cong \mathbb{K}^{r+1} \tag{2.5.8}$$

is regular and hence constant by Corollary 2.4.8. The unique element in the image is a vector which belongs to every 1 dimensional (vector) subspace of  $V$ . Since  $\dim V \geq 2$  it follows that it is the zero vector.  $\square$

## 2.6 Exercises

Let  $V$  be a  $\mathbb{K}$  vector space of finite dimension, and let  $0 \leq h \leq \dim V$ . The *Grassmannian*

$$\mathrm{Gr}(h, V) := \{W \subset V \mid \dim W = h\}.$$

is the set of subvector spaces of  $V$  of dimension  $h$ . The Zariski topology on  $\mathrm{Gr}(h, V)$  is defined as follows. Let  $\mathrm{Fr}(h, V)$  be the set of ordered lists of linearly independent vectors  $v_1, \dots, v_h \in V$ . We define the left action

$$\begin{aligned} \mathrm{GL}_h(\mathbb{K}) \times \mathrm{Fr}(h, V) &\longrightarrow \mathrm{Fr}(h, V) \\ ((a_{ij}), \{v_1, \dots, v_h\}) &\mapsto \{\sum_{i=1}^h a_{1i}v_i, \sum_{i=1}^h a_{2i}v_i, \dots, \sum_{i=1}^h a_{hi}v_i\} \end{aligned} \quad (2.6.9)$$

The quotient for the equivalence relation defined by the above action is the map

$$\begin{aligned} \mathrm{Fr}(h, V) &\xrightarrow{\pi} \mathrm{Gr}(h, V) \\ v_1, \dots, v_h &\mapsto \mathrm{span}(v_1, \dots, v_h) \end{aligned} \quad (2.6.10)$$

Since  $\mathrm{Fr}(h, V) \subset V^h$  (as an open subset) it inherits a Zariski topology from  $V^h \cong \mathbb{A}^{h \cdot \dim V}$ . The Zariski topology on  $\mathrm{Gr}(h, V)$  is the quotient topology.

**Exercise 2.6.1.** The goal of the exercise is to provide  $\mathrm{Gr}(h, V)$  with the structure of an algebraic variety. Let  $U \subset V$  be a vector subspace of dimension  $\dim V - h$ , i.e. an element of  $\mathrm{Gr}(\dim V - h, V)$ . Let  $\mathrm{Gr}(h, V)_U \subset \mathrm{Gr}(h, V)$  be the subset of  $W$  which are transverse to  $U$ .

- (a) Show that  $\mathrm{Gr}(h, V)_U$  is open.
- (b) Show that the action of  $\mathrm{Hom}(V/U, U)$  on  $\mathrm{Gr}(h, V)_U$  defined by

$$\begin{aligned} \mathrm{Hom}(V/U, U) &\longrightarrow \mathrm{Gr}(h, V)_U \\ (f, W) &\mapsto \{w + \varphi(\bar{w}) \mid w \in W\} \end{aligned} \quad (2.6.11)$$

is simply transitive ( $\bar{w}$  is the equivalence class of  $w$  in  $V/U$ ), and hence it gives a bijection

$$\varphi_U: \mathrm{Hom}(V/U, U) \rightarrow \mathrm{Gr}(h, V)_U. \quad (2.6.12)$$

To be precise there is one such bijection for each choice of  $W \in \mathrm{Gr}(h, V)_U$ , but they are all equivalent for what follows. Show that  $\varphi_U$  is a homeomorphism, and that the collection of  $\mathrm{Gr}(h, V)_U$ 's and homeomorphisms  $\varphi_U$  is an algebraic atlas of  $\mathrm{Gr}(h, V)$ . Thus we have given  $\mathrm{Gr}(h, V)$  the structure of an algebraic prevariety.

- (c) Prove that  $\mathrm{Gr}(h, V)$  is an algebraic variety, i.e. that it is of finite type and separated. (It might help to unwind the definitions above for  $V = \mathbb{K}^n$ , replacing  $\{v_1, \dots, v_h\} \in \mathrm{Fr}(h, V)$  by the  $h \times n$  matrix whose rows are the  $v_i$ 's.)
- (d) Prove that  $\mathrm{Gr}(h, V)$  is irreducible. (Recall that prevarieties of finite type have an irreducible decomposition.)

**Exercise 2.6.2.** The goal of the exercise is to show that  $\mathrm{Gr}(h, V)$ , with the structure of algebraic variety provided by Exercise 2.6.1, is a projective variety.

1. Let  $v_1, \dots, v_a \in V$  be linearly independent, and let  $\alpha \in \bigwedge^h V$ . Prove that  $v_i \wedge \alpha = 0$  for all  $i \in \{1, \dots, a\}$  if and only if there exists  $\beta \in \bigwedge^{h-a} V$  such that  $\alpha = v_1 \wedge \dots \wedge v_a \wedge \beta$ .
2. For  $\alpha \in \bigwedge^h V$ , let  $m_\alpha$  be the linear map

$$\begin{aligned} V &\xrightarrow{m_\alpha} \bigwedge^{h+1} V \\ v &\mapsto v \wedge \alpha \end{aligned}$$

Show that if  $\alpha \neq 0$ , then the kernel of  $m_\alpha$  has dimension at most  $h$ , and that  $\dim \ker(m_\alpha) = h$  if and only if  $\alpha$  is *decomposable*, i.e.  $\alpha = w_1 \wedge \dots \wedge w_h$ , where  $w_1 \wedge \dots \wedge w_h \in V$  are linearly independent.

3. If  $W \in \mathrm{Gr}(h, V)$  then  $\bigwedge^h W$  is a 1-dimensional subspace of  $\bigwedge^h V$ , i.e. a point of  $\mathbb{P}(\bigwedge^h V)$ . Hence we have a well defined *Plücker map*

$$\begin{aligned} \mathrm{Gr}(h, V) &\xrightarrow{\mathcal{P}} \mathbb{P}(\bigwedge^h V) \\ W &\mapsto \bigwedge^h W. \end{aligned}$$

Show that

$$\text{im}(\mathcal{P}) = \left\{ [\alpha] \in \mathbb{P} \left( \bigwedge^h V \right) \mid \dim(\ker m_\alpha) \geq h \right\}, \quad (2.6.13)$$

and if  $[\alpha] \in \text{im}(\mathcal{P})$ , then  $[\alpha] = \bigwedge^h \ker(m_\alpha)$ . Conclude that  $\mathcal{P}$  is injective and that  $\text{im}(\mathcal{P})$  is closed in  $\mathbb{P}(\bigwedge^h V)$ .

4. Prove that the Plücker map defines an isomorphism  $\text{Gr}(h, V) \xrightarrow{\sim} \text{im}(\mathcal{P})$  between algebraic varieties, and hence  $\text{Gr}(h, V)$  is a projective variety.

Let

$$\text{Gr}(k, \mathbb{P}(V)) := \{L \subset \mathbb{P}(V) \mid L \text{ is a linear subspace, } \dim L = k\}. \quad (2.6.14)$$

We have natural identification

$$\begin{array}{ccc} \text{Gr}(k+1, V) & \longrightarrow & \text{Gr}(k, \mathbb{P}(V)) \\ W & \mapsto & \mathbb{P}(W) \end{array} \quad (2.6.15)$$

Thus  $\text{Gr}(k, \mathbb{P}(V))$  is a projective variety.

**Exercise 2.6.3.** Let  $V$  be a 4-dimensional  $\mathbb{K}$  vector space and

$$\text{Gr}(1, \mathbb{P}(V)) \xrightarrow{\mathcal{P}} \mathbb{P}(\bigwedge^2 V) \cong \mathbb{P}^5$$

the Plücker map.

1. Prove that the image of  $\mathcal{P}$  is a non degenerate quadric hypersurface, i.e. that the ideal of  $\text{im } \mathcal{P}$  is generated by a non degenerate quadratic polynomial  $F$ .
2. Let  $X \subset \text{Gr}(1, \mathbb{P}(V))$ . Prove that  $\mathcal{P}(X)$  is a line if and only if  $X$  is a pencil of lines, i.e. the set of lines containing point  $p$  and belonging to a plane  $\Lambda$  containing  $p$ .
3. Let  $X \subset \text{Gr}(1, \mathbb{P}(V))$ . Prove that  $\mathcal{P}(X)$  is a plane if and only if one of the following holds:
  - a)  $X$  is the set of lines containing a point  $p$ .
  - b)  $X$  is the set of lines contained in a plane  $\Lambda$ .

**Exercise 2.6.4.** 1. Let  $X \subset \mathbb{P}^n$  be closed. Given  $0 \leq k \leq n$  let

$$F_k(X) := \{\Lambda \in \text{Gr}(k, \mathbb{P}^n) \mid \Lambda \subset X\}. \quad (2.6.16)$$

Prove that  $F_k(X)$  is a closed subset of  $\text{Gr}(k, \mathbb{P}^n)$ .

2. Let  $X = V(Z_0 Z_3 - Z_1 Z_2) \subset \mathbb{P}^3$  be a non degenerate quadric surface. Describe  $F_1(X) \subset \text{Gr}(1, \mathbb{P}^3) \subset \mathbb{P}^5$ .