

On complex algebraic singularities of some genuinely nonlinear PDEs

Denys Dutykh* and Éric Leichtnam

Abstract. *In this manuscript, we highlight a new phenomenon of complex algebraic singularities formation for solutions of a large class of genuinely nonlinear Partial Differential Equations (PDEs). We start from a unique CAUCHY datum which is holomorphic ramified like $x_1^{\frac{1}{k+1}}$ (and its powers) around the **smooth** locus $x_1 = 0$ and is sufficiently singular. Then, we expect the existence of a solution which should be holomorphic ramified around the **singular** locus \mathfrak{S} defined by the vanishing of the discriminant of an algebraic equation of degree $k + 1$. Notice, moreover, that the monodromy of the CAUCHY datum is **Abelian**, whereas one of the solutions is **non-Abelian** and that \mathfrak{S} depends on the CAUCHY datum in contrast to the LERAY principle (stated for linear problems only). This phenomenon is due to the fact that the PDE is genuinely nonlinear and that the CAUCHY datum is sufficiently singular. First, we investigate the case of the inviscid BURGERS Equation (iBE). Later, we state a general Conjecture 9.2, which describes the expected phenomenon. We view this Conjecture 9.2 as a working programme allowing us to develop interesting new Mathematics. We also state Conjecture 7.1, which is a particular case of the general Conjecture 9.2 but keeps all the flavour and difficulty of the subject. Then, we propose a new algorithm with a map \mathbf{F} such that a fixed point of \mathbf{F} would give a solution to the problem associated with Conjecture 7.1. Then, we perform convincing, elaborate numerical tests which suggest that a BANACH norm should exist for which the mapping \mathbf{F} should be a contraction so that the solution (with the above specific algebraic structure) should be unique. This work is a continuation of [36].*

Preface

In this manuscript, we investigate the process of singularity formation in two examples of fully (or genuinely) nonlinear Partial Differential Equation (PDE)s starting from a single algebraic singular CAUCHY datum. Namely, we consider specific ramified initial value (or CAUCHY-type) problems. In order to better understand the real singularities, our approach consists in looking at what happens in the complex domain. Thus, we formally assume that space and time variables take complex values. The analysis of the PDEs in the complex domain sheds new light on the origin of familiar real singularities. In the case of the inviscid BURGERS Equation (iBE) (with an algebraic singular CAUCHY datum), we are able to carry out all computations analytically using two completely different methods:

2020 Mathematics Subject Classification: 35A21 (primary), 35L03, 35L15 (secondary).

Keywords: nonlinear PDEs, initial value problem, complex singularities, ramification.

*Corresponding author

CAUCHY–KOVALEVSKAYA theory and contact geometry on the space of 1–jets. We propose a generalisation to the case of genuinely nonlinear PDEs of orders two or higher, and we state several conjectures. Moreover, for a second order PDE, we propose an iterative scheme, which allows us to construct efficiently approximate complex solutions. The convergence of this scheme is empirically demonstrated on a sufficient number of numerical examples, which is a good indication in favour of our conjectures. We stress that the proposed algorithm is entirely new. This scheme might also be used to show the existence and uniqueness of solutions to the ramified Initial Value Problem (IVP) corresponding to our conjectures.

In more precise mathematical terms, in this manuscript, we highlight a new phenomenon of complex algebraic singularities formation for solutions of genuinely nonlinear Partial Differential Equation (PDE)

$$P(u)(t, \mathbf{x}) = 0, \quad (t, \mathbf{x}) \in \mathbb{C} \times \mathbb{C}^n$$

starting from a unique CAUCHY datum, which is sufficiently singular. More precisely, we start from a unique CAUCHY datum $u(0, \mathbf{x})$ which is ramified near the origin like $x_1^{\frac{1}{k+1}}$ around the smooth hyper surface $x_1 = 0$ and assume that P is simply characteristic with respect to the co-normal of $x_1 = 0$. Then, we expect the existence of a solution to $P(u) = \underline{0}$ of a specific form (ansatz) $u(t, \mathbf{x}) = \mathcal{A}(t, \mathbf{x}, z(t, \mathbf{x}))$, where \mathcal{A} is a holomorphic function near the origin and $z(t, \mathbf{x})$ is a solution of an algebraic equation

$$z^{k+1}(t, \mathbf{x}) - a_{k-1}(t, \mathbf{x})z^{k-1}(t, \mathbf{x}) - \dots - a_0(t, \mathbf{x}) = 0. \quad (*)$$

Thus, the solution is holomorphic ramified around the singular locus defined by the discriminant (swallow-tail) of Equation (*). Since we search for a solution under a special form which is ramified along a single (singular) hyper-surface, we need only a single CAUCHY datum. Actually, “morally” the solution is unique once we have fixed the choice of a root to the polynomial equation defined by the simple characteristic hypothesis. The fact that the solution is ramified around a singular locus and has a non-ABELIAN monodromy, whereas the CAUCHY datum is ramified around a smooth locus and has ABELIAN monodromy is a (new) phenomenon which is due to the facts that the CAUCHY datum is sufficiently singular and that the PDE $P(u) = \underline{0}$ is genuinely nonlinear. Let us provide two examples to illustrate the concept of genuine nonlinearity, while the more general definition will be given in the formulation of Conjecture 9.2. Let u be a holomorphic function

$$\begin{aligned} u : \mathbb{C}^{n+1} &\longrightarrow \mathbb{C} \\ (t, x_1, x_2, \dots, x_n) &\mapsto u(t, x_1, x_2, \dots, x_n). \end{aligned}$$

Then, the following ‘initial’ value problem is genuinely nonlinear

$$u_{tt} - u_{x_1}(u_{x_1 x_1} + u_{x_2 x_2} + \dots + u_{x_n x_n}) = \underline{0}, \quad u(0, \mathbf{x}) = c_1 x_1 + c_2 x_1^{1+\frac{1}{3}},$$

and the next one is not

$$u_{tt} - u_{x_2}(u_{x_1 x_1} + u_{x_2 x_2} + \dots + u_{x_n x_n}) = \underline{0}, \quad u(0, \mathbf{x}) = c_1 x_1 + c_2 x_1^{1+\frac{1}{3}},$$

where $c_{1,2} \in \mathbb{C}^\times$.

The full generalization of this phenomenon is stated in Conjecture 9.2 for genuinely nonlinear PDE of order greater or equal to two. Simpler versions of this Conjecture are stated in Conjecture 7.1 and Conjecture 7.4. But Conjecture 7.1 keeps all the flavour and the conceptual difficulty of Conjecture 9.2. Conjecture 7.1 considers the following PDE

$$u_{tt} - u_x u_{xx} = \underline{0}$$

along with the CAUCHY datum

$$u(0, x) = \sum_{j=1}^{N_0} c_j x^{1 + \frac{j-1}{3}},$$

where $N_0 \in \mathbb{N}_{>2}$, $c_{1,2} \in \mathbb{C}^\times$ and $\{c_j\}_{j=3}^{N_0} \subseteq \mathbb{C}$ are constants. Moreover, a root of the polynomial equation associated with the simple characteristic hypothesis is chosen. Then, Conjecture 7.1 states the existence of an algebraic solution $u(t, x)$ ramified along a cusp

$$4p^3(t, x) - 27q^2(t, x) = 0$$

depending on the CAUCHY datum and the previous choice. This conjecture is a working programme that allows the development of interesting new mathematics. The algebraic and geometric studies carried out in Sections 5 and 6 allow us to construct, in Section 7, a new algorithm with a map \mathbf{F} such that a fixed point of \mathbf{F} provides a solution to Conjecture 7.1. In Section 8, we carry out various elaborate numerical experiments which show quite convincingly that the iterations $\mathbf{F} \circ m(w_0)$ seem to converge (as $m \rightarrow +\infty$) to a unique fixed point. This suggests that a family of semi-norms should exist, allowing the construction of a BANACH norm for which \mathbf{F} would be contracting. However, this is a very difficult problem, much harder than the one treated successfully in [33] corresponding to the case of algebraic equations of degree two (see also Section 2 for a short reminder).

In the case of the inviscid BURGERS Equation (iBE), we are able to carry out all the computations analytically using two completely different methods: the CAUCHY-KOVALEVSKAYA theorem and the contact geometry approach on the space of 1-jets. This new phenomenon is exhibited very clearly in the case of the iBE. For the most general case of genuinely nonlinear PDEs of the order higher or equal to two, we state the general Conjecture 9.2, which describes the new phenomenon previously mentioned very precisely. Conjecture 9.2 is a very difficult problem. We view it as a motivation to develop new Mathematics. Indeed, in this paper, we try to lay the foundations of proof. We also state Conjectures 7.1 and 7.4, which are particular cases of the general Conjecture 9.2. The Conjecture 7.1 deals with the case $k = 2$ and

$$P(u) := u_{tt} - u_x u_{xx} = \underline{0}, \tag{**1}$$

$$u(0, x) = \sum_{j=1}^{N_0} c_j x^{1 + \frac{j-1}{3}}, \tag{**2}$$

where $N_0 \geq 3$, $c_{1,2} \in \mathbb{C}^\times$ and $c_j \in \mathbb{C}$, ($3 \leq j \leq N_0$) are constants, and $(t, x) \in \mathbb{C}^2$. Morally, Conjecture 7.1 is not much easier than Conjecture 9.2, but its simpler formulation allows for direct numerical verifications.

We revisit and improve the geometric study of the algebra $\mathcal{O}[z]$, where $z^3 = pz + q$ and \mathcal{O} denotes the algebra of germs of holomorphic functions $\{a(p, q)\}$ near the origin $\mathbf{0} \in \mathbb{C}^2$. This study allows us to construct a new algorithm (and the corresponding ansatz) along with a mapping \mathbf{F} such that a fixed point of \mathbf{F} gives a solution to the Problem (**₁), (**₂). We provide various numerical experiments show that the iterations $\mathbf{F}^{\circ m}(w_0)$ seem to converge numerically (when $m \rightarrow +\infty$) to a fixed point which does not depend on w_0 . This provides strong empirical evidence that this algorithm will provide a unique solution to the Problem (**₁), (**₂) of the form $\mathcal{A}(t, \mathbf{x}, z(t, \mathbf{x}))$ once a choice of a root to the simply characteristic equation (of degree two here) has been chosen. It remains a very difficult open problem to find appropriate a BANACH (normed) algebra for which

$$\|\mathbf{F}(w) - \mathbf{F}(v)\| \leq c\|w - v\|,$$

with $0 < c < 1$, which would give a proof of Conjecture 7.1.

Abu Dhabi, UAE
 Saint-Jean-de-Chevelu, France
 June 2024

Denys Dutykh
Eric Leichtnam

1. Introduction

The process of formation of singularities to solutions of nonlinear Partial Differential Equations is a huge and important topic which has stimulated a lot of interesting works (see *e.g.* [47, 54]). We begin by recalling some known results on this topic in order to briefly describe the landscape in which our results and goals will be realized.

Even infinitely smooth initial data under the dynamics of a nonlinear PDE will not remain in general smooth for all times. Henceforth, the topic of the singularity formation in solutions to nonlinear PDEs has been central in the study of these equations. Perhaps the wave-breaking phenomenon is the most familiar and far from being completely studied, *natural* singularity formation processes [4]. Moreover, everyone can easily see the wave-breaking process.

An attempt to classify various *finite time* singularities in PDEs has been made in [12]. Physical (and several other) applications have clearly motivated this classification. The question of eventual blow-up in 3D incompressible EULER and NAVIER–STOKES equations is open and central to many current theoretical and numerical investigations [23, 30, 40, 46]. We also refer to [8] as an excellent review of available numerical approaches to detect complex singularities of PDEs. Complex singularities of the LORENZ dynamical system (with complex time) have been studied in [52].

The present manuscript is devoted in the first place to the study of complex algebraic singularities in the inviscid BURGERS Equation (iBE). Namely, we study the algebraic singularity blow-up under the dynamics of some first and second-order genuinely nonlinear PDEs. Among the second-order PDEs, we focus on a particular second-order PDE belonging to the celebrated family of p -systems described in some detail below.

The iBE can be written as [7]:

$$u_t - uu_x = \underline{0}, \quad (1.1)$$

where in the (classical) real case

$$\begin{aligned} u : \mathbb{R}_{\geq t_0} \times \mathbb{R} &\longrightarrow \mathbb{R}, \\ (t, x) &\mapsto u(t, x). \end{aligned}$$

The subscripts $(-)_t$ and $(-)_x$ denote the usual partial derivatives $\frac{\partial}{\partial t}$ and $\frac{\partial}{\partial x}$ respectively. In order to obtain an Initial Value Problem (IVP), Equation (1.1) has to be completed by an appropriate initial condition:

$$u(t_0, x) = u_0(x), \quad x \in \mathbb{R}.$$

It is well-known that the iBE will develop the gradient catastrophe in finite time from a generic initial condition. More precisely, the existence time $t_0 \leq t < t_s$ depends on the initial condition:

$$t_s = \frac{1}{\sup_{x \in \mathbb{R}} u_{0,x}(x)}.$$

Please note that the real singularity never occurs (in forward time) if $u_{0,x}$ takes strictly negative values for all finite $x \in \mathbb{R}$. Below, for the sake of simplicity, we shall take $t_0 = 0$.

The real singularity is point-like in the sense that it occurs locally at a single point and at a fixed time t_s . In contrast to real singularities, we shall demonstrate below that complex algebraic singularities take place on sets of positive dimensions (*i.e.* analytical hyper-surfaces in \mathbb{C}^2 in our case). For the sake of completeness, we would like to mention that some singularities may even happen on sets of fractional HAUSDORFF dimensions [13]. This mechanism is conjectured for the NAVIER–STOKES equations in 3D. We can also remember an old idea of J. LERAY that singularities in the NAVIER–STOKES equations could be related to the phenomenon of turbulence [5]. However, this idea was not followed, up to now and to the best of our knowledge, by any significant results.

In a similar line of thinking, we shall consider also in the present manuscript the following second-order nonlinear hyperbolic equation:

$$u_{tt} - u_x u_{xx} = \underline{0}, \quad (1.2)$$

whose IVP requires traditionally two data to be specified:

$$u(0, x) = u_0(x), \quad x \in \mathbb{R}, \quad (1.3)$$

$$u_t(0, x) = u_1(x), \quad x \in \mathbb{R}. \quad (1.4)$$

However, we shall consider a ramified CAUCHY problem with only one (algebraic) CAUCHY datum (1.3) and we shall seek a solution $(t, x) \mapsto u(t, x)$ of a very special (algebraic) form so that only the first CAUCHY datum is needed. Both considered Equations (1.1) and (1.2) come from the same family of PDEs discussed in Section 3. Mathematically speaking, we discuss in the present study the hyperbolic sub-family of genuinely nonlinear PDEs. The general notion of genuinely nonlinear PDE that we use is stated as Assumption (2) in Conjecture 9.2. For example, the following PDE

$$u_{tt} - u_x u_{xx} = \underline{0}$$

is genuinely nonlinear, whereas

$$u_{tt} - uu_{xx} = \underline{0}$$

is not genuinely nonlinear but only quasilinear. Other examples will be given in Section 3.

Example 1.1. Let us consider the following CAUCHY ‘toy’ problem in \mathbb{C}^2 :

$$\begin{aligned} \frac{\partial u}{\partial t} &= u^2, \\ u(0, x) &= \frac{1}{x}. \end{aligned}$$

It is not difficult to see that the exact solution to this problem is

$$u(t, x) = \frac{1}{x - t}.$$

This simple example shows that it is absolutely crucial to assume that CAUCHY data (1.3) is bounded if we want to establish the existence of a holomorphic solution ramified along a characteristic hyper-surface. However, we mention that this toy example is not representative of our study for the following reasons:

1. The singular locus $\mathfrak{S} := \{t \in \mathbb{C} \mid (t, t)\}$ is a regular hyper-surface.
2. The hyper-surface \mathfrak{S} is not characteristic for the differential operator $\frac{\partial}{\partial t}$.

Henceforth, our example has some serious limitations.

It is well-known that the behaviour of the real-valued and complex-valued solutions is quite different. We shall mention the example of the 2D viscid BURGERS system. For real-valued solutions O. LADYZHENSKAYA proved¹ in 1963 global existence and uniqueness result in SOBOLEV spaces [29]. Later, it turned out that complex-valued solutions differ drastically from real ones. In particular, LI and SINAI proved in [37] using the renormalization group method that complex-valued solutions (in n D) can develop finite time singularities. This was proven earlier in [45] for gradient-like solutions. The 2D complex BURGERS equation was studied in [39], where an open set of a six-parameter family of initial conditions is constructed such that the corresponding solutions exhibit blowups in finite time. Finally, complex-valued initial conditions have been showing *numerically* to develop a singularity in finite time in complete agreement with theory. These results have also been extended to 3D complex-valued NAVIER–STOKES equations [38]. The complex singularities in 2D EULER equations have been studied in [19].

Our study differs at least in two important respects from the previously described line of thinking:

1. We consider hyperbolic and inviscid equations starting with the iBE.
2. We complexify not only the dependent variables but also the independent ones.

The viscous BURGERS equation with real-time and complex spatial variables has been considered in [17] in the context of semi-groups. Similarly, the linear wave (or D’ALEMBERT) and telegraph equations were treated in [16]. Perhaps the complex view on PDEs can be traced back to the work of S. KOVALEVSKAYA who considered the CAUCHY problem (CP) for the usual (linear) heat equation with complex temporal and spatial variables [28]. In particular, she showed that there are examples of holomorphic non-entire initial conditions such that the power series solution does not converge in any neighbourhood of $t_0 \in \mathbb{C}$. The BOREL

¹This result is a kind of folklore theorem because the original reference [29] does not contain a very detailed proof.

summability of these divergent solutions has been studied much later [41]. However, our study's goal is to shed new light from the complex geometry angle on the formation of real singularities in fully nonlinear PDEs. The complex iBE appeared surprisingly also in random matrix theory [43] and random surface models [27].

In the present work, we focus first on the classical iBE and the following ramified IVP:

$$\begin{aligned} u_t - uu_x &= \underline{0}, \\ u(0, x) &= x^{\frac{1}{3}}. \end{aligned}$$

We show that the last IVP possesses solutions of the form $u = a + z$, where a is a regular holomorphic function and z verifies the cubic equation $z^3 = pz + q$. The obtained solution can be analytically continued along any path in a small neighbourhood of the origin $\mathbf{0} \in \mathbb{C}^2$ and originating from $(0, x_0) \in \mathbb{C}^2$ and avoiding the cusp singularity $4p^3 - 27q^2 = \underline{0}$. This is explained in detail in Section 4. Thus, the solution u is ramified along the *singular* locus $4p^3 - 27q^2 = \underline{0}$ and its monodromy group² (cf. Remarks 4.3 and 4.4) is the (non-commutative) permutation group S_3 (cf. Remark 4.3) whereas the CAUCHY datum $x^{\frac{1}{3}}$ is ramified around a *smooth* locus with a commutative monodromy group, namely $\mathbb{Z}/3\mathbb{Z}$. This phenomenon (including the changing of the monodromy group) is due both to the genuine nonlinearity of the iBE and the fact the algebraic CAUCHY datum is sufficiently singular (see Section 4.1 and Theorem 6.4).

However, the main goal of our study is to formulate a convincing numerical convergence evidence along with the generalisation of this phenomenon for genuinely nonlinear PDEs of the order two and higher (see Conjectures 7.1, 7.4 and 9.2).

A first generalization is stated in Conjecture 7.1 which considers the Partial Differential Equation (PDE) (1.2) with the initial condition

$$u(0, x) = \sum_{j=1}^{N_0} c_j x^{1 + \frac{j-1}{3}},$$

²Informally speaking, the monodromy group is a group of transformations that encodes what happens with data when we turn around the singularity. More formally, a monodromy is a representation of the fundamental group. Let \mathbf{V} be a holomorphic connected variety. Let us take $x_0 \in \mathbf{V}$ and u be a holomorphic germ defined at point x_0 , which can be prolonged along any closed loop issued from x_0 . If γ is any such closed loop at x_0 , by $\gamma \cdot u$ we denote the result of this prolongation. Finally, by \mathbb{E} , we denote the vector space generated by all such elements $\gamma \cdot u$. If $\dim \mathbb{E} < +\infty$, we say that u is of *finite determination*. The homotopy equivalence classes $[\gamma]$ form the fundamental group $\pi_1(\mathbf{V})$. Finally, the monodromy is defined as the following map:

$$\begin{aligned} \pi_1(\mathbf{V}) &\longrightarrow \mathbf{GL}(\mathbb{E}), \\ \gamma &\mapsto (u \mapsto \gamma \cdot u). \end{aligned}$$

Of course, it has to be shown that all these maps are well-defined, and the result does not depend on the representative γ of the homotopy class $[\gamma]$. Fortunately, it can be done without any difficulties.

where $N_0 \in \mathbb{N}_{>2}$, $c_{1,2} \in \mathbb{C}^\times$ and $\{c_j\}_{j=3}^{N_0} \subseteq \mathbb{C}$ are constants. It states the existence of an algebraic solution $u(t, x)$ ramified along a cusp $4p^3(t, x) - 27q^2(t, x) = 0$ depending on the CAUCHY datum. We strongly believe that Conjectures 7.1, 7.4 and the most general Conjecture 9.2 are important problems requiring the development of interesting new Mathematics.

This manuscript is the first step in which we shall provide solid mathematical foundations and numerical evidence to believe that they are true. Especially regarding the Conjecture 7.1, we shall construct in Section 7 a new (and highly non-trivial) algorithm, allowing us to obtain quite convincing numerical tests supporting the convergence of this new algorithm in Section 8. Moreover, we sketch a tentative strategy to theoretically address the general Conjecture 9.2. In Sections 5 and 6, we revisit the general results of [36] in the specific case of the cubic equation $z^3 = pz + q$ and make them more precise. The big advantage here is that most computations can be done almost explicitly, which allows us to highlight the underlying structures, which will allow us to construct a new practical algorithm from Section 7. More precisely, in Section 5 we revisit the ring³ $\mathcal{O}[[z]]$ (with $z^3 = pz + q$) and we introduce the primitive operator ∂_q^{-1} . The reason why the derivation ∂_q operator governs the structure of the ring $\mathcal{O}[[z]]$ is elucidated in Section 6. In the same Section, we establish the fact that (z, z^2) is a solution of a holonomic \mathcal{D} -module with the characteristic variety \mathbf{V} which is included in the union of the zero section and the co-normal⁴ to the cusp $4p^3 - 27q^2 = 0$:

$$\mathbf{V} \cap \mathbf{T}_0^* \mathbb{C}^2 = (0, 0; 0, \mathbb{C}).$$

In fact, using a deep result of KASHIWARA, we are able to prove an even stronger result stating that the characteristic variety \mathbf{V} is the union of the zero section and the co-normal to the cusp. The characteristic variety is an important geometric invariant of a \mathcal{D} -module. The integrability of the characteristic variety is a central result in the theory of \mathcal{D} -modules. Loosely speaking, this result says that the singular support of a \mathcal{D} -module is an involutive sub-variety in the co-tangent bundle. The involutivity property may be seen under the sheaf angle: the ideal sheaf defining the singular support is closed with respect to the natural POISSON bracket on the co-tangent bundle [50].

Let us remind you of some additional relevant background material that will help you understand the sequel of this section. We denote by \mathcal{D} the sheaf of holomorphic differential operators on $\mathbb{C}^2 \cong \{(p, q)\}$. The vector space of sections of \mathcal{D} over an open subset \mathcal{U} is denoted by $\mathcal{D}(\mathcal{U})$.

Definition 1.2. Let M be a sheaf of \mathcal{D} -modules (\mathcal{D} -module for short) over an open subset \mathcal{U} of \mathbb{C}^2 . The \mathcal{D} -module \mathcal{M} is said to be coherent if for any sheaf

³Throughout this article, the word “ring” means “ring with identity”.

⁴Let us remind that the co-normal is constituted of all co-tangent vectors which annihilate on the tangent space to a given manifold (here, an analytic curve or surface). In the case of a non-smooth manifold or variety \mathbf{V} , the co-normal of \mathbf{V} is defined as the closure of the co-normal of \mathbf{V}^{reg} , where \mathbf{V}^{reg} designates the set of regular points of \mathbf{V} .

homomorphism of \mathcal{D} -modules

$$\phi : \mathcal{D}^m \longrightarrow \mathcal{M}$$

and for any point z of \mathcal{U} , one can find an open neighbourhood \mathcal{U}_z of z and a finite number of sections s_1, s_2, \dots, s_r (over \mathcal{U}_z) of the sheaf $(\ker \phi)$ such that for any open subset $\mathcal{W} \subseteq \mathcal{U}_z$, the restrictions $s_1|_{\mathcal{W}}, s_2|_{\mathcal{W}}, \dots, s_r|_{\mathcal{W}}$ generate $(\ker \phi)(\mathcal{W})$ as a module over $\mathcal{D}(\mathcal{W})$.

The fact that above the origin, there is only one line (the co-normal to ∂_q) in \mathbf{V} is absolutely crucial for our constructions. In particular, it explains why the primitive ∂_q^{-1} exists and why the “micro-local singularities”⁵ of the product z^2 remain under the control above the origin $\mathbf{0} \in \mathbb{C}^2$. This observation will constitute one of the key ingredients in the future proof of Conjecture 7.1. However, before engaging in this huge endeavour, we thought obtaining convincing numerical tests to converge our new iterative algorithm was indispensable. Indeed, its convergence was not obvious *a priori*, and its proof will constitute a big challenge. We end with a short heuristic explanation of why we believe that if we impose a second CAUCHY datum to Conjecture 7.1, then there will be no solution in general. Indeed, from the linear result [36], two cusps will show up. The products of u and its derivatives will provide a spreading of singularities (associated with the co-normal of each of the cusps) over the origin, which will not be possible to control. We shall return to this question in the forthcoming works.

The present manuscript is organized as follows. In Section 2, we briefly review some results of [33], which proves Conjecture 9.2 in the case of the algebraic equation of degree two and which provides some foundational material for an approach of Conjectures 7.1 and 7.4. In Section 3, we review an interesting family of genuinely nonlinear PDE to which our Conjectures should apply. First, we understand the formation of complex singularities in the iBE using the CAUCHY-KOVALEVSKAYA theory in Section 4.2, then, for the sake of completeness, we obtain the same results using completely different methods of contact geometry in Section 4.3. The reason for tackling the same problem from two completely different angles is that it allows us to explore different ideas underlying the general conjecture formulated below. Some brief reminder on the ring of formal series with holomorphic germs⁶ coefficients are made in Section 5, and some remarks on holonomic \mathcal{D} -modules are presented in Section 6. The second order nonlinear PDE (1.2) is analyzed in Section 7, and a completely new algorithm to approximate ramified solutions is presented in Section 7.1. The convergence of this algorithm would prove a particular case of the general Conjecture 9.2. We demonstrate numerically in Section 8 that the proposed algorithm converges in practice, which constitutes one

⁵We take this expression in the quotes because the distributions z and z^2 are not yet well-defined in the real case *stricto sensu*. It will be a topic of our future works.

⁶The notion of a germ of a mathematical object captures the local properties of that object. The germ is precisely the equivalence class of objects that share the same local property. In order to implement the idea of germs, the space has to be at least topological to give sense to the word ‘local’. The name ‘germ’ was introduced into Mathematics in the continuation of the *sheaf* metaphor.

of the main achievements of the present work. Some possible further generalizations are discussed in Section 9. This study’s main conclusions and perspectives are outlined in Section 10. Finally, in Appendix C, we show how our theory relates to the classical theory of shock waves in an elementary example, see also the end of Subsection 4.3.1.

2. Review of existing results

In this Section, we remind the general theory of singular solutions to genuinely nonlinear PDEs, which was initiated by the second author some thirty years ago, and the construction of holomorphic ramified solutions that have been considered even earlier, see *e.g.* [53, 22, 24]. Certain linear ramified CAUCHY problems were considered in [34] and second-order semi-linear problems in [35]. The linear ramified differential operators were considered in [10] in the setting of sheaves theory. They demonstrated independently a weaker version of the results presented in [36]. HAMADA considered the homogeneous linear case even earlier in [21]. The micro-local existence result was established in [6] for certain classes of linear differential operators. The same homogeneous case was investigated later using the contact transformation approach in [26]. The linear homogeneous case was also considered using the tools of the micro-local theory of sheaves in [11].

The reminders will allow us to understand better the context and set up of Conjectures 7.1, 7.4 and 9.2. Actually, Theorem 2.8 (to be recalled below) is the particular case of Conjectures 7.1 and 9.2 obtained by restricting oneself to algebraic equations of the second order. This is an additional good reason in favour of Conjecture 7.1.

Let us consider an operator \mathcal{P} of the order m :

$$u \mapsto \mathcal{P}(u) \stackrel{\text{def}}{=} \sum_{|\sigma|=m} P_\sigma(\mathbf{x}, u, \partial_{\mathbf{x}}^\beta u) \partial_{\mathbf{x}}^\sigma u + \mathcal{R}(\mathbf{x}, \partial_{\mathbf{x}}^\beta u)$$

where we use the multi-indices⁷ σ and β with $|\beta| \leq m - 1$. The use of multi-index notation is often attributed in the literature to Laurent SCHWARTZ. All functions P_σ and \mathcal{R} are supposed to be holomorphic in their arguments $\forall \mathbf{x}$ in some vicinity $\mathcal{U}_{\mathbf{0}}$ of $\mathbf{0} \in \mathbb{C}^{n+1}$, $\forall \partial_{\mathbf{x}}^\beta u$ in some vicinity of $u_0^\beta \stackrel{\text{def}}{=} \partial_{\mathbf{x}}^\beta u(\mathbf{0})$, $\beta \in \mathbb{N}^{n+1}$. The question of local solutions to equation $\mathcal{P}u = \underline{0}$ can be naturally

⁷If $\beta = (\beta_0, \beta_1, \dots, \beta_n) \in \mathbb{Z}^{n+1}$ be a multi-index. By $|\cdot|$ we denote the *height* of the multi-index defined as

$$|\beta| \stackrel{\text{def}}{=} \sum_{j=0}^n |\beta_j|.$$

The derivation operator with respect to the variable x_j , $j \in (n + 1)^\square$ is denoted by ∂_{x_j} . If $\beta \in \mathbb{N}^{n+1}$ is a multi-index of derivation, then we shall write:

$$\partial^\beta \stackrel{\text{def}}{=} \partial_{x_0}^{\beta_0} \circ \partial_{x_1}^{\beta_1} \circ \dots \circ \partial_{x_n}^{\beta_n}.$$

asked. The solution $u(\mathbf{x})$ is holomorphic and ramified around a hyper-surface \mathfrak{S} defined by equation $s(\mathbf{x}) = 0$ and passing by $\mathbf{0} \in \mathbb{C}^{n+1}$. This solution has the following analytical form [33]:

$$u(\mathbf{x}) = a(\mathbf{x}) + \sum_{k=0}^{+\infty} b_k(\mathbf{x}) s^{\gamma_k}(\mathbf{x}), \quad (2.1)$$

where the sequence $\{\gamma_k\}_{k=0}^{+\infty}$ takes its values in $\mathbb{R}_{>0}$ and is strictly increasing and tending to $+\infty$. The functions a and $b_k, \forall k \in \mathbb{N}$ are holomorphic in $\mathcal{U}_{\mathbf{0}} \ni \mathbf{0}$. The series (2.1) is convergent on the covering space of $\mathcal{U}_{\mathbf{0}} \setminus \mathfrak{S}$. The following assumption is also adopted: the hyper-surface \mathfrak{S} is simply characteristic for the linearized equation \mathcal{P}_{lin} of the genuinely nonlinear operator \mathcal{P} in every point $a(\mathbf{x})$. In other words, the principal symbol p_m of \mathcal{P}

$$p_m(\mathbf{x}, \bar{\xi}) \stackrel{\text{def}}{=} \sum_{|\sigma|=m} P_{\sigma}(\mathbf{x}, a(\mathbf{x}), \partial_{\mathbf{x}}^{\beta} a(\mathbf{x})) \bar{\xi}^{\sigma}, \quad |\beta| \leq m - 1,$$

verifies two conditions:

- $p_m(\mathbf{x}, ds(\mathbf{x}))|_{\mathfrak{S}} \equiv 0$,
- $\partial_{\bar{\xi}} p_m(\mathbf{0}, ds(\mathbf{0}))|_{\mathfrak{S}} \neq 0$.

The last condition means that \mathfrak{S} is simply characteristic for \mathcal{P}_{lin} . This condition may be viewed as a holomorphic analogue of the condition for a real differential operator to be strictly hyperbolic with respect to a real hypersurface containing the CAUCHY data.

We also suppose that we know the ‘initial’ data $\partial_t^{\beta} a|_{t=0} (\forall |\beta| \leq m - 1)$, $b_k|_{t=0} (\forall k \geq 0)$ and $s|_{t=0}$, where $t = 0$ is the equation a complex smooth hyper-surface \mathcal{T} transversal to the vector field $\partial_{\bar{\xi}} p_m(\mathbf{x}, ds(\mathbf{x}))$. Below, we consider two separate cases.

2.1. Weakly singular solutions

In this case, we take a holomorphic function a and a hyper-surface \mathfrak{S} (given by equation $s(\mathbf{x}) = 0$) verifying preceding conditions, we assume $\mathcal{P}(a) \equiv \underline{0}$ and we seek for weakly singular solutions of type (2.1) with

$$\gamma_k := m + (k + 1)\mu, \quad \mu \in]0, 1[$$

so that the constructed solution u admits m bounded derivatives. A change of variables allows us to assume that \mathfrak{S} is defined by the equation

$$s(\mathbf{x}) = y_1 = 0,$$

where we decompose $\mathbf{x} \equiv (t, \mathbf{y})$ and we introduce the following notation:

$$1 - \frac{\mathbf{y}}{R} \stackrel{\text{def}}{=} \prod_{j=1}^n \left(1 - \frac{y_j}{R}\right), \quad R > 0.$$

By analogy,

$$\frac{1}{1 - \frac{\mathbf{y}}{R}} \stackrel{\text{def}}{=} \prod_{j=1}^n \frac{1}{1 - \frac{y_j}{R}}, \quad R > 0.$$

Definition 2.1. Let $u = \sum_{j=1}^{+\infty} u_j y^j$ and $v = \sum_{j=1}^{+\infty} v_j y^j$ be two formal power series with complex coefficients. We shall say that $u \preceq v$ if and only if

$$|u_j| \leq |v_j|, \quad \forall j \in \mathbb{N}.$$

The last definition can be easily extended to the multi-variable case:

Definition 2.2. Let $p \in \mathbb{N}^\times$ and $u = \sum_{\beta \in \mathbb{N}^p} u_\beta \mathbf{y}^\beta$ and $v = \sum_{\beta \in \mathbb{N}^p} v_\beta \mathbf{y}^\beta$ be two formal power series with complex coefficients. We shall say that $u \preceq v$ if and only if

$$|u_\beta| \leq |v_\beta|, \quad \forall \beta \in \mathbb{N}^p.$$

Definition 2.3. Let Υ be a formal variable. Given $R > 0, \varepsilon > 0, p \in \mathbb{N}$ and $d \in \mathbb{N}$, we denote by $\mathbb{A}^d(R, \varepsilon, p)$ the set of formal power series of the type:

$$u = u(\mathbf{x}, \Upsilon) = \sum_{j=0}^{+\infty} \sum_{k=1}^{+\infty} u_{k,j}(t, \mathbf{y}) \Upsilon^{k\mu+j+d}, \quad 0 < \mu < 1,$$

where functions $u_{k,j}$ are holomorphic in some neighbourhood of $\mathbf{0} \in \mathbb{C}^{n+1}$ and verify

$$u_{k,j} \preceq \sum_{l=0}^{+\infty} \frac{t^l}{\left(1 - \frac{\mathbf{y}}{R}\right)^{k+j+l}} \cdot \frac{(k+j+l)!}{l!(k+j+p)!} \cdot C_{kjl}^p \quad (2.2)$$

such that

$$\|u\|_{p,d} \stackrel{\text{def}}{=} \sum_{k,j,l \in \mathbb{N}} C_{kjl}^p \varepsilon^{k+2j+l} < +\infty, \quad (2.3)$$

where $C_{kjl}^p(u)$ are the smallest non-negative real constants allowing to bound functions $u_{k,j}$ in (2.2). We also introduce the following related definitions:

$$\mathbb{A}^d(R) \stackrel{\text{def}}{=} \bigcup_{\varepsilon > 0} \mathbb{A}^d(R, \varepsilon, p), \quad \mathbb{A}^d \stackrel{\text{def}}{=} \bigcup_{R > 0} \mathbb{A}^d(R).$$

To each element $u \in \mathbb{A}^d(R, \varepsilon, p)$ we put non-injectively in correspondence a function

$$u(\mathbf{x}) = \sum_{j=0}^{+\infty} \sum_{k=1}^{+\infty} u_{k,j} y_1^{k\mu+j+d}.$$

It is not difficult to see that if $u_1 \in \mathbb{A}^{d_1}$ and $u_2 \in \mathbb{A}^{d_2}$, then $u_1 \cdot u_2 \in \mathbb{A}^{d_1+d_2}$. If β is a multi-index in \mathbb{N}^{n+1} of the height $\leq d$ then the operator ∂^β takes the elements from \mathbb{A}^d and sends them to $\mathbb{A}^{d-|\beta|}$. Since \mathfrak{S} is characteristic, then \mathcal{P}_{lin} sends \mathbb{A}^d to \mathbb{A}^{d-m+1} . The following property specifies the geometric domain of the convergence associated with the norm $\|\cdot\|_{p,d}$ defined in (2.3):

Lemma 2.4 ([33]). *Let $u \in \mathbb{A}^d(R, \varepsilon, p)$. Then, every $u_{k,j}$ is a holomorphic function in the domain*

$$\Omega \stackrel{\text{def}}{=} \left\{ (t, \mathbf{y}) \in \mathbb{C}^{n+1} \mid |y_j| < R, \quad |t| < \varepsilon \prod_{j=1}^n \left(1 - \frac{|y_j|}{R} \right) \right\},$$

and, for every compact $K \subset \Omega$ there exists a constant $C_K > 0$ such that

$$\sup_K |u_{k,j}| \leq C_K^{k+j+1}.$$

Conversely, let $\{u_{k,j}\}$ be a sequence of holomorphic functions defined on an open neighbourhood $\mathcal{U}_{\mathbf{0}}$ of the point $\mathbf{0} \in \mathbb{C}^{n+1}$ such that

$$\sup_{\mathcal{U}_{\mathbf{0}}} |u_{k,j}| \leq C^{k+j+1}.$$

Then,

$$\sum_{j=0}^{+\infty} \sum_{k=1}^{+\infty} u_{k,j} \Upsilon^{k\mu+j+d} \in \mathbb{A}^d.$$

Then, finally, we may state the first result:

Theorem 2.5 ([33]). *Take $u_0 \in \mathbb{A}^m$ and let \mathcal{J} be an analytic hyper-surface passing by $\mathbf{0} \in \mathbb{C}^{n+1}$ being defined by the analytical equation $t = 0$. This hyper-surface is transversal to the field $\partial_{\bar{\xi}} p_m(\mathbf{x}, dy_1)$ (and, thus, to \mathfrak{S}). Then, there exists $r \in \mathbb{A}^m(\mathbb{C}^{n+1})$ such that*

$$\mathcal{P}(a(\mathbf{x}) + r(\mathbf{x})) \equiv \underline{0} \quad \text{and} \quad r|_{t=0} = u_0.$$

The proof consists, roughly speaking, of applying the fixed point iterations.

Remark 2.6. The obtained solution $a(\mathbf{x}) + r(\mathbf{x})$ in the previous Theorem is ramified around a smooth locus (independent of the value of μ) because the CAUCHY datum is not sufficiently singular.

2.2. Strongly singular solutions

In this Section we are looking for solutions of type (2.1) with

$$\gamma_k \stackrel{\text{def}}{=} m - \frac{1}{2} + \frac{k}{2}.$$

We stress out that the function a is not necessarily a solution of equation $\mathcal{P}(a) = \underline{0}$ anymore if $b_0 \neq \underline{0}$. In this Section, a and \mathfrak{S} will be considered as problem unknowns on the same footing with $\{b_k\}_{k=0}^{+\infty}$.

Definition 2.7. Let S be a formal variable. By $\mathbb{B}^{m-1}(\mathbb{C}^{n+1})$ we designate the algebra of formal series of the form:

$$r(\mathbf{x}, S) = \sum_{k=0}^{+\infty} b_k(\mathbf{x}) S^{\frac{k+2m-1}{2}}.$$

We assume also that all $\{b_k\}_{k \in \mathbb{N}}$ are holomorphic functions defined on the same neighbourhood \mathcal{U}_0 of the point $\mathbf{0} \in \mathbb{C}^{n+1}$ and there exists a constant $C > 0$ such that

$$|b_k(\mathbf{x})| \leq C^{k+1}, \quad \forall k \in \mathbb{N}, \quad \forall \mathbf{x} \in \mathcal{U}_0.$$

Below, we shall substitute the formal variable S by the holomorphic function s vanishing at the origin $\mathbf{0} \in \mathbb{C}^{n+1}$ and defining the characteristic hyper-surface for the operator \mathcal{P}_{lin} .

For any $|\beta| = 1$, we define the action of ∂^β on \mathbb{B}^{m-1} by conveying that

$$\partial^\beta (S^{\frac{k+2m-1}{2}}) = \frac{k+2m-1}{2} S^{\frac{k+2m-3}{2}} \partial^\beta s(\mathbf{x}).$$

The action of ∂^β on \mathbb{B}^{m-1} is defined recursively for any $|\beta| \leq m$. To each element $r(\mathbf{x}, S) \in \mathbb{B}^{m-1}(\mathbb{C}^{n+1})$ we associate non-injectively a function $\mathbf{x} \mapsto r(\mathbf{x}, s(\mathbf{x}))$ defined near the origin.

Let $\mathbf{x} \equiv (t, \mathbf{y}) \in \mathbb{C} \times \mathbb{C}^n$ and we assume that the operator \mathcal{P} has the form $\mathcal{P} = \partial_t^m + \mathcal{Q}(t, \mathbf{y}, \partial^\beta)$ with $|\beta| \leq m$ and $\beta \neq (m, 0, \dots, 0)$. The principal symbol of the linearized operator $\mathcal{P}_{\text{lin}}(a)$ of \mathcal{P} at a has the form $p_m(t, \mathbf{y}, \partial^\beta \tau, \bar{\xi})$ with $|\beta| \leq m - 1$. Finally, we consider a point $(\tau_0, \bar{\xi}^0) \in (\mathbb{C} \times \mathbb{C}^n) \setminus \mathbf{0}$ together with CAUCHY data $\{a_j(\mathbf{y})\}_{j=0}^{m-1}$ holomorphic in some neighbourhood of $\mathbf{0} \in \mathbb{C}^n$. We assume that

$$p_m(0, \partial^\sigma a_j(\mathbf{0}), \tau^0, \bar{\xi}^0) = \underline{0}, \quad \partial_\tau p_m(0, \partial^\sigma a_j(\mathbf{0}), \tau^0, \bar{\xi}^0) \neq \underline{0},$$

where $j + |\sigma| \leq m - 1$. We choose also a root $\tau_1(\mathbf{x}, u_\beta, \bar{\xi})$ of the equation $\tau \mapsto p_m(\mathbf{x}, u, \tau, \bar{\xi}) = 0$ holomorphic near the point $(0, \partial^\sigma a_j(\mathbf{0}), \bar{\xi}^0)$. Finally, let s_1 be a holomorphic function near $\mathbf{0} \in \mathbb{C}^n$ such that

$$s_1(\mathbf{0}) = 0, \quad \text{d} s_1(\mathbf{0}) = \bar{\xi}^0.$$

We can state the main result:

Theorem 2.8 ([33]). *Let $\sum_{k=0}^{+\infty} b_k^0(\mathbf{y}) S^{\frac{k+2m-1}{2}} \in \mathbb{B}^{m-1}(\mathbb{C}^n)$. Then, there exists a neighbourhood of the point $\mathbf{0} \in \mathbb{C}^{n+1}$ along with holomorphic functions $s, a, \{b_k\}_{k=0}^{+\infty}$ defined on it such that s is characteristic for \mathcal{P}_{lin} and*

$$\begin{aligned} b_k(0, \mathbf{y}) &\equiv b_k^0, \quad \forall k \in \mathbb{N}, \\ \partial_t^j a(0, \mathbf{y}) &\equiv a_j, \quad j \in (m+1)^\square, \\ s(0, \mathbf{y}) &\equiv s_1, \\ \partial_t s(0) &\equiv \tau^0. \end{aligned}$$

The series

$$r(\mathbf{x}, S) \stackrel{\text{def}}{=} \sum_{k=0}^{+\infty} b_k(\mathbf{x}) S^{\frac{k+2m-1}{2}} \in \mathbb{B}^{m-1}$$

and $\mathbf{x} \mapsto u(\mathbf{x}) \equiv a(\mathbf{x}) + r(\mathbf{x}, s(\mathbf{x}))$ is a solution of the equation $\mathcal{P}(u) = \underline{0}$.

The space $\mathbb{B}^{m-1}(\mathbb{C}^{n+1})$ is endowed with appropriate norm to define a BANACH algebra. This Theorem is proved by applying the fixed point PICARD iteration, which also gives a practical algorithm converging to the required solution [35].

Remark 2.9. Since the CAUCHY datum $u_0 \equiv u(0, \mathbf{y})$ is sufficiently singular, the hyper-surface $s = \underline{0}$ around which the solution is ramified depends on u_0 and cannot be prescribed in advance. Moreover, since u_0 is ramified like $s_1^{\frac{1}{2}}$, the algebraic equation $z^2 + bz + c = \underline{0}$ will inevitably show up in the construction of the solution. The discriminant of this equation is equal to $\Delta := b^2 - 4c$. Observe that $\Delta = \underline{0}$ defines a smooth locus with respect to the variables (b, c) . That is why Theorem 2.8 is easier to prove than its generalizations to the algebraic equations of order $k \geq 3$, cf. Conjectures 7.1, 7.4 and 9.2.

3. An interesting family of genuinely nonlinear PDEs

We consider a family of PDEs with the unknown function $u : \mathbb{R} \times \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}$:

$$\mathcal{L}_m(u) \stackrel{\text{def}}{=} \partial_t^m u - \partial_x^{m-1} u \partial_x^m u = \underline{0}, \quad (3.1)$$

where $m \geq 1$ is an integer parameter and $\partial_{(-)}^m$ denotes the m^{th} order partial derivatives with respect to independent variables t or x . Let us introduce the auxiliary variables:

$$v_1(t, x) \stackrel{\text{def}}{=} \partial_x^{m-1} u(t, x), \quad v_m(t, x) \stackrel{\text{def}}{=} \partial_t^{m-1} u(t, x).$$

Then, we have

$$\begin{aligned} \partial_t^{m-1} v_1 - \partial_x^{m-1} v_m &= \underline{0}, \\ \partial_t v_m - \partial_x \left(\frac{1}{2} v_1^2 \right) &= \underline{0}. \end{aligned}$$

We shall also introduce the functions $\{v_j\}_{j=2}^{m-1}$ satisfying the following system:

$$\begin{aligned} \partial_t v_1 - \partial_x v_2 &= \underline{0}, \\ \partial_t v_2 - \partial_x v_3 &= \underline{0}, \\ &\vdots \\ \partial_t v_m - \partial_x \left(\frac{1}{2} v_1^2 \right) &= \underline{0}. \end{aligned}$$

For regular solutions (*i.e.* of the class C^1), the last system can be rewritten under the following matrix form:

$$\partial_t \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_m \end{pmatrix} - \begin{pmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ v_1 & 0 & 0 & \cdots & 0 \end{pmatrix} \cdot \partial_x \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_m \end{pmatrix} = \mathbf{0}, \quad (3.2)$$

where $\mathbf{0}$ is a function taking a constant (zero) vector value, *i.e.*

$$\begin{aligned} \mathbf{0} : \mathbb{R}_{\geq 0} \times \mathbb{R} &\longrightarrow \mathbb{R}^m, \\ (t, x) &\mapsto (0, 0, \dots, 0)^\top. \end{aligned}$$

Formulations (3.1) and (3.2) are equivalent to each other on the set of sufficiently regular solutions.

It is not difficult to see that this system matrix is nothing else but the transposed companion matrix of the polynomial

$$\lambda^m - v_1 \in \mathbb{R}[\lambda].$$

If we assume that $v_1 > 0$ (just for definiteness), the eigenvalues of this matrix are all simple and given by:

$$\lambda_k \stackrel{\text{def}}{:=} v_1^{\frac{1}{m}} e^{i \frac{2\pi k}{m}}, \quad k \in m^\square.$$

The eigenvector associated with the eigenvalue λ_k is a direction in the kernel of the following linear operator:

$$\begin{pmatrix} -\lambda_k & 1 & 0 & \cdots & 0 \\ 0 & -\lambda_k & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ v_1 & 0 & 0 & \cdots & -\lambda_k \end{pmatrix}$$

or simply the following vector:

$$\omega_k \stackrel{\text{def}}{:=} (1, \lambda_k, \dots, \lambda_k^{m-1})^\top$$

We can easily verify that

$$\langle d\lambda_k, \omega_k \rangle = \frac{\lambda_k}{m v_1} \neq 0,$$

which means that the k^{th} characteristic field is *genuinely nonlinear* according to the definition from [31, Section 5]. It is also obvious that System (3.2) is not hyperbolic since it admits complex characteristic speeds if $m > 2$. Below, we shall consider in some detail the two hyperbolic cases $m \in \{1, 2\}$. The

former corresponds to the familiar iBE. The latter deserves more attention. It can be written as the following system of two equations:

$$\partial_t v_1 - \partial_x v_2 = \underline{0}, \quad (3.3)$$

$$\partial_t v_2 - \partial_x (\sigma(v_1)) = \underline{0}. \quad (3.4)$$

In the last system (also known as the p -system), we can recognize a particular case of nonlinear elasticity written in LAGRANGIAN variables, where v_1 is the deformations field and v_2 corresponds to the velocity field. Finally, the mechanical constraint $\sigma(w) \stackrel{\text{def}}{=} \frac{1}{2} w^2$ is induced by the deformation v_1 . See [9, Chapter 7] and, in particular, Equation (7.1.11) therein for more details. It is known that System (3.3) is hyperbolic when $\frac{d\sigma}{dw} \equiv w > 0$, which becomes the condition $v_1 > 0$ in our case. The two characteristic fields corresponding to speeds

$$\lambda_{\pm} = \pm\sqrt{v_1}$$

and to eigenvectors

$$\omega_{\pm} = (1, \pm\sqrt{v_1})^T$$

are genuinely nonlinear as in the general case (*i.e.* $m > 2$).

4. Inviscid Burgers equation

In this Section we consider a particular IVP for the iBE (1.1):

$$u_t - u u_x = \underline{0}, \quad (4.1a)$$

$$u(0, x) = x^{\frac{1}{3}}. \quad (4.1b)$$

The main particularity of our study is that we consider the space and time variables to be complex, including the unknown function u :

$$u : \mathbb{C} \times \mathbb{C} \longrightarrow \mathbb{C}.$$

The derivatives with respect to x and t are understood from now on in the sense of complex analysis. The initial condition containing an algebraic branching point in the complex domain was taken on purpose. Consequently, the IVP (4.1) is called the *ramified CAUCHY problem*. Solving the ramified CAUCHY problem (4.1) means by definition to find an open neighbourhood $\mathcal{U}_{\mathbf{0}}$ of $\mathbf{0} \in \mathbb{C}^2$ along with a germ of the holomorphic function $u(t, x)$ at a point of $\mathcal{U}_{\mathbf{0}} \cap \{t = 0\}$ (different from the origin) satisfying (4.1a) and (4.1b) such that $u(t, x)$ can be continued holomorphically along any path in $\mathcal{U}_{\mathbf{0}}$ which does not meet a certain characteristic hyper-surface which restricts to $x = 0$ when $t = 0$. Below, we describe two different approaches to study the IVP (4.1) because both of them contain the foundational ideas to approach the Conjecture 7.1.

4.1. A linear digression

It is often helpful to consider first the linear problem before tackling the fully nonlinear formulation. It often brings some useful insight. The linear counterpart to IVP (4.1) reads

$$\begin{aligned}u_t - u_x &= 0, \\u(0, x) &= x^{\frac{1}{3}}.\end{aligned}$$

It is clear to see that the last IVP has the following *exact* solution:

$$u(t, x) = (t + x)^{\frac{1}{3}}.$$

Moreover, the singular locus of this solution is the following affine algebraic variety:

$$\mathfrak{S}_s \stackrel{\text{def}}{=} \{(t, x) \in \mathbb{C}^2 \mid t + x = 0\}.$$

It is clear to see that this locus is completely regular as a geometric object. This is a characteristic property of linear equations. Below in Section 4.2, we shall demonstrate that in the genuinely **nonlinear** case (4.1), the corresponding singular locus is **singular** itself, and we shall describe its algebraic singularity explicitly. We emphasize that Example 1.1 was (simply) nonlinear and also had a completely regular locus. Also, the GALOIS group of the field extension of the CAUCHY data and of the solution is the same ABELIAN group:

$$\mathbf{Gal}(\mathbb{C}(x^{\frac{1}{3}}) : \mathbb{C}(x)) = U_3.$$

This observation is to be compared with the genuinely nonlinear case described below. Moreover, the solution $u(t, x)$ and the CAUCHY datum $u(0, x)$ share the same monodromy group.

Let us underline one more aspect in which linear and genuinely nonlinear problems differ. If we remove from \mathbb{C}^2 the singular locus and we compute the fundamental group of the obtained domain, we shall obtain:⁸

$$\pi_1(\mathbb{C}^2 \setminus \mathfrak{S}_s) = \mathbb{Z},$$

which is an ABELIAN group. In the genuinely nonlinear case, the fundamental group will be non-commutative.

⁸Indeed, this result is not difficult to obtain

$$\begin{aligned}\pi_1(\mathbb{C}^2 \setminus \mathfrak{S}_s) &= \pi_1(\mathbb{C}^2 \setminus \{x = 0 \mid (t, x) \in \mathbb{C}^2\}) = \pi_1(\mathbb{C} \times \mathbb{C} \setminus \{0\}) \\ &= \pi_1(\mathbb{C} \setminus \{0\}) = \mathbb{Z}.\end{aligned}$$

4.2. The Cauchy–Kovalevskaya approach

In the present section, we employ the CAUCHY–KOVALEVSKAYA theory in order to study the Initial Value Problem (IVP) (4.1). We shall even obtain a slightly more general result by considering the following IVP:

$$u_t - uu_x = \underline{0}, \quad (4.2a)$$

$$u(0, x) = a_0(x) + x^{\frac{1}{3}}. \quad (4.2b)$$

Theorem 4.1. *Consider the IVP (4.1) for the iBE. Then, there exist three holomorphic functions $(t, x) \mapsto a(t, x)$, $(t, x) \mapsto p(t, x)$ and $(t, x) \mapsto q(t, x)$ defined in some neighbourhood of the point $(0, 0) \in \mathbb{C}^2$ such that*

$$a(0, x) = a_0(x), \quad p(0, x) = 0, \quad q(0, x) = x \quad (4.3)$$

and

$$u(t, x) \stackrel{\text{def}}{=} a(t, x) + z(t, x) \quad (4.4)$$

being a unique solution to Equation (4.1) with the complex auxiliary function $(t, x) \mapsto z(t, x)$ verifying the following algebraic relation⁹

$$z^3 = pz + q. \quad (4.5)$$

Proof. First of all, we check that the ansatz (4.4) verifies the initial condition (4.2). Indeed, by taking the limit $t \rightarrow 0$ in Equation (4.5), we obtain

$$z^3(0, x) = p(0, x) \cdot z(0, x) + q(0, x) \stackrel{(4.3)}{\equiv} x.$$

Thus, $z(0, x) = x^{\frac{1}{3}}$. Substituting this result into the solution ansatz (4.4), we obtain:

$$u(0, x) := a(0, x) + z(0, x) \stackrel{(4.3)}{\equiv} a_0(x) + x^{\frac{1}{3}}.$$

By differentiating the algebraic relation (4.5) with respect to t and x , we may compute the partial derivatives of the function z :

$$z_t = \frac{p_t z + q_t}{3z^2 - p}, \quad z_x = \frac{p_x z + q_x}{3z^2 - p}. \quad (4.6)$$

Let us substitute the solution ansatz (4.4) into the iBE (4.1a):

$$a_t + z_t - (a + z)(a_x + z_x) = \underline{0}.$$

⁹This cubic relation can be beneficially seen as an algebraic equation $f(z) := z^3 - pz - q = 0$ over some field \mathbb{k} of characteristic different from 2 and 3. We would like to mention here that the GALOIS group associated with this equation depends on the fact whether its discriminant $4p^3 - 27q^2$ is a square or not in \mathbb{k} . In the former case, $\mathbf{Gal}_{\mathbb{k}}(f) = \mathbb{Z}/3\mathbb{Z}$ and in the latter case $\mathbf{Gal}_{\mathbb{k}}(f) = S_3$, i.e. the symmetric group over a finite set of 3 symbols. Since we are working over the field of complex numbers $\mathbb{k} := \mathbb{C}(p, q)$, the GALOIS group is the latter.

By substituting partial derivatives (4.6) and multiplying both sides by the denominator $3z^2 - p$, we obtain:

$$(3z^2 - p)a_t + zp_t + q_t - (a + z)(a_x(3z^2 - p) + zp_x + q_x) = \underline{0}.$$

Now, by equating the coefficients in front of equal powers of z , we have the following sequence of equalities:

$$\begin{aligned} z^0 : & \quad -pa_t + q_t + \cancel{apa_x} - aq_x = \underline{0}, \\ z^1 : & \quad p_t + \cancel{pa_x} - ap_x - q_x = \underline{0}, \\ z^2 : & \quad a_t - \cancel{aa_x} - \frac{1}{3}p_x = \underline{0}, \\ z^3 : & \quad 3a_x = \underline{0}. \end{aligned}$$

The last equation can be substituted into three previous ones to obtain:

$$\begin{aligned} q_t &= aq_x + pa_t \equiv aq_x + \frac{1}{3}pp_x, \\ p_t &= ap_x + q_x, \\ a_t &= \frac{1}{3}p_x. \end{aligned} \tag{4.7}$$

The first two equations are not incompatible because co-tangent vectors dp and dq are linearly independent by continuity in the vicinity of the point $(0, 0) \in \mathbb{C}^2$. Indeed,

$$(p_t, p_x)(0, 0) \equiv (1, 0), \quad (q_t, q_x)(0, 0) \equiv (a_0(0), 1).$$

It is not difficult to see that the last two co-vectors are linearly independent for any value of $a_0(0) \in \mathbb{C}$. Thus, after applying the classical CAUCHY-KOVALEVSKAYA theorem to System (4.7), we obtain the required local existence and uniqueness result. \square

Corollary 4.2. *The local existence and uniqueness for the IVP (4.1) is obtained by simply choosing $a_0 := \underline{0}$.*

Remark 4.3. It is instructive to have a look at the obtained solution from the point of view of GALOIS theory. We already established above that the GALOIS group associated with the initial data is the ABELIAN group U_3 . The dynamics of the iBE transforms this initial data into the local analytic solution (4.4). The GALOIS group associated to the solution u can be easily computed:

$$\mathbf{Gal}(\mathbb{C}(p, q)[z] : \mathbb{C}(p, q)) = S_3.$$

We stress that the last group is obviously different from U_3 and, additionally, it is non-commutative.

Investigating the situations where the solution established in the last Theorem 4.1 becomes singular is also very interesting. Let us fix some values of (p_0, q_0) of $(p, q) \in \mathbb{C}^2$ for which z_0 is a simple root of Equation (4.5). Then,

by the classical implicit function theorem, there exists a holomorphic function $(p, q) \mapsto z(p, q)$ defined in the vicinity of the point (p_0, q_0) of the complex plane. Henceforth, the singular locus for Equation (4.5) will consist of the following points:

$$\Delta \stackrel{\text{def}}{:=} \{ (p, q) \in \mathbb{C}^2 \mid \text{Equation (4.5) has a double root} \}.$$

It is not difficult to describe this set analytically. For this, we eliminate z from the following system of equations:

$$\begin{cases} z^3 - pz - q = 0, \\ 3z^2 - p = 0. \end{cases}$$

The last two equations can be seen as a system of linear equations with respect to parameters (p, q) . It admits the following parametric solution:

$$\begin{cases} q = -2z^3, \\ p = 3z^2. \end{cases}$$

Henceforth, the singular locus is

$$\Delta := \{ (p, q) \in \mathbb{C}^2 \mid 4p^3 - 27q^2 = 0 \}.$$

It is not difficult to recognize here the swallow tail (cusp) singularity [55], which coincides with the cusp in this particular case ($n = 2$, see below).

Remark 4.4. The Theorem 4.1 establishes the local existence of the solution in some polydisc D around the origin $(0, 0) \in \mathbb{C}^2$. Let us take the solution germ $u(t, x)$ at $(t_0, x_0) \in D \setminus \Delta^<(\{\mathbf{0}\})$. If γ denotes a closed loop around the point (t_0, x_0) which does not intersect the singular locus $\Delta^<(\{\mathbf{0}\})$, then one can obtain the analytical continuation of the germ $u(t, x)$ along the loop γ [14]. In other terms, this defines the action of the first homotopy (or fundamental) group $\pi_1(\mathcal{U}_0 \setminus \Delta^<(\{\mathbf{0}\}))$ on the solution (4.4):

$$\gamma \cdot u := a + \gamma \cdot z,$$

since the regular part a is not ramified. We note that this group action turns out to be *transitive* (see below Theorem 4.8). We say that $\gamma \cdot z$ is the monodromy action of γ on z . Depending on the chosen loop $\gamma \in D \setminus \Delta^<(\{\mathbf{0}\})$, this action may bring us to a different root of the algebraic Equation (4.5). This situation is schematically depicted in Figure 1. We also mention that this action can be lifted to the universal covering.

It is well-known (see *e.g.* [1]) that the fundamental group¹⁰ of the complex

¹⁰Both Authors of this manuscript are deeply convinced that the mathematical reality is connected. We have some reasons to think that there are some deep hidden connections between the problem considered in our study and the number theory. Indeed, it is known that the braid group \mathbf{Br}_3 is the universal central extension of the modular group $\mathbf{PSL}_2(\mathbb{Z}) \stackrel{\text{def}}{:=} \mathbf{SL}_2(\mathbb{Z})/Z(\mathbf{SL}_2(\mathbb{Z})) \equiv \mathbf{SL}_2(\mathbb{Z})/\{\pm\mathbf{1}\}$ (here by $\mathbf{1}$ we understand the unit element of $\mathbf{SL}_2(\mathbb{Z})$). The last modular group is deeply rooted in the theory of modular forms and newforms in the number theory. However, at the current stage, we are unable to make this statement more precise.




Figure 1: A schematic illustration for the analytical continuation of a solution germ defined at $x_0 \in D$ along a closed path $\gamma \subseteq D$.

plane without the cusp is

$$\pi_1(\mathbb{C}^2 \setminus \Delta^<(\{0\})) = \mathbf{Br}_3,$$

the braid group with three strands is an infinite non-commutative group (in contrast to the linear case discussed above).

Let us clarify our terminology. Consider the following polynomial equation:¹¹

$$F(\mathbf{x}, z) := z^{n+1} - x_n z^{n-1} - x_{k-1} z^{n-2} - \dots - x_2 z - x_1 = \underline{0}, \quad (4.8)$$

where $\mathbf{x} \stackrel{\text{def}}{=} (x_1, x_2, \dots, x_n) \in \mathbb{C}^n$. By $\Delta(\mathbf{x})$ we denote the discriminant of the polynomial Equation (4.8) [18]. By swallow tail we shall designate the analytical hyper-surface $\Delta^<(\{0\})$. We provide below the most general definition of the swallow tail which suits our purposes:

Definition 4.5 ([36]). An analytical hyper-surface \mathfrak{S} in \mathbb{C}^{n+1} defined in some open neighbourhood \mathcal{U}_0 is called the swallow tail with the edge $\mathbf{0} \in \mathbb{C}^{n+1}$ if there exists $k \in \mathbb{N}_{\geq 1}$ and k holomorphic functions

$$g_j : \mathcal{U}_0 \rightarrow \mathbb{C}, \quad g_j(\mathbf{0}) = 0, \quad j \in k^\square,$$

such that the differentials $\{dg_j(\mathbf{0})\}_{j=1}^k$ are linearly independent and \mathfrak{S} is defined as the locus of the equation

$$\Delta(g_1(\mathbf{x}), g_2(\mathbf{x}), \dots, g_k(\mathbf{x})) = \underline{0},$$

¹¹We would like to mention that the GALOIS group of this polynomial equation is the symmetric group S_{n+1} over a finite set of $n + 1$ symbols, which is the most non-commutative finite group, in the sense that the centre of $Z(S_n)$ is trivial $\forall n \geq 3$. The point is that if the underlying GALOIS group is non-commutative, then the analytical prolongation of the solution will be non-commutative as well, *i.e.* the solution value will depend on the order of loops starting and returning to a given point.

where, as above, $\Delta(g_1, g_2, \dots, g_k)$ denotes the discriminant of the following polynomial equation in z [18]:

$$z^{k+1} - g_k z^{k-1} - \dots - g_2 z - g_1 = \underline{0}.$$

We define the co-normal $\mathcal{N}(\mathfrak{S})$ of \mathfrak{S} to be the closure in $\mathbf{T}^* \mathcal{U}_0 \setminus \mathbf{0}$ of the co-normal $\mathcal{N}(\mathfrak{S}_{\text{reg}})$, where $\mathfrak{S}_{\text{reg}}$ is the smooth (regular) part of \mathfrak{S} , *i.e.*

$$\mathfrak{S}_{\text{reg}} \stackrel{\text{def}}{=} \{ \mathbf{y} \in \mathbb{C}^k \mid \Delta(\mathbf{y}) = 0, \quad d\Delta(\mathbf{y}) \neq 0 \}.$$

It can be also shown that for any polydisc D centered around $\mathbf{0} \in \mathbb{C}^k$, $D \cap \mathfrak{S}_{\text{reg}}$ is a connected domain. This property is important from the topological point of view.

Remark 4.6. Let $\mathbf{x}_0 \in \mathcal{U}_0 \setminus \mathfrak{S}$. Consider a holomorphic germ¹² at the point $(g_1^{(0)}, g_2^{(0)}, \dots, g_n^{(0)}) := (g_1(\mathbf{x}_0), g_2(\mathbf{x}_0), \dots, g_n(\mathbf{x}_0))$:

$$(g_1, g_2, \dots, g_n) \mapsto z(g_1, g_2, \dots, g_n),$$

where z is the solution to the polynomial Equation (4.8). The classical implicit function theorem ensures the existence of this germ. The germ $z(g_1, g_2, \dots, g_n)$ is a holomorphic function ramified around $\Delta^<(\{0\})$ and it can be prolonged holomorphically along any path starting from the point $(g_1(\mathbf{x}_0), g_2(\mathbf{x}_0), \dots, g_n(\mathbf{x}_0))$ and avoiding the swallow tail $\Delta^<(\{0\})$. Then, the germ in \mathbf{x}_0 :

$$\mathbf{x} \mapsto z(g_1(\mathbf{x}), g_2(\mathbf{x}), \dots, g_n(\mathbf{x}))$$

can be prolonged holomorphically along any path starting from \mathbf{x}_0 and belonging to $\mathcal{U}_0 \setminus \mathfrak{S}$. These facts are rigorously established in [36, Theorem 3.2]:

Theorem 4.7. *Let $\{z_j\}_{j=1}^{n+1}$ denote the set of solutions to polynomial Equation (4.8). Then, for $\forall j \in (n+1)^\square$, the germ z_j defined at \mathbf{x}_0 can be holomorphically prolonged along any path starting from \mathbf{x}_0 and belonging to $\mathbb{C}^n \setminus \Delta^<(\{0\})$.*

¹²Consider a point x of a complex topological manifold \mathbb{V} . Consider also two maps $f, g : \mathcal{U}_x^{(f), (g)} \rightarrow \mathbb{C}$ defined on some neighbourhoods (not necessarily the same) of the point $x \in \mathbb{V}$. These functions define the same germ (or they belong to the same equivalence class) if there is an open neighbourhood $\mathcal{U} \subseteq \mathcal{U}_x^{(f)} \cap \mathcal{U}_x^{(g)}$ such that

$$f(u) = g(u), \quad \forall u \in \mathcal{U}.$$

This fact can be denoted as $f \sim_x g$ and the germ generated by the function f is defined as

$$[f]_x \stackrel{\text{def}}{=} \{ g : \mathcal{U}_x \rightarrow \mathbb{C} \mid f \sim_x g \}.$$

The set of germs at a point x can be obviously endowed with the ring structure over the field \mathbb{C} . Here, we discussed the scalar case. The generalization to the vectorial case is straightforward. We would like to mention that the set of germs does not possess a non-trivial topology. Henceforth, it makes little sense to speak of the convergence of sequences of germs.

Proof. See [36, p. 40]. □

Basically, we came to the point where we may define a group epimorphism from the fundamental group on the permutation group of simple roots to the algebraic Equation (4.8):

Theorem 4.8 ([36]). *Let D be a polydisc of some positive radius centered around $\mathbf{0} \in \mathbb{C}^n$ and γ is a closed path drawn in $D \setminus \Delta^<(\{\mathbf{0}\})$ starting and ending at \mathbf{x}_0 :*

$$\begin{aligned} [0, 1] &\longrightarrow D \setminus \Delta^<(\{\mathbf{0}\}), \\ t &\mapsto \gamma(t) \end{aligned}$$

such that $\gamma(0) = \gamma(1) \equiv \mathbf{x}_0$. By $\gamma \cdot z(\mathbf{x}_0)$ we denote the holomorphic germ obtained from $z(\mathbf{x}_0)$ after the analytical continuation along the loop γ . The implicit function theorem and the fact that the roots of (4.8) are simple in $D \setminus \Delta^<(\{\mathbf{0}\})$ allow us to define the mapping:

$$\begin{aligned} S_{n+1} &\longrightarrow S_{n+1} \\ z(\mathbf{x}_0) &\mapsto \gamma \cdot z(\mathbf{x}_0), \end{aligned}$$

where S_{n+1} is the permutation group on $n + 1$ symbols $\{z_1, z_2, \dots, z_{n+1}\}$. Then, the following mapping

$$\begin{aligned} \pi_1(D \setminus \Delta^<(\{\mathbf{0}\})) &\longrightarrow S_{n+1}, \\ \gamma &\mapsto (z_j(\mathbf{x}_0) \mapsto \gamma z_j(\mathbf{x}_0)) \end{aligned}$$

defines the desired group epimorphism.

4.2.1. Uniformization of the local solution

It is known that the CAUCHY–KOVALEVSKAYA theory provides only the *local* analytic solutions. However, we work in a complex analytic setting where a germ can be continued along any path, avoiding singularities. Henceforth, the local solution obtained in the previous Section can be continued analytically along any path starting in some neighbourhood in origin and avoiding the swallow tail singularity $\Delta^<(\{\mathbf{0}\})$. The analytical continuation theorem ensures that two solution germs at a point x obtained from two *homotopic* paths starting at x_0 are identical. This fundamental theorem of the analytical continuation guarantees that the *uniform* solution is well-defined in each homotopy class of paths [14].

We shall use this analytical continuation technique in order to lift the obtained local solution to the global one on the universal covering. Let us denote the universal covering¹³ of $\mathcal{X} \stackrel{\text{def}}{=} \mathbb{C}^2 \setminus \Delta^<(\{\mathbf{0}\})$ by \mathcal{Y} . In the previous Section, we already mentioned that the fundamental group of \mathcal{X} is \mathbf{Br}_3 and the GALOIS

¹³Let \mathcal{X} be a connected topological manifold. We shall say that a topological manifold \mathcal{Y} is a covering of \mathcal{X} if there exists a surjection $\pi : \mathcal{X} \rightarrow \mathcal{Y}$ such that $\forall x \in \mathcal{X}, \exists \mathcal{U}_x$ an open

group associated to Equation (4.5) is S_3 . It is known from the theory of groups that there exists a group epimorphism¹⁴

$$\chi : \mathbf{Br}_3 \twoheadrightarrow S_3.$$

It is also a well-known fact that $\ker \chi \trianglelefteq \mathbf{Br}_3$ is a normal sub-group and $\mathbf{Br}_3 / \ker \chi \cong S_3$. Moreover, for every $\gamma \in \ker \chi$, the monodromy action of $\chi(\gamma)$ fixes each solution z of (4.5). We introduce also an intermediate covering $\mathcal{R} \stackrel{\text{def}}{=} \mathcal{Y} / \ker \chi$. It is composed of \mathcal{Y} elements orbits by the (sub-)group action $\ker \chi$. It is also the RIEMANN surface which defines the GALOIS covering of the structural automorphisms group (*i.e.* $\ker \chi$). This situation is represented in the following commutative diagram:

$$\begin{array}{ccc} \pi_1(\mathcal{X}) & \longrightarrow & \mathcal{Y} \\ & & \searrow \pi \\ & & \mathcal{R} \\ & \swarrow \pi_\chi & \\ & \mathcal{X} & \end{array}$$

Hence, the solution germ u on \mathcal{X} induces a uniform holomorphic function $u \circ \pi_\chi$ on the intermediate covering \mathcal{R} . In a similar way, the solution germ u can be lifted to the universal covering \mathcal{Y} by taking the composition $u \circ \pi$. We added a little illustration of the uniformization process in Figure 2. This completes our short description of the solution uniformization.

4.3. The contact geometry approach

In this Section, we recover the same swallow tail singularity using the contact geometry approach. It can also be seen as a holomorphic method of characteristics.

Consider a complex space¹⁵

$$\mathbb{C}^5 \equiv \mathbb{C}^{2+2+1} \stackrel{\text{def}}{=} \{(t, x, u, \tau, \xi) \in \mathbb{C}^5\},$$

together with a holomorphic contact form

$$\omega \stackrel{\text{def}}{=} du - \tau dt - \xi dx,$$

neighbourhood with the property that $\pi^{-1}(\{\mathcal{U}_x\}) = \cup_{i \in I} V_i$ is open in \mathcal{Y} and $\forall i \in I$ the maps $\pi_i : V_i \rightarrow \mathcal{U}_x$ are diffeomorphisms. A covering is said to be the *universal covering* if \mathcal{Y} is also simply connected (we also remind that a space is called ‘simply connected’ if any closed loop is homotopic to a point). The equivalence classes of homotopic paths basically give universal covering. The term ‘the universal covering’ is not an abuse of the language because all universal coverings of space are isomorphic to each other.

¹⁴An epimorphism is simply a surjection subject to some additional properties.

¹⁵We may think of it as a 1-jet space of a complex function $(t, x) \mapsto u(t, x)$.




Figure 2: *Illustration of the solution uniformization on the universal covering \mathcal{Y} .*

which defines the contact structure on \mathbb{C}^5 . Let us also consider a first-order holomorphic PDE:

$$\mathcal{F}(t, x, u, u_t, u_x) \equiv \mathcal{F}(t, x, u, \tau, \xi) = \underline{0}.$$

Typically, the practical application we are interested in is given by the inviscid BURGERS Equation (iBE):

$$\mathcal{F}(t, x, u, \tau, \xi) := \tau - u\xi = \underline{0}. \quad (4.9)$$

It is assumed that the family $\{d\mathcal{F}, \omega\}$ is free¹⁶ and \mathcal{F} is a real smooth function of its arguments.

The contact field $\mathcal{H}_{\mathcal{F}}$ is defined by the following condition [2]:

$$d\omega(\mathcal{H}_{\mathcal{F}}, h) \equiv d\mathcal{F} \cdot h, \quad \forall h \in \ker \omega.$$

In our case, the contact field can be easily computed explicitly [32, Chapter V, Section §1]:

$$\mathcal{H}_{\mathcal{F}} := \left(\frac{\partial \mathcal{F}}{\partial \tau}, \frac{\partial \mathcal{F}}{\partial \xi}, \tau \frac{\partial \mathcal{F}}{\partial \tau} + \xi \frac{\partial \mathcal{F}}{\partial \xi}, \right. \quad (4.10)$$

$$\left. - \left(\frac{\partial \mathcal{F}}{\partial t} + \tau \frac{\partial \mathcal{F}}{\partial u} \right), - \left(\frac{\partial \mathcal{F}}{\partial x} + \xi \frac{\partial \mathcal{F}}{\partial u} \right) \right). \quad (4.11)$$

By ϕ^t we denote the flow map along the vector field $\mathcal{H}_{\mathcal{F}}$:

$$\begin{cases} \frac{d}{dt} \phi^t(z) \Big|_{t=0} = \mathcal{H}_{\mathcal{F}}(z), \\ \phi^0(z) = z. \end{cases}$$

¹⁶In the contrary case, the contact field $\mathcal{H}_{\mathcal{F}}$ will be identically equal to zero.

The integral curves of this flow are called the *characteristics*. By construction, the function \mathcal{F} is constant along this flow. We can easily check it by direct computation:

$$\begin{aligned} \frac{d}{dt} \left(\mathcal{F} \left(\phi^t(z) \right) \right) &\stackrel{(4.10)}{=} \mathcal{F}_t \mathcal{F}_\tau + \mathcal{F}_x \mathcal{F}_\xi + \mathcal{F}_u (\tau \mathcal{F}_\tau + \xi \mathcal{F}_\xi) \\ &\quad + \mathcal{F}_\tau (-\mathcal{F}_t - \tau \mathcal{F}_u) + \mathcal{F}_\xi (-\mathcal{F}_x - \xi \mathcal{F}_u) \equiv \underline{0}. \end{aligned}$$

Obviously, the last identity is the consequence of the fact that

$$d\mathcal{F} \cdot \mathcal{H}_{\mathcal{F}} := d\omega(\mathcal{H}_{\mathcal{F}}, \mathcal{H}_{\mathcal{F}}) = \underline{0}.$$

It is not difficult to check that $\mathcal{H}_{\mathcal{F}} \in \ker \omega$.

We recall that the LEGENDRIAN sub-variety \mathbb{V} in our case is a holomorphic, possibly singular, integral sub-variety of $\ker \omega$ of dimension $\dim \mathbb{V} = 2$ such that

$$\begin{aligned} \mathbf{T}\mathbb{V} &\subseteq \ker \omega, \\ \mathbb{V} &\subseteq \mathcal{F}^{-1}(0). \end{aligned} \tag{4.12}$$

We shall say that \mathbb{V} is the generalized solution in the sense of LIE. We know that a contact transformation allows obtaining a classical solution locally given by the function graph. If the canonical projection

$$\begin{aligned} \pi : \mathbb{V} &\longrightarrow \mathbb{C}^2, \\ (t, x, u, \tau, \xi) &\mapsto (t, x) \end{aligned}$$

is of (maximal) rank 2, then \mathbb{V} is a 1-jet $j^1(u)$ of a complex holomorphic function u , *i.e.*

$$\mathbb{V} \equiv \left\{ (t, x) \in \mathbb{C}^2 \mid (t, x, u(t, x), u_t(t, x), u_x(t, x)) \right\}.$$

We remind that all considerations in this article are local in the vicinity of $(0, 0) \in \mathbb{C}^2$. The LEGENDRIAN \mathbb{V} should be understood in the same sense.

The LEGENDRIAN \mathbb{V} can be also defined by a *generating function* [2], *i.e.* $(\tau, x) \mapsto \mathcal{S}(\tau, x)$:

$$\mathbb{V} = \left\{ (\tau, x) \in \mathbb{C}^2 \mid (-\mathcal{S}_\tau, \tau, \mathcal{S} - \tau \mathcal{S}_\tau, x, \mathcal{S}_x) \right\}.$$

It is not difficult to check that conditions (4.12) hold. For example,

$$d(\mathcal{S} - \tau \mathcal{S}_\tau) - \tau d(-\mathcal{S}_\tau) - \mathcal{S}_x dx = d\mathcal{S} - \mathcal{S}_\tau d\tau - \mathcal{S}_x dx \equiv \underline{0}.$$

4.3.1. Application to the iBE

In this case, the first order holomorphic PDE was defined above in Equation (4.9). In this case, the contact field becomes:

$$\begin{aligned} \mathcal{H}_{\mathcal{F}} &:= \left(1, -u, \underbrace{\tau - u\xi}_{\equiv \underline{0} \text{ by (4.9)}}, \tau \xi, \xi^2 \right). \end{aligned} \tag{4.13}$$

Using the iBE (4.9) and CAUCHY data (4.2), we may reconstruct the 1-jet of the function u at $t = 0$:

$$j^1(u) \cap \{t = 0\} \equiv \left(0, x, x^{\frac{1}{3}}, \frac{1}{3} x^{-\frac{1}{3}}, \frac{1}{3} x^{-\frac{2}{3}} \right) \stackrel{\text{def}}{=} z_0.$$

The last point z_0 will be used as an initial condition in finding the flow $\phi^s(z_0)$ (we change the flow parameter t to s , because t is already employed in the iBE). To simplify computations, we notice also that the third component in $\mathcal{H}_{\mathcal{F}}$ vanishes (cf. Equation (4.13)) since \mathcal{F} conserves its value along the flow trajectories:

$$\begin{cases} \frac{d}{dt} \phi^s(z_0) \Big|_{s=0} = \mathcal{H}_{\mathcal{F}}(z_0) \equiv (1, -u, 0, \tau \xi, \xi^2), \\ \phi^0(z_0) = z_0. \end{cases}$$

The flow can be explicitly written in this case:

$$\begin{aligned} \phi^s(z_0) &\equiv (t(s), x(s), u(s), \tau(s), \xi(s)) \\ &:= \left(s, x - x^{\frac{1}{3}} s, x^{\frac{1}{3}}, \frac{x^{\frac{1}{3}}}{3x^{\frac{2}{3}} - s}, \frac{1}{3x^{\frac{2}{3}} - s} \right). \end{aligned}$$

Hence, $u \equiv x^{\frac{1}{3}}$. We introduce a new variable

$$y \stackrel{\text{def}}{=} x - x^{\frac{1}{3}} t \equiv u^3 - ut.$$

In these new variables, the flow becomes:

$$\phi^s(z_0) \equiv \left(t, y, u, \frac{u}{3u^2 - t}, \frac{1}{3u^2 - t} \right),$$

with an additional algebraic relation:

$$u^3 - ut - y = 0. \quad (4.14)$$

The singularity occurs when $t \rightarrow 3u^2$. It is not difficult to see that by definition of variable y , and we have in the limit $y \rightarrow -2u^3$. After eliminating u , we obtain an important algebraic relation between t and y on the singularity:

$$4t^3 - 27y^2 = 0. \quad (4.15)$$

It corresponds to the swallow tail (cusp) singularity seen above in Section 4.2.

Going back to Equation (4.14), we can write:

$$u^3 - ut - y = (u - z)(u^2 + zu - t + z^2), \quad (4.16)$$

$$y = z(z^2 - t), \quad (4.17)$$

where z is a chosen zero of the polynomial (4.14) of the variable u . The discriminant of the trinomial $u \mapsto u^2 + zu - t + z^2$ is equal to $4t - 3z^2$. Therefore, Equation (4.16) (in variable u) has a double root either when $4t - 3z^2 = 0$ (i.e. the trinomial has a double root) or when z is also zero of the trinomial, namely $t \equiv 3z^2$.

Now, we are going to explain schematically how the holomorphic ramification of the solution (in the complex domain) can explain the appearance of a shock of the iBE $u_t - uu_x = 0$ in the real domain, i.e. $(t, x) \in \mathbb{R}^2$. Indeed, consider two small enough real numbers $0 < x_1 < x_0$. Then, the real solution $\mathbb{R}^2 \ni (t, x) \mapsto u(t, x)$ of the real iBE with CAUCHY datum $u(0, x) = x^{\frac{1}{3}}$ will have a shock at time $t_0 > 0$ if

$$x_0 - x_0^{\frac{1}{3}} t_0 = x_1 - x_1^{\frac{1}{3}} t_0 \quad (4.18)$$

since, in this case, the solution u will have two different values $x_0^{\frac{1}{3}}$ and $x_1^{\frac{1}{3}}$ at the same location $(t_0, x_0 - x_0^{\frac{1}{3}} t_0)$. From Equation (4.18) we deduce

$$t_0 = \frac{x_0 - x_1}{x_0^{\frac{1}{3}} - x_1^{\frac{1}{3}}} \equiv x_0^{\frac{2}{3}} + x_0^{\frac{1}{3}} x_1^{\frac{1}{3}} + x_1^{\frac{2}{3}}.$$

Therefore, since $0 < x_1 < x_0$, we get the following inequalities:

$$3x_1^{\frac{2}{3}} < t_0 < 3x_0^{\frac{2}{3}}.$$

Now consider the holomorphic germ $u(t, x)$ of the solution to the iBE such that $u(0, x) = x^{\frac{1}{3}}$. First, we consider the line $(t, y \equiv x_0 - x_0^{\frac{1}{3}} t)$ for $t \in [0, t_0]$ and consider (4.16) with $z = x_0^{\frac{1}{3}}$. Along this line, $z = x_0^{\frac{1}{3}}$ will never be a double root of Equation (4.16) because $t_0 < 3x_0^{\frac{2}{3}}$. Consequently, along this line, the determination of the solution will not change when we cross the vanishing locus of the discriminant (4.15) of Equation (4.16). Now, consider the other line $(t, y \equiv x_1 - x_1^{\frac{1}{3}} t)$ for $t \in [0, t_0]$ and consider (4.16) with $z = x_1^{\frac{1}{3}}$. Then, along this line at the time instance $t_1 = 3x_1^{\frac{2}{3}}$ we cross the locus (4.15) and $z = x_1^{\frac{1}{3}}$ becomes a double root of Equation (4.16) as explained above. Then, for $t > t_1 \equiv 3x_1^{\frac{2}{3}}$, the determination of the solution of the inviscid BURGERS Equation (iBE) changes along the remaining part of the line, which will explain the appearance of a shock wave taking place at time $t = t_0$. In Appendix C we explain this phenomenon on a simpler example. Overall, we may conclude that holomorphic ramified singularities do not allow us to predict what will happen **after** the shock wave appearance in the *real* solution.

Remark 4.9. If in CAUCHY problem (4.1) we take another ramified initial data

$$u(0, x) = x^{\frac{1}{5}},$$

then direct computations show that the algebraic relation between u , t and y would be

$$u^5 - ut - y = 0.$$

4.4. Generalizations

A natural multidimensional generalization of the IVP (4.2) for an unknown function

$$\begin{aligned} u : \mathbb{C} \times \mathbb{C}^n &\longrightarrow \mathbb{C}, \\ (t, x_1, x_2, \dots, x_n) &\mapsto u(t, x_1, x_2, \dots, x_n) \end{aligned}$$

is the following one:

$$u_t - u(u_{x_1} + u_{x_2} + \dots + u_{x_n}) = \underline{0}, \tag{4.19a}$$

$$u(0, x_1, x_2, \dots, x_n) = u_0(x_1, x_2, \dots, x_n) + x_1^{\frac{1}{p}}, \tag{4.19b}$$

for some holomorphic function u_0 in the vicinity of $(0, 0, \dots, 0) \in \mathbb{C}^n$. The right-hand side in the multi-dimensional PDE could be a polynomial in variables t , x and u . In the homogeneous case, it is not difficult to show the following

Theorem 4.10. *Consider the IVP (4.19). Then, there exist p germs of holomorphic functions a_0, a_1, \dots, a_{p-1} in the vicinity of $(t, x_1, x_2, \dots, x_n) \equiv (t, \mathbf{x}) := (0, 0, \dots, 0) \in \mathbb{C}^{n+1}$ along with a holomorphic function $(t, \mathbf{x}) \mapsto v(t, \mathbf{x})$ defined in the vicinity of the origin such that $u(t, \mathbf{x}) \stackrel{\text{def}}{=} v(t, \mathbf{x}) + z(t, \mathbf{x})$ is the unique solution to (4.19). The function $(t, \mathbf{x}) \mapsto z(t, \mathbf{x})$ is defined as a solution to the following algebraic equation (with t and \mathbf{x} being seen as parameters):*

$$z^p = a_{p-1}(t, \mathbf{x})z^{p-1} + \dots + a_1(t, \mathbf{x})z + a_0(t, \mathbf{x})$$

together with ‘initial’ conditions:

$$\begin{aligned} a_0(0, x_1, x_2, \dots, x_n) &= x_1, \\ a_j(0, x_1, x_2, \dots, x_n) &= \underline{0}, \quad j \in (p-1)^\square. \end{aligned}$$

Proof. This can be done by analogy with the proof of Theorem 4.1. □

5. A ring of algebraic convergent power series

In the present Section, we glimpse the very rich and complex geometry of the ring¹⁷ of formal power series. The goal of this Section is to provide a differential study

¹⁷If necessary, this ring can also be considered an algebra over the field of complex numbers \mathbb{C} .

of the algebra $\mathcal{O}[[z]]$ (with $z^3 = pz + q$). This study will exhibit some of the tools allowing us to construct the Algorithm of Section 7.1, which will provide a test of our Conjecture 7.1.

To fix the ideas, let us consider the following ring [51]:

Definition 5.1. We define

$$\mathcal{O}[[z]] \stackrel{\text{def}}{=} \left\{ \sum_{j=0}^{+\infty} a_j(p, q) z^j \right\},$$

where the coefficients $\{a_j(p, q)\}_{j \in \mathbb{N}}$ are the germs of analytic functions at the origin¹⁸ $(0, 0) \in \mathbb{C}^2$ and the complex variable z satisfies the algebraic equation (4.5). The algebra of these germs is denoted by \mathcal{O} . We require also that there exists $\exists \varepsilon > 0$ (depending on the $\{a_j(p, q)\}_{j \in \mathbb{N}}$) such that

$$\sum_{j=0}^{+\infty} \sup_{\substack{|p| < \varepsilon \\ |q| < \varepsilon}} |a_j(p, q)| \varepsilon^j < +\infty.$$

By applying the WEIERSTRASS division theorem [44], we may show that this ring is isomorphic to the following reduced form:

$$\mathcal{O}[z] \stackrel{\text{def}}{=} \left\{ a(p, q) + b(p, q)z + c(p, q)z^2 \mid z^3 = pz + q \right\},$$

for some germs of analytic functions a , b and c . Our goal consists of building some form of the differential and integral calculus on $\mathcal{O}[[z]]$ (or, in our case, equivalently on $\mathcal{O}[z]$).

Remark 5.2. Notice that we may remove the explicit dependence on variable q in coefficients a , b and c since according to Equation (4.5), $q = z^3 - pz$ and the coefficients (a, b, c) may be re-expanded in the second variable $q \mapsto \{a, b, c\}(p, q)$ after the substitution of the new expression for q . Finally, after collecting all the terms, we obtain the equivalent representation of the ring using a formal power series:

$$\mathcal{O}[z] \stackrel{\text{def}}{=} \left\{ \sum_{j=0}^{+\infty} b_j(p) z^j \mid z^3 = pz + q \right\}.$$

It is not difficult to see that the same remark applies also to the ring $\mathcal{O}[[z]]$ that we considered from the beginning:

$$\sum_{j=0}^{+\infty} a_j(p, q) z^j \equiv \sum_{j=0}^{+\infty} b_j(p) z^j.$$

Namely, we have demonstrated the following

¹⁸All germs in this study are implicitly supposed to be defined at the origin $(0, 0) \in \mathbb{C}^2$.

Proposition 5.3. *Every element $\sum_{j=0}^{+\infty} a_j(p, q) z^j \in \mathcal{O}[[z]]$ may be written under the form $\sum_{j=0}^{+\infty} b_j(p) z^j$ for some appropriately chosen holomorphic coefficients $\{b_j(p)\}_{j=0}^{+\infty}$.*

Now, we start to construct the differential calculus over the rings $\mathcal{O}[[z]]$ and $\mathcal{O}[z]$:

Lemma 5.4. *The partial derivatives of the solution z with respect to coefficients p and q are given by:*

$$\frac{\partial z}{\partial p} = \frac{z}{3z^2 - p}, \quad \frac{\partial z}{\partial q} = \frac{1}{3z^2 - p}.$$

Proof. By taking the differential of Equation (4.5), we obtain:

$$(3z^2 - p) dz = z dp + dq$$

or

$$dz = \frac{z}{3z^2 - p} dp + \frac{1}{3z^2 - p} dq.$$

After identifying the left and right-hand sides, we obtain the required result. \square

Remark 5.5. Lemma 5.4 (and Section 4.2) shows that the derivatives of a branch of $z(p, q)$ blow up when the point (p, q) tends to a point of the discriminant locus $\Delta^<(\{0\})$.

5.1. Primitivization

For the reasons which will become clearer below, the goal of this Section is to introduce and study the primitive of the ring $\mathcal{O}[z]$ with respect to the variable q :

$$\partial_q^{-1} \mathcal{O}[z] \stackrel{\text{def}}{=} \{a + bz + cz^2 \in \mathcal{O}[z] \mid \partial_q(a + bz + cz^2) \in \mathcal{O}[z]\}.$$

The following lemma shows that the derivative ∂_q of an element of $\mathcal{O}[z]$ belongs to $\mathcal{O}[z]$ only under certain conditions:

Lemma 5.6. *Let z be a solution to Equation (4.5) and $\Delta := 4p^3 - 27q^2$ be its discriminant. Take an element $a + bz + cz^2 \in \mathcal{O}[z]$. Then,*

$$\partial_q(a + bz + cz^2) \in \mathcal{O}[z]$$

if and only if the discriminant Δ divides $6cq - 2bp$ and $9bq - 4cp^2$.

Proof. Let us show the main steps of computations which yield the required result. First of all, we differentiate the element $a + bz + cz^2 \in \mathcal{O}[z]$ with respect to q and we use the result from Lemma 5.4:

$$\partial_q(a + bz + cz^2) = a_q + b_q z + c_q z^2 + \frac{b + 2cz}{3z^2 - p}.$$

It is clear that the derivative $\partial_q(a + bz + cz^2) \in \mathcal{O}[z]$ if and only if the last fraction can be presented as

$$\frac{b + 2cz}{3z^2 - p} \equiv \alpha + \beta z + \gamma z^2$$

for some holomorphic germs α , β and γ . The system of algebraic equations to determine these germs is obtained from the following obvious relation:

$$b + 2cz = (\alpha + \beta z + \gamma z^2)(3z^2 - p).$$

After expanding the powers of z higher than two using Equation (4.5) and identifying the coefficients in front of z^0 , z^1 and z^2 , we obtain the following system of equations:

$$\begin{aligned} 3\beta q - \alpha p &= b, \\ 3\gamma q + 2\beta p &= 2c, \\ 3\alpha + 2\gamma p &= \underline{0}. \end{aligned}$$

Simple algebraic manipulations yield:

$$\begin{aligned} \alpha &:= -\frac{2p(6cq - 2bp)}{\Delta}, \\ \beta &:= \frac{9bq - 4cp^2}{\Delta}, \\ \gamma &:= \frac{3(6cq - 2bp)}{\Delta}. \end{aligned}$$

This yields the conclusion of this Lemma. \square

Similar conditions can be formulated for the derivation with respect to p . However, it is out of the scope of our study.

The next question we have to discuss is the integral calculus on $\mathcal{O}[z]$ in the sense of the operation inverse to the differentiation. Let us focus on the independent variable¹⁹ q and introduce the following

Definition 5.7. An element $w(z) \in \mathcal{O}[z]$ is called a primitive of $v(z) \in \mathcal{O}[z]$ if two conditions²⁰ are satisfied:

- $\partial_q w = v$,
- $w(0) = 0$.

This fact will be denoted as $w = \partial_q^{-1} v$ and, by definition, we have $\partial_q \circ \partial_q^{-1} = \mathbb{1}_{\mathcal{O}[z]}$.

¹⁹The reason is that the integration only with respect to this variable q can be shown to be well-defined.

²⁰The second condition is imposed to remove any ambiguity with respect to constant terms.

To illustrate this definition, we provide one concrete example:

$$\partial_q^{-1} z = \frac{3qz + pz^2}{4}.$$

The last formula can be obtained by specifying Equation (5.1) to z . Otherwise, it can be checked by a direct verification.²¹

In the following two Lemmas 5.8 and 5.9, we show that this definition is meaningful and effective. However, first of all, we make use of the Remark 5.2 and switch back to the isomorphic ring object $\mathcal{O}[[z]]$ since it is directly used in our construction presented in Section 7.1:

Lemma 5.8. *The primitive $\partial_q^{-1} v, \forall v \in \mathcal{O}[[z]]$ is unique.*

Proof. In order to show this statement, it is enough to demonstrate that $\partial_q w = 0$ and $w(0) = 0$ imply that $w = \underline{0} \in \mathcal{O}[[z]]$. Indeed, from the second condition, we conclude that

$$w = \sum_{j=1}^{\infty} c_j(p) z^j$$

for some germs $\{c_j(p)\}_{j=1}^{\infty}$. Let us compute its derivative (which exists because w is a primitive):

$$\partial_q w = \sum_{j=1}^{\infty} j c_j(p) \frac{z^{j-1}}{3z^2 - p} = \underline{0}.$$

We multiply both sides of the last equality by $3z^2 - p$:

$$\sum_{j=1}^{\infty} j c_j(p) z^{j-1} = \underline{0}.$$

By using the unicity theorem for power series, we conclude that

$$c_j(p) \equiv \underline{0}, \quad \forall j \geq 1.$$

□

Lemma 5.9. *Let $\sum_{j=0}^{\infty} a_j(p) z^j \in \mathcal{O}[[z]]$. Then, the primitive ∂_q^{-1} of this element with respect to q belongs also to $\mathcal{O}[[z]]$ and can be explicitly computed (see Equation (5.1)).*

²¹Indeed, the following sequence of equalities holds:

$$\begin{aligned} \partial_q \circ \partial_q^{-1} z &= \partial_q \left(\frac{3qz + pz^2}{4} \right) = \frac{1}{4} \left(\frac{2pz}{3z^2 - p} + 3z + \frac{3q}{3z^2 - p} \right) \\ &= \frac{9z^3 - pz + 3q}{4(3z^2 - p)} = \frac{9z^3 - pz + 3(z^3 - pz)}{4(3z^2 - p)} \\ &= \frac{12z^3 - 4pz}{4(3z^2 - p)} = \frac{4z(3z^2 - p)}{4(3z^2 - p)} \equiv z. \end{aligned}$$

Proof. The proof is constructive. Let us assume that there exists another element $\sum_{j=0}^{\infty} b_j(p) z^j \in \mathcal{O}[[z]]$ such that

$$\partial_q \left(\sum_{j=0}^{\infty} b_j(p) z^j \right) = \sum_{j=0}^{\infty} a_j(p) z^j$$

and we shall construct it. By interchanging the summation and derivation operators, and using Lemma 5.4, we obtain:

$$\sum_{j=1}^{\infty} b_j(p) \frac{j z^{j-1}}{3z^2 - p} = \sum_{j=0}^{\infty} a_j(p) z^j.$$

By multiplying both sides by $3z^2 - p$ and rearranging the terms, we obtain:

$$\sum_{j=0}^{\infty} (j+1) b_{j+1}(p) z^j = \sum_{j=0}^{\infty} (3a_{j-2} - p a_j(p)) z^j,$$

where we assume the following convention $a_{-1} \equiv a_{-2} \stackrel{\text{def}}{=} \underline{0}$. It can be readily deduced that

$$b_{j+1}(p) := \frac{1}{j+1} (-p a_j(p) + 3a_{j-2}(p)), \quad \forall j \in \mathbb{N}.$$

Henceforth, we obtain the required result:

$$\partial_q^{-1} \left(\sum_{j=0}^{\infty} a_j(p) z^j \right) = \underbrace{\sum_{j=1}^{\infty} \frac{1}{j} (-p a_{j-1}(p) + 3a_{j-3}(p)) z^j}_{\stackrel{\text{def}}{=} w \in \mathcal{O}[[z]]}. \quad (5.1)$$

Once we constructed the primitive given in the right-hand side of the above equation, it is not difficult to check by the direct calculation that we have found the desired element in $\mathcal{O}[[z]]$:

$$\partial_q w := \partial_q \left(\sum_{j=1}^{\infty} \frac{1}{j} (-p a_{j-1}(p) + 3a_{j-3}(p)) z^j \right) = \sum_{j=0}^{\infty} a_j(p) z^j.$$

This is left to the reader as a simple exercise²². The found primitive w in Equa-

²²For the sake of completeness of our manuscript, we provide here this simple computation:

$$\begin{aligned} \partial_q w &= \sum_{j=1}^{\infty} \frac{1}{j} (3a_{j-3} - p a_{j-1}) \frac{j z^{j-1}}{3z^2 - p} = \sum_{j=1}^{\infty} (3a_{j-3} - p a_{j-1}) \frac{z^{j-1}}{3z^2 - p} \\ &= \frac{1}{3z^2 - p} \left(3 \sum_{j=1}^{\infty} a_{j-3} z^{j-1} - p \sum_{j=1}^{\infty} a_{j-1} z^{j-1} \right) \\ &= \frac{1}{3z^2 - p} \left(3z^2 \sum_{j=0}^{\infty} a_j z^j - p \sum_{j=0}^{\infty} a_j z^j \right) \equiv \sum_{j=0}^{\infty} a_j z^j. \end{aligned}$$

tion (5.1) satisfies also obviously the second condition $w(0) = 0$. \square

It is worthwhile to observe that one cannot define the primitive with respect to the other independent variable p . It is due, in particular, to the non-uniqueness as it follows from this simple example (employing again Lemma 5.4):

$$\partial_p q \equiv \partial_p(z^3 - pz) = 3z^2 \frac{z}{3z^2 - p} - z - p \frac{z}{3z^2 - p} \equiv \underline{0} = \partial_p \underline{0}.$$

Hence, we just showed the following

Proposition 5.10. *Let $w \in \mathcal{O}[[z]]$. Then, two conditions*

- $\partial_p w \equiv \underline{0}$,
- $w(0) = 0$

do not necessarily imply that $w \equiv \underline{0}$.

Proof. See above. \square

A conceptual explanation of the non-existence of the primitive ∂_p^{-1} will be given at the very end of Section 6.

The primitive with respect to q has another good property that deserves to be mentioned here: p -differentiable in the ring $\mathcal{O}[[z]]$. The following computation demonstrates this property and gives explicitly the formula for the p -derivative:

$$\begin{aligned} \partial_p \circ \partial_q^{-1} \left(\sum_{j=0}^{\infty} a_j z^j \right) &= \partial_p \left(\sum_{j=1}^{\infty} \frac{1}{j} (3a_{j-3} - pa_{j-1}) z^j \right) \\ &= \underbrace{\sum_{j=1}^{\infty} \frac{1}{j} (3a'_{j-3} - pa'_{j-1} - a_{j-1}) z^j}_{(*)} + \underbrace{\sum_{j=1}^{\infty} \frac{z^j}{3z^2 - p} (3a_{j-3} - pa_{j-1})}_{(**)}, \end{aligned}$$

where by prime $(\cdot)'$ we denote the derivative with respect to p . The first term $(*)$ is regular and belongs to $\mathcal{O}[[z]]$. The second term $(**)$ can be further transformed to the regular form:

$$\begin{aligned} (**): & \Rightarrow \sum_{j=1}^{\infty} \frac{z^j}{3z^2 - p} (3a_{j-3} - pa_{j-1}) \\ &= \frac{1}{3z^2 - p} \left(3 \sum_{j=0}^{\infty} a_j z^{j+3} - p \sum_{j=0}^{\infty} a_j z^{j+1} \right) \\ &= \frac{z}{3z^2 - p} \left(3z^2 \sum_{j=0}^{\infty} a_j z^j - p \sum_{j=0}^{\infty} a_j z^j \right) \equiv z \sum_{j=0}^{\infty} a_j z^j. \end{aligned}$$

Hence, to summarize, we just demonstrated the following result:

$$\partial_p \circ \partial_q^{-1} \left(\sum_{j=0}^{\infty} a_j z^j \right) = \sum_{j=1}^{\infty} \frac{1}{j} (3a'_{j-3} - pa'_{j-1} - a_{j-1}) z^j + z \sum_{j=0}^{\infty} a_j z^j.$$

More generally, one can similarly show that

Proposition 5.11. *Let $u = \partial_q^{-k} v$ for some element $v \in \mathcal{O}[[z]]$ and $k \in \mathbb{N}$. Then, $\partial_p^\alpha \circ \partial_q^\beta u \in \mathcal{O}[[z]]$ provided that $\alpha + \beta \leq k$.*

6. Holonomic coherent \mathcal{D} -modules

In this Section we perform the explicit construction of a coherent²³ holonomic²⁴ \mathcal{D} -module whose geometry is underlying our solution strategy of Conjecture 7.1. The goal is to study the singularities of the obtained solution. More precisely, we show that the vector²⁵ $(z, z^2)^\top$ is a solution of a holonomic \mathcal{D} -module whose characteristic variety \mathbf{V} is included into the union of the zero section and of the co-normal to the swallow tail. In the case of a smooth hyper-surface defined by $x = \underline{0}$, the multivalued function $x^{\frac{1}{3}}$ is a solution of the equation

$$x u_x - \frac{u}{3} = \underline{0}.$$

So, morally, the “micro-local” singularities of $x^{\frac{1}{3}}$ live in the co-normal of $x = \underline{0}$ and this property is one of the ingredients of the proof of Theorem 2.8. Therefore, in order to understand the Conjecture 7.1, it is important to work out the analogue of this property in the case of z satisfying $z^3 = pz + q$. This is the goal of this section.

²³We remind that a \mathcal{D} -module m is said to be *coherent* if for any $m \in \mathbb{N}^\times$ and any \mathcal{D} -module homomorphism

$$\mathcal{D}^m \stackrel{\text{def}}{:=} \underbrace{\mathcal{D} \times \mathcal{D} \times \dots \times \mathcal{D}}_{m \text{ times}} \longrightarrow m$$

has a kernel which is locally a \mathcal{D} -module of finite type.

²⁴As a reminder, a coherent \mathcal{D} -module is said to be *holonomic* if its characteristic variety \mathbf{V} is LAGRANGIAN. In other words, in every regular point $x \in \mathbf{V}$, we have

$$\mathbf{T}_x \mathbf{V} = (\mathbf{T}_x \mathbf{V})^\top,$$

where the orthogonal complement $(\cdot)^\top$ is taken in the sense of the natural symplectic structure on the co-tangent bundle $\mathbf{T}^* \mathbf{V}$. We remind that in general one has only the *involution* property of the characteristic variety [15]:

$$(\mathbf{T}_x \mathbf{V})^\top \subseteq \mathbf{T}_x \mathbf{V}.$$

This is a symplectic geometry view on the involutivity. More algebraic points of view are also possible, see *e.g.* [15, 50].

²⁵After some simplification, the solution ansatz we consider looks like $a + bz + cz^2$. So, one could be inclined to consider instead the vector $(\underline{1}, z, z^2)^\top$. However, the first component is completely regular, and since we are interested in singularities, we retain only the two last components.

So, we consider the algebraic Equation (4.5) whose discriminant Δ is given by

$$\Delta := 4p^3 - 27q^2.$$

The co-normal²⁶ n to $\mathfrak{S} \stackrel{\text{def}}{=} \Delta^<(\{0\})$ can be easily computed:

$$n(\mathfrak{S}) := \overline{n(\mathfrak{S}_{\text{reg}})} = \{(3z^2, -2z^3, z\lambda, \lambda) \mid (z, \lambda) \in \mathbb{C}^2\} \subseteq \mathbf{T}^*\mathbb{C}^2, \tag{6.1}$$

where we use the following coördinates to parametrize the co-tangent bundle:

$$\mathbf{T}^*\mathbb{C}^2 \equiv \{(p, q, \xi_1, \xi_2)\} \cong \mathbb{C}^4.$$

In the sequel, the co-normal $n(\mathfrak{S})$ will be always understood as the closure $\overline{n(\mathfrak{S}_{\text{reg}})}$ according to its definition.

Remark 6.1. The fact that the co-normal $n(\mathfrak{S})$ to the swallow tail contains only one complex direction²⁷ above the origin is absolutely crucial in our study.

Lemma 6.2. *Let us define the following polynomials on the co-tangent bundle $\mathbf{T}^*\mathbb{C}^2$:*

$$P_1(p, q; \xi_1, \xi_2) \stackrel{\text{def}}{=} \frac{p}{3}\xi_2^2 - \xi_1^2, \quad m_1 \stackrel{\text{def}}{=} \deg(P_1) = 2, \tag{6.2a}$$

$$P_2(p, q; \xi_1, \xi_2) \stackrel{\text{def}}{=} \frac{q}{2}\xi_2^3 + \xi_1^3, \quad m_2 \stackrel{\text{def}}{=} \deg(P_2) = 3, \tag{6.2b}$$

$$P_3(p, q; \xi_1, \xi_2) \stackrel{\text{def}}{=} (4p^3 - 27q^2)\xi_1^2, \quad m_3 \stackrel{\text{def}}{=} \deg(P_3) = 2, \tag{6.2c}$$

whose respective degrees in variables (ξ_1, ξ_2) are denoted by $m_k, j \in 3^\square$. Then,

$$\bigcap_{j=1}^3 P_j^<(\{0\}) = n(\mathfrak{S}) \cup \{(p, q, 0, 0)\} \subseteq \mathbf{T}^*\mathbb{C}^2.$$

Proof. Due to the continuity of polynomial functions, it is enough to check the statement on the regular part of the co-normal $n(\mathfrak{S})$. First, let us prove the inclusion \subseteq . So, consider a point

$$(p, q; \xi_1, \xi_2) \in \bigcap_{j=1}^3 P_j^<(\{0\}).$$

Observe that if $\xi_2 = 0$, then $P_1(p, q; \xi_1, \xi_2) = 0$ implies that $\xi_1 = 0$. Consequently, the point $(p, q; \xi_1, \xi_2)$ belongs to the zero section. If we assume now that $4p^3 - 27q^2 \neq 0$, then $P_3(p, q; \xi_1, \xi_2) = 0$ implies $\xi_1 = 0$. We draw similarly the conclusion that the point $(p, q; \xi_1, \xi_2)$ belongs to the zero section

²⁶Informally speaking, the co-normal $n(\mathfrak{S})$ plays the rôle of the characteristic variety for the quasi-linear operator (7.1a) defined below.

²⁷It would be more accurate to speak about a line, of course. In the real case the singular directions are $(0, \pm 1)$, which lie along the same line.

again. Now assume that $\xi_2 \neq 0$ and $4p^3 - 27q^2 = 0$. Then, there exists $z \in \mathbb{C}$ such that $(p, q) \equiv (3z^2, -2z^3)$. If $z = 0$, then from $P_1(p, q; \xi_1, \xi_2) = 0$ we deduce that $\xi_1 = 0$ and $(0, 0; 0, \xi_2)$ belongs to $\mathcal{N}(\mathfrak{S})$ as desired. If $z \neq 0$, then we can deduce from the equalities $P_{1,2}(p, q; \xi_1, \xi_2) = 0$:

$$\xi_1 = \frac{\xi_1^3}{\xi_1^2} = \frac{-\frac{q}{2}\xi_2^3}{\frac{p}{3}\xi_2^2} = z\xi_2.$$

This proves the inclusion. Now let us prove the reverse inclusion \supseteq . The zero section is clearly included in $\cap_{j=1}^3 P_j^<(\{0\})$.

So, let us consider instead $(3z^2, -2z^3, z\lambda, \lambda) \in \mathcal{N}(\mathfrak{S})$. Then, an easy computation shows that

$$P_j(3z^2, -2z^3, z\lambda, \lambda) \equiv \underline{0}, \quad \forall j \in \mathbb{3}^\square.$$

This proves the Lemma. □

Theorem 6.3. *There exists an open polydisc D centered around $\mathbf{0} \in \mathbb{C}^2$ such that $\forall k \in \mathbb{3}^\square$ and $\forall (i, j) \in 2^\square \times 2^\square$ we can find a differential operator R_{ij}^k with holomorphic coefficients on D such that:*

1. *The operator $R_{ij}^k, \forall (i \neq j)$ is of the order $m_k - 1$ at most.*
2. *The differential operator R_{ii}^k is of the order m_k and admits the polynomial $P_k(p, q; \xi_1, \xi_2)$ as its principal symbol.*
3. *The following identity holds:*

$$\begin{pmatrix} R_{11}^k & R_{12}^k \\ R_{21}^k & R_{22}^k \end{pmatrix} \cdot \begin{pmatrix} z \\ z^2 \end{pmatrix} \equiv \begin{pmatrix} \underline{0} \\ \underline{0} \end{pmatrix}.$$

Proof. This theorem can be proved along the same lines as [36, Theorem 4.37]. □

6.1. The holonomic \mathcal{D} -module construction

The holonomic \mathcal{D} -module \mathcal{M} will be constructed in the proof of the following

Theorem 6.4. *The characteristic variety²⁸ \mathbf{V} of \mathcal{M} is included in*

$$\bigcap_{j=1}^3 P_j^{\leq}(\{0\}) \equiv n(\mathfrak{S}) \cup \{(p, q, 0, 0)\}$$

and

1. *The \mathcal{D} -module \mathcal{M} is holonomic.*
2. *The vector (z, z^2) defines a holomorphic solution to \mathcal{M} on every simply connected open subset of $\mathbb{C}^2 \setminus \Delta^<(\{0\})$.*

Proof. We denote by \mathcal{D} (respectively by $\mathcal{D}^{(j)}$) the sheaf of holomorphic differential operators (of the order not greater than j , respectively) on the polydisc D . The grading of \mathcal{D} is defined as

$$\mathbf{gr} \mathcal{D} \stackrel{\text{def}}{=} \bigoplus_{j=-1}^{+\infty} \frac{\mathcal{D}^{(j+1)}}{\mathcal{D}^{(j)}}, \quad \mathcal{D}^{(-1)} \equiv \{0\}.$$

Algebraically speaking, this object is isomorphic to the polynomial algebra $\mathcal{O}[\xi_1, \xi_2]$.

We consider a \mathcal{D} -module morphism ϕ defined as:

$$\begin{aligned} \phi : \mathcal{D}^6 &\longrightarrow \mathcal{D}^2 \\ (Q_\ell^k)_{\substack{k \in 3 \square \\ \ell \in 2 \square}} &\mapsto \sum_{\substack{k \in 3 \square \\ \ell \in 2 \square}} (Q_\ell^k R_{\ell,1}^k, Q_\ell^k R_{\ell,2}^k). \end{aligned}$$

We are particularly interested in the co-kernel of this morphism:

$$\mathcal{M} \stackrel{\text{def}}{=} \text{coker } \phi \Rightarrow \text{codom } \phi / \phi_{>}(\{\text{dom } \phi\}) \equiv \mathcal{D}^2 / \phi_{>}(\{\mathcal{D}^6\})$$

along with the canonical projection

$$\pi : \mathcal{D}^2 \longrightarrow \mathcal{M}.$$

²⁸It might be useful to say a few words about the characteristic variety of a \mathcal{D} -module in case a reader is not familiar with this notion. The annihilator of \mathcal{M} is defined as

$$\mathbf{Ann} \mathcal{M} \stackrel{\text{def}}{=} \{P \in \mathbf{gr} \mathcal{D} \mid P \cdot m = \underline{0}, \forall m \in \mathbf{gr} \mathcal{M}\}.$$

It is an ideal in $\mathbf{gr} \mathcal{D}$. However, this ideal generally depends on the choice of a good filtration. So, one has to consider instead its *radical*:

$$\sqrt{\mathbf{Ann} \mathcal{M}} \stackrel{\text{def}}{=} \{P \in \mathbf{gr} \mathcal{D} \mid P^k \in \mathbf{Ann} \mathcal{M} \text{ for some } \exists k \in \mathbb{N}\}.$$

It can be shown that the object $\sqrt{\mathbf{Ann} \mathcal{M}}$ does not depend on the filtration. The set of common zeros of $\sqrt{\mathbf{Ann} \mathcal{M}}$ is referred to as the characteristic variety \mathbf{V} of \mathcal{M} :

$$\mathbf{V}(\sqrt{\mathbf{Ann} \mathcal{M}}) \stackrel{\text{def}}{=} \{(x_0, \bar{\xi}_0) \mid P(x_0, \bar{\xi}_0) \equiv 0, \forall P \in \sqrt{\mathbf{Ann} \mathcal{M}}\}.$$

We stress out that \mathcal{M} is a \mathcal{D} -module by construction. Theorem 6.3 allows us to define a \mathcal{D} -module sheaf homomorphism by the formula:

$$\begin{aligned} \mathcal{M} &\longrightarrow \mathcal{O}[z] \\ (\mathcal{A}, \mathcal{B}) &\mapsto \mathcal{A}z + \mathcal{B}z^2, \end{aligned}$$

where the right-hand side is a ramified solution existing on $D \setminus \Delta^<(\{0\})$ defined by the pair (z, z^2) . Then part (2) of the Theorem follows from Theorem 4.7. We introduce also a *good*²⁹ filtration $\{m_j\}_{j \in \mathbb{Z}_{\geq -1}}$ on m in the following way [42]:

$$m_{-1} \stackrel{\text{def}}{=} \{0\}, \quad m_j \stackrel{\text{def}}{=} \pi(\mathcal{D}^{(j)}, \mathcal{D}^{(j)}) \quad \forall j \geq 0, \quad m \equiv \bigcup_{j=-1}^{+\infty} m_j.$$

Consider $k \in 3\mathbb{Z}$ and $(\mathcal{A}, \mathcal{B}) \in m_{j+1}$ with $j \geq -1$. The grading of m is

$$\mathbf{gr} m \stackrel{\text{def}}{=} \bigoplus_{j=-1}^{+\infty} \frac{m^{(j+1)}}{m^{(j)}}.$$

The action of \mathcal{D} on m induces the action of $\mathbf{gr} \mathcal{D}$ on $\mathbf{gr} m$ in the obvious way. In m_{k+1+m_k} / m_{k+m_k} we have:

$$P_k \cdot (\mathcal{A}, \mathcal{B}) = \mathcal{A} \cdot (R_{1,1}^k, R_{1,2}^k) + \mathcal{B} \cdot (R_{2,1}^k, R_{2,2}^k) = \underline{0}.$$

Hence, the multiplication of m_{k+1} / m_k by P_k induces a zero mapping into m_{k+1+m_k} / m_{k+m_k} .

The characteristic variety of m minus the zero section $\{(p, q, 0, 0)\}$ is included into a LAGRANGIAN sub-variety of $\mathbf{T}^*\mathbb{C}^2 \setminus \{(p, q, 0, 0)\}$, so m is holonomic. This completes the proof of the Theorem. \square

Of course, a couple of holomorphic ramified functions (z, z^2) does not define a (uniform) distribution, but morally, Theorem 6.4 means that the “micro-local singularities” of (z, z^2) live in the co-normal to the swallow tail $\mathcal{N}(\mathfrak{S})$.

²⁹A filtration of m is an increasing sequence of sub-modules $\{m_j\}_{j \in \mathbb{Z}_{\geq -1}}$ of m verifying two properties:

- $m \equiv \bigcup_{j=-1}^{+\infty} m_j$.
- $\mathcal{D}^{(k)} \cdot m_j \subseteq m_{j+k}, \forall (k, j) \in \mathbb{N}^2$.

We say that a filtration $\{m_j\}_{j \in \mathbb{Z}_{\geq -1}}$ on m is *good* if m_j is coherent $\forall j \in \mathbb{Z}_{\geq -1}$ and there exists $\exists k_0 \in \mathbb{Z}_{\geq -1}$ such that $\forall k \geq k_0$:

$$\mathcal{D}^{(k)} \cdot m_j = m_{k+j}.$$

We remind that, in general, one only has a weaker property:

$$\mathcal{D}^{(k)} \cdot m_j \subseteq m_{k+j}.$$

We reiterate the important fact there exists only one direction³⁰ in $\mathcal{N}(\mathfrak{S})$ above $(0, 0) \in \mathbb{C}^2$, namely $(0, \lambda)$, $\lambda \in \mathbb{C}$. The principal symbol of ∂_q (respectively, ∂_p) does not vanish (respectively, vanishes) on $\{(0, 0; 0, \lambda) \mid \lambda \in \mathbb{C}^\times\}$. This explains why the primitive ∂_q^{-1} exists on $\mathcal{O}[[z]]$ whereas ∂_p^{-1} does not exist as we saw above in Section 5. Lastly, Theorem 6.4 shows morally that the singularities of the product $z \cdot z$ do not spread over the origin and remain confined in $\mathcal{N}(\mathfrak{S})$.

6.2. A computational digression

Let P_i and P_j ($i, j \in 3^\square$) be any two polynomials from Lemma 6.2. Theorem 6.4 shows that the characteristic variety \mathbf{V} is contained in

$$\mathbf{V} \subseteq P_i^<(\{0\}), \quad \mathbf{V} \subseteq P_j^<(\{0\}).$$

Hence, $\mathbf{T}_x \mathbf{V} \subseteq \ker dP_{i,j}(\mathbf{x})$, where $\mathbf{x} \in \mathbf{V}$ is a smooth point of the characteristic variety \mathbf{V} . Then, if $\omega \stackrel{\text{def}}{=} \sum_{k=1}^2 dx_k \wedge d\xi_k$ denotes the standard symplectic 2-form on $\mathbf{T}_x \mathbf{V}$, then $\forall h \in \mathbf{T}_x \mathbf{V}$:

$$dP_{i,j}(\mathbf{x}) \cdot h \equiv \omega(\mathcal{H}_{P_{i,j}}(\mathbf{x}), h) = 0, \tag{6.3}$$

where $\mathcal{H}_{P_{i,j}}$ is the following vector field:

$$\mathcal{H}_{P_{i,j}} \stackrel{\text{def}}{=} \sum_{k=1}^2 \left(\frac{\partial P_{i,j}}{\partial \xi_k} \frac{\partial}{\partial x_k} - \frac{\partial P_{i,j}}{\partial x_k} \frac{\partial}{\partial \xi_k} \right).$$

The second equality in (6.3) shows that $\mathcal{H}_{P_{i,j}}(\mathbf{x}) \in (\mathbf{T}_x \mathbf{V})^\perp$, where the orthogonal complement is taken in the sense of the natural symplectic structure on the (co-)tangent bundle $\mathbf{T}^* \mathbf{V}$. Due to the involutivity³¹ property of the characteristic variety \mathbf{V} [15], we have that

$$(\mathbf{T}_x \mathbf{V})^\perp \subseteq \mathbf{T}_x \mathbf{V}.$$

Henceforth, $\mathcal{H}_{P_{i,j}}(\mathbf{x}) \in \mathbf{T}_x \mathbf{V}$. Consequently, we obtain the following result:

$$\omega(\mathcal{H}_{P_i}(\mathbf{x}), \mathcal{H}_{P_j}(\mathbf{x})) \equiv \{P_i, P_j\}(\mathbf{x}) = 0,$$

where the POISSON bracket operator definition is implied by the symplectic structure:

$$\{P, Q\} \stackrel{\text{def}}{=} \sum_{k=1}^2 \left(\frac{\partial P}{\partial \xi_k} \frac{\partial Q}{\partial x_k} - \frac{\partial P}{\partial x_k} \frac{\partial Q}{\partial \xi_k} \right).$$

Since \mathbf{x} is a smooth point of \mathbf{V} , Theorem 6.4 allows to show that $\bigcap_{j=1}^3 P_j^<(\{0\})$ and \mathbf{V} coincide in a small neighbourhood of \mathbf{x} . Then, by HILBERT Nullstellensatz

³⁰It would be more accurate to speak about a line, of course. In the real case the singular directions are $(0, \pm 1)$, which lie along the same (complex) line.

³¹We remind that if the characteristic variety \mathbf{V} is LAGRANGIAN, then one has an even stronger property $(\mathbf{T}_x \mathbf{V})^\perp = \mathbf{T}_x \mathbf{V}$.

and Theorem 6.4, one deduces that the POISSON brackets $\{P_i, P_j\}$ ($i, j \in 3^{\square}$) must belong at least to the radical ideal $\sqrt{\mathcal{G}}$, where $\mathcal{G} \stackrel{\text{def}}{=} \langle P_1, P_2, P_3 \rangle$. We are going to check this property using modern computer algebra methods which are deeply based on the GRÖBNER bases methods [48].

First of all, let us initialize and load the library to work with polynomial ideals:

```
restart :
with(PolynomialIdeals) :
```

Then, we define the polynomials from Lemma 6.2:

```
P[1] := (p, q, r, s) -> 1/3*p*s^2 - r^2;
P[2] := (p, q, r, s) -> 1/2*q*s^3 + r^3;
P[3] := (p, q, r, s) -> (4*p^3 - 27*q^2)*r^2;
```

along with the procedure which allows to compute the POISSON bracket of two smooth functions:

```
PBracket := proc(P,Q)
  description "Returns the Poisson bracket of two
  functions P and Q";
  return simplify(diff(P(p, q, r, s), r)*diff(Q(p, q, r,
  s), p) - diff(P(p, q, r, s), p)*diff(Q(p, q, r, s),
  r) + diff(P(p, q, r, s), s)*diff(Q(p, q, r, s), q)
  - diff(P(p, q, r, s), q)*diff(Q(p, q, r, s), s));
end proc;
```

Then, we define the polynomial ideal \mathcal{G} :

```
I := PolynomialIdeal(P[1](p, q, r, s), P[2](p, q, r, s),
  P[3](p, q, r, s));
```

and check whether the ideal \mathcal{G} is radical:

```
IsRadical(I);
```

The last function returns the logical value **false**. Henceforth, we construct its radical:

```
RI := Radical(I);
```

We have to compute also the corresponding pairwise POISSON brackets of polynomials $P_{1,2,3}$:

```
P12 := (p, q, r, s) -> PBracket(P[1], P[2]);
P13 := (p, q, r, s) -> PBracket(P[1], P[3]);
P23 := (p, q, r, s) -> PBracket(P[2], P[3]);
```

Finally, we can check whether the POISSON brackets belong to the radical ideal $\sqrt{\mathcal{G}}$:

```

IdealMembership(P12(p, q, r, s), RI);
IdealMembership(P13(p, q, r, s), RI);
IdealMembership(P23(p, q, r, s), RI);

```

All these functions return the value **true**. This motivates us to establish an even stronger result comparing to Theorem 6.4.

It is also interesting to note that $\{P_1, P_2\}$ and $\{P_2, P_3\}$ belong even to the ideal \mathcal{G} . However, it is not the case for $\{P_1, P_3\}$. These facts can be simply checked as follows:

```

IdealMembership(P12(p, q, r, s), I);
IdealMembership(P23(p, q, r, s), I);
IdealMembership(P13(p, q, r, s), I);

```

The first two calls return **true** while the last one returns the value **false**. However, it is not difficult to check that $\{P_1, P_3\}^2$ already belongs to \mathcal{G} :

```

IdealMembership(P13(p, q, r, s)^2, I);

```

which returns the value **true**.

6.3. A stronger result

In fact, we were able to prove a stronger result compared to Theorem 6.4. One actually has the equality in Theorem 6.4:

Theorem 6.5. *Under conditions of Theorem 6.4, one has*

$$\mathbf{V} \setminus \{(p, q, 0, 0)\} = \mathcal{N}(\mathfrak{S}) \setminus \{(p, q, 0, 0)\},$$

where $\{(p, q, 0, 0)\}$ refers to the zero section.

Proof. In Theorem 6.4 we already established the inclusion $\mathbf{V} \setminus \{(p, q, 0, 0)\} \subseteq \mathcal{N}(\mathfrak{S}) \setminus \{(p, q, 0, 0)\}$ and from it we deduce that \mathcal{M} is holonomic. Let

$$\begin{aligned} \pi : \mathbf{T}^*\mathbb{C}^2 &\longrightarrow \mathbb{C}^2 \\ (\mathbf{x}, \bar{\xi}) &\mapsto \mathbf{x} \end{aligned}$$

denote the standard projection. We shall prove our statement by contradiction. So, let us assume that $\mathcal{N}(\mathfrak{S}) \setminus \{(p, q, 0, 0)\}$ is not included in $\mathbf{V} \setminus \{(p, q, 0, 0)\}$ and let us deduce a contradiction. Consider a point

$$(p_0, q_0; \xi_1^0, \xi_2^0) \in \mathcal{N}(\mathfrak{S}) \setminus (\mathbf{V} \cup \{(p, q, 0, 0)\}).$$

By homogeneity, for any $\mu \in \mathbb{C}^\times$, the point $(p_0, q_0; \mu \xi_1^0, \mu \xi_2^0)$ does not belong to \mathbf{V} . Since $\pi^{-1}\{(p_0, q_0)\} \cap \mathcal{N}(\mathfrak{S})$ is a complex line and since $\mathbf{V} \setminus \{(p, q, 0, 0)\} \subseteq \mathcal{N}(\mathfrak{S}) \setminus \{(p, q, 0, 0)\}$, we immediately deduce that $(p_0, q_0) \notin \pi(\mathbf{V} \setminus \{(p, q, 0, 0)\})$. Now recall that the characteristic variety minus the zero

section is (always) a homogeneous closed subset of $\mathbf{T}^*\mathbb{C}^2 \setminus \{(p, q, 0, 0)\}$. Therefore, there exists an open neighbourhood $\mathcal{U}^{(0)}$ of $(p_0, q_0) \in \mathbb{C}^2$ such that

$$\pi(\mathbf{V} \setminus \{(p, q, 0, 0)\}) \cap \mathcal{U}^{(0)} = \emptyset.$$

Consider now a WHITNEY stratification of \mathbb{C}^2 :

$$\mathbb{C}^2 = \bigcup_{\lambda \in \Lambda} X_\lambda$$

such that $\mathbf{V} \subseteq \bigcup_{\lambda \in \Lambda} \mathbf{N}^*X_\lambda$, where X_λ are smooth (open) sub-manifolds and \mathbf{N}^*X_λ denotes the co-normal bundle to X_λ . Recall also that $\{X_\lambda\}_{\lambda \in \Lambda}$ form a partition of \mathbb{C}^2 . Since $\pi(\mathbf{V} \setminus \{(p, q, 0, 0)\}) \cap \mathcal{U}^{(0)} = \emptyset$, we can assume that $(p, q, 0, 0) \in X_{\lambda_0}$, where X_{λ_0} is an open subset of \mathbb{C}^2 so that its co-normal is the zero section. Then, consider an open simply connected neighbourhood $\mathcal{U}^{(\lambda_0)}$ of (p_0, q_0) which is included in X_{λ_0} . Consider also an open simply connected subset $\mathcal{W}^{(\lambda_0)}$ included in $\mathcal{U}^{(\lambda_0)} \setminus \Delta^{<}(\{0\})$. Now we can apply to \mathcal{M} the first theorem of KASHIWARA [25, Theorem 3.1] (see also a brief reminder below). It states, in particular, that the restriction of

$$\text{Ext}_{\mathcal{D}}^0(\mathcal{M}, \mathcal{O}) \equiv \text{Hom}_{\mathcal{D}}(\mathcal{M}, \mathcal{O})$$

to X_{λ_0} is a local system. Since by Theorem 6.4, the vector (z, z^2) is a holomorphic solution of \mathcal{M} on $\mathcal{W}^{(\lambda_0)}$, KASHIWARA's theorem implies that one can analytically continue (z, z^2) to $\mathcal{W}^{(\lambda_0)}$. But this is impossible because if we make the analytic continuation of $\partial_q z = \frac{1}{3z^2 - p}$ along a path which ends up at (p_0, q_0) , it will blow up. This contradiction, therefore, shows that the reverse inclusion is satisfied, and this Theorem is proved. \square

For the sake of the exposition completeness, we formulate the following fundamental result of M. KASHIWARA on which our Theorem 6.5 is based:

Theorem 6.6 ([25]). *Let \mathfrak{M} be a maximally overdetermined system on a complex manifold X and $X = \bigcup X_\alpha$ be a stratification of X satisfying the regularity conditions of H. WHITNEY such that the singular support of \mathfrak{M} is contained in the union of co-normal projective bundles of the strata. Then, the restriction of $\text{Ext}_{\mathcal{D}_X}^i(\mathfrak{M}, \mathcal{O}_X)$ to each stratum is a locally constant sheaf of finite rank.*

In the Theorem above, the following notations were used:

\mathcal{D}_X The sheaf of differential operators of *finite order* on the complex manifold X .

\mathcal{O}_X The sheaf of holomorphic functions on X , which is a left coherent \mathcal{D}_X -module.

\mathfrak{M} The coherent \mathcal{D}_X -module.

Ext The right derived functor of the hom-functor.

7. Second order PDE

In this section, we study the next representative of the PDE family (3.1) with $m = 2$. Typically, we would like to develop the methods which would allow us to study the following ramified CP³² for a genuinely nonlinear second order PDE:

$$u_{tt} - u_x u_{xx} = \underline{0}, \tag{7.1a}$$

$$u(0, x) = \sum_{j=1}^{N_0} c_j x^{1 + \frac{j-1}{3}}, \tag{7.1b}$$

with $N_0 \in \mathbb{N}_{>2}$, $c_{1,2} \in \mathbb{C}^\times$ and $\{c_j\}_{j=3}^{N_0} \subseteq \mathbb{C}$ some constants. Namely, we would like to provide sufficient evidence that the following conjecture holds. This is the main goal of this section:

Conjecture 7.1. Consider the ramified IVP (7.1). Then, for each choice of a root to the algebraic equation³³ $\tau^2 - u_x(0, 0) \cdot 1^2 = \underline{0}$ (where we put $\xi \leftarrow 1 \neq 0$ by homogeneity of the principal symbol), there exist holomorphic functions $p(t, x)$, $q(t, x)$ and $\{a_j(t, x)\}_{j=0}^2$ defined in a neighbourhood of $(0, 0) \in \mathbb{C}^2$ with

$$p(0, x) = \underline{0}, \quad q(0, x) = x$$

such that the following assertion holds: the problem (7.1) admits a local solution of the form:

$$u(t, x) = \partial_q^{-1} \left(\sum_{j=0}^2 a_j(t, x) z^j \right), \tag{7.2}$$

where $z(t, x)$ satisfies the following algebraic equation:

$$z^3 = p(t, x)z + q(t, x).$$

Proof. Open problem. □

Remark 7.2. Please notice that the second order ramified IVP (7.1) contains only one CAUCHY data. To compensate this fact, we impose a special form (*i.e.* ansatz) for the solution that we are seeking. The condition $c_1 \neq 0$ is necessary in order to ensure that the equation $\tau^2 - u_x(0, 0) \cdot 1^2 = \underline{0}$ has two distinct roots. The condition $c_2 \neq 0$ means that the CAUCHY datum is sufficiently singular. It is crucial to ensure that the matrix \mathbf{M} defined in Equation (7.11) in the eikonal Equation (7.10) is invertible. Actually, the Theorem 2.5 from Section 2.1 shows that if the constant c_2 is equal to zero, then the solution will be ramified around a smooth hyper-surface.

³²We underline the fact that the term CAUCHY problem (CP) is understood here in the *incomplete* (or loose) sense specified in the Introduction Section 1 since our CAUCHY datum contains only one initial condition for a second order problem in time (7.1a).

³³We have to assume that $u_x(0, 0) \neq 0$ in order to have two distinct roots.

The choice of a root in the algebraic equation will allow fixing the value of $q_t(0, 0)$. Then, it is very likely that the solution under the form (7.2) will be unique. Notice that the functions $p(t, x)$ and $q(t, x)$ will depend on the CAUCHY datum.

Despite the fact that we do not have proof for this conjecture, we shall present in this Section strong evidence that it is true. Our symbolic–numeric computations will be performed in the case of $c_1 = 1$ and $c_2 = \frac{3}{4}$ for the sake of convenience.

In the sequel, we shall rather write the solution ansatz without applying the WEIERSTRASS division theorem [44], *i.e.*

$$\sum_{j=0}^2 a_j(t, x) z^j \equiv \sum_{k=0}^{\infty} b_k(t, x) z^k.$$

Despite the apparent complication, expressing the iterative process in this way will be easier.

7.1. Construction of iterations

Motivated by Lemma 5.8 and Theorem 6.5, we extract from the PDE (7.1a) the following characteristic equation:

$$q_t(0, 0)^2 - u_x(0, 0)q_x(0, 0) = 0.$$

Observe that $q_x(0, 0) = 1$ and that the choice $c_1 = 1$ necessarily implies $u_x(0, 0) = 1$. Then, in the previous equation, we shall choose the root $q_t(0, 0) = 1$ for the rest of this section.

The approach we adopt in this section is greatly inspired by Sections 5 and 6. We would like to construct a solution to Equation (7.1a) in the following form, motivated by Lemma 5.9:

$$u(t, x) = \sum_{k=1}^{\infty} \frac{1}{k} \left(-p(t, x)b_{k-1}(t, x) + 3b_{k-3}(t, x) \right) z^k \in \mathcal{O}[[z]], \quad (7.3)$$

where $p(t, x)$, $q(t, x)$ and $\{b_k(t, x)\}_{k=0}^{\infty}$ are holomorphic functions of its two variables $(t, x) \in \mathbb{C}^2$ defined in the vicinity $\mathcal{U}_{(0,0)}$ of $(0, 0) \in \mathbb{C}^2$ that verify the following ‘initial’ conditions:

$$p(0, x) \equiv \underline{0}, \quad q(0, x) \equiv x, \quad q_t(0, 0) = 1, \quad (7.4a)$$

$$b_0(0, x) \equiv \underline{1}, \quad b_1(0, x) \equiv \underline{1}, \quad b_k(0, x) \equiv \underline{0}, \quad \forall k \geq 2, \quad (7.4b)$$

and

$$z^3 = pz + q, \quad \forall (t, x) \in \mathcal{U}_{(0,0)} \subseteq \mathbb{C}^2.$$

The ‘initial’ conditions (7.4) correspond to the CAUCHY datum (7.1b) with $c_1 = 1$ and $c_2 = \frac{3}{4}$:

$$u(0, x) = x + \frac{3}{4} x^{\frac{4}{3}}. \quad (7.5)$$

Several other initial conditions will be considered below in Section 8.

It seems impossible to construct directly such a solution $u(t, x)$ having a property that the series (7.3) is convergent $\forall (t, x) \in \mathcal{U}_{(0,0)}$. Instead, we propose to construct this solution $u(t, x)$ using the fixed point iterative scheme.

Using the ansatz (7.3) and this scheme, we shall see that the solution is completely determined if we have the following collection of holomorphic functions:

$$\mathfrak{C} \stackrel{\text{def}}{=} \left(p(t, x), q(t, x), \{b_k(t, x)\}_{k=0}^\infty \right) \cap (7.4). \tag{7.6}$$

By construction, the functions from this collection \mathfrak{C} verify the ‘initial conditions’ (7.4). Thus, if we know these functions, we can easily recover the solution $u(t, x)$ thanks to (7.3). Hence, the collection (7.6) will be called the solution data.

Now, we will give several computations in order to understand this situation better. First, one has:

$$z_t(p(t, x), q(t, x)) = \frac{z p_t + q_t}{3z^2 - p}.$$

Next, for $u(t, x)$ as defined in Equation (7.3), one checks that

$$u_t = \sum_{k=0}^\infty \left(\left(1 - \frac{1}{k}\right) b_{k-1} p_t + b_k q_t + \frac{1}{k} (3b_{k-3,t} - p b_{k-1,t}) \right) z^k,$$

where by convention $b_j \equiv 0$ for $j < 0$, so the terms proportional to $\frac{1}{k}$ with $k = 0$ are absent from the sum above. One can easily obtain a similar formula for u_x . Next, after a lengthy computation, one gets:

$$\begin{aligned} u_{tt} &= \sum_{k=0}^\infty \left(2 \left(1 - \frac{1}{k}\right) p_t b_{k-1,t} + \left(1 - \frac{1}{k}\right) b_{k-1} p_t^2 + b_k q_{tt} + \right. \\ &\quad \left. 2q_t b_{k,t} + \frac{1}{k} (3b_{k-3,tt} - p b_{k-1,tt}) \right) z^k + \\ &\quad \frac{1}{3z^2 - p} \sum_{k=0}^\infty \left((k-1) p_t^2 b_{k-1} + 2k b_k p_t q_t + (k+1) q_t^2 b_{k+1} \right) z^k. \end{aligned}$$

Naturally, a similar formula holds for u_{xx} as well. Then, one checks that the coefficient of $\frac{1}{3z^2 - p}$ in $u_{tt} - u_x u_{xx}$ is equal to

$$\begin{aligned} &\sum_{k=0}^\infty \left((k-1) p_t^2 b_{k-1} + 2k b_k p_t q_t + (k+1) q_t^2 b_{k+1} \right. \\ &\quad \left. - \sum_{j=0}^k \left[\left(1 - \frac{1}{j}\right) b_{j-1} p_x + b_j q_x + \frac{1}{j} (3b_{j-3,x} - p b_{j-1,x}) \right] \right) \\ &\quad \left[(k-j-1) p_x^2 b_{k-j-1} + 2(k-j) b_{k-j} p_x q_x + (k-j+1) q_x^2 b_{k-j+1} \right] z^k. \end{aligned} \tag{7.7}$$

The first naive tentative to solve Conjecture 7.1 consists in replacing $u(t, x)$ by (7.3) in the nonlinear differential operator $u \mapsto u_{tt} - u_x u_{xx}$. One obtains a certain expression ($\hat{\mathfrak{X}}$). Unfortunately, it seems extremely difficult to construct directly $u(t, x)$ as in Equation (7.3) such that ($\hat{\mathfrak{X}}$) $\equiv \underline{0}$ and (7.7) be divisible by $3z^2 - p$ in $\mathcal{O}[[z]]$. Henceforth, we introduce an iterative procedure where we replace in one specific place of the expression ($\hat{\mathfrak{X}}$) the function $p(t, x)$ by $\hat{p}(t, x)$, respectively, $q(t, x)$ by $\hat{q}(t, x)$ and $\{b_k(t, x)\}_{k=0}^\infty$ by $\{\hat{b}_k(t, x)\}_{k=0}^\infty$. This procedure is better explained in Appendix B. As a result, we obtain the new expression:

$$(\hat{\mathfrak{X}}) = \sum_{k=0}^{\infty} \alpha_k z^k + \frac{\sum_{k=0}^{\infty} \beta_k z^k}{3z^2 - p}.$$

Then, by requiring that $3z^2 - p$ divides $\sum_{k=0}^{\infty} \beta_k z^k$ we obtain the eikonal Equation (7.10) which gives us the new components (\hat{p}, \hat{q}) . Roughly speaking, (7.10) means that the cusp is characteristic for the PDE. By requiring that ($\hat{\mathfrak{X}}$) vanishes, we obtain the following discrete ‘time’ (but infinite dimensional) dynamical system:

$$\left(\hat{p}(t, x), \hat{q}(t, x), \{\hat{b}_k(t, x)\}_{k=0}^\infty \right) := \mathbf{F} \left(p(t, x), q(t, x), \{b_k(t, x)\}_{k=0}^\infty \right).$$

Henceforth, we may say that the ideas of geometric optics inspire the construction of iterations. A similar procedure was employed by WAGSCHAL in [53].

Then, by the construction of the mapping \mathbf{F} , its fixed point gives us the desired solution to Conjecture 7.1. In this Section we shall construct the mapping \mathbf{F} and implement it in a computer algebra system `Maple`TM:

$$\mathbf{F} : \mathfrak{C} \longrightarrow \mathfrak{C}, \tag{7.8}$$

$$\left(p(t, x), q(t, x), \{b_k(t, x)\}_{k=0}^\infty \right) \mapsto \left(\hat{p}(t, x), \hat{q}(t, x), \{\hat{b}_k(t, x)\}_{k=0}^\infty \right).$$

As we already mentioned, the mapping \mathbf{F} is constructed such that if we have a fixed point

$$\mathbf{F} \left(p(t, x), q(t, x), \{b_k(t, x)\}_{k=0}^\infty \right) \equiv \left(p(t, x), q(t, x), \{b_k(t, x)\}_{k=0}^\infty \right),$$

then the solution (7.3) reconstructed from this data verifies (7.1a) (cf. Lemma 7.3). Additionally, since we impose Conditions (7.4b), this mapping will yield that $(q_t(0, 0))^2 = 1$. The fact that we have chosen $q_t(0, 0) = 1$ corresponds to the choice of a root to equation $\tau^2 - \partial_x u(0, 0) \cdot 1^2 = \underline{0}$ in Conjecture 7.1 (where we replaced $\xi \leftarrow 1 \neq 0$ by homogeneity of the principal symbol).

We define the following family of functions:

$$\begin{aligned} \mathcal{B}_k \stackrel{\text{def}}{=} & \sum_{j=0}^k \left[\left(1 - \frac{1}{j} \right) p_x b_{j-1} + q_x b_j + \frac{1}{j} \left(-p b_{j-1, x} + 3 b_{j-3, x} \right) \right] \times \\ & \left[(k - j - 1) p_x^2 b_{k-j-1} + 2(k - j) p_x q_x b_{k-j} + (k - j + 1) q_x^2 b_{k-j+1} \right] \\ & - (k - 1) \cdot p_t^2 b_{k-1}, \quad \forall k \in \mathbb{N}, \end{aligned}$$

where we additionally use the convention:

$$b_{-j} \equiv 0, \forall j \geq 1. \tag{7.9}$$

The *eikonal equation*³⁴ is defined as

$$\begin{pmatrix} \hat{p}_t \\ \hat{q}_t \end{pmatrix} = \mathbb{M}^{-1} \cdot \begin{pmatrix} \sum_{k=0}^{+\infty} \left(\frac{p}{3}\right)^k \mathcal{B}_{2k} \\ \sum_{k=0}^{+\infty} \left(\frac{p}{3}\right)^k \mathcal{B}_{2k+1} \end{pmatrix}. \tag{7.10}$$

Since functions \hat{p} and \hat{q} belong to the class \mathfrak{E} , they verify the following initial conditions:

$$\hat{p}(0, x) = 0, \quad \hat{q}(0, x) = x.$$

The functional 2×2 matrix \mathbb{M} is defined as

$$\mathbb{M} \stackrel{\text{def}}{=} \begin{pmatrix} 4 \sum_{k=0}^{+\infty} \left(\frac{p}{3}\right)^k k q_t b_{2k} & \sum_{k=0}^{+\infty} \left(\frac{p}{3}\right)^k (2k+1) q_t b_{2k+1} \\ 2 \sum_{k=0}^{+\infty} \left(\frac{p}{3}\right)^k (2k+1) q_t b_{2k+1} & 2 \sum_{k=0}^{+\infty} \left(\frac{p}{3}\right)^k (k+1) q_t b_{2k+2} \end{pmatrix}. \tag{7.11}$$

We underline that the inversion of the last matrix is necessary to obtain an explicit form of the eikonal System (7.10) solved with respect to the time derivatives. It is not difficult to see (by substituting the necessary initial data and conventions) that the first Equation in (7.10) yields $\hat{p}_t(0, 0) = \frac{1}{2}$.

Let us define also the following family of functions:

$$\begin{aligned} \mathcal{A}_k &\stackrel{\text{def}}{=} (k-1)b_{k-1}p_t^2 + 2kb_k\hat{p}_tq_t + (k+1)\hat{q}_tq_tb_{k+1} - \\ &\sum_{j=0}^k \left[\left(1 - \frac{1}{j}\right)b_{j-1}p_x + b_jq_x + \frac{1}{j}(-pb_{j-1,x} + 3b_{j-3,x}) \right] \times \\ &\left[(k-j-1)b_{k-j-1}p_x^2 + 2(k-j)b_{k-j}p_xq_x + (k-j+1)b_{k-j+1}q_x^2 \right], \\ &\forall k \in \mathbb{N}. \end{aligned}$$

Also, for any $k \in \mathbb{N}$ we define:

$$\mathcal{C}_k \stackrel{\text{def}}{=} \sum_{j=0}^{\infty} \mathcal{A}_{k+2+2j} \left(\frac{p}{3}\right)^j.$$

³⁴Strictly speaking, the term ‘*eikonal equation*’ is not defined for genuinely nonlinear PDEs and singular hyper-surfaces. However, we have good reasons to think that what we are doing is the right generalization of this notion to our case.

Now, we are ready to write down equations which define the family of functions $\{\hat{b}_k(t, x)\}_{k=0}^{\infty}$:

$$\begin{aligned} \frac{\partial \hat{b}_k}{\partial t} = & \frac{1}{2q_t} \left\{ - \left(2 - \frac{2}{k}\right) b_{k-1,t} p_t - \left(1 - \frac{1}{k}\right) b_{k-1} p_{tt} - b_k q_{tt} \right. \\ & - \frac{1}{k} (-p b_{k-1,tt} + 3b_{k-3,tt}) - \mathcal{C}_k \\ & + \sum_{j=0}^k \left[\left(1 - \frac{1}{j}\right) b_{j-1} p_x + b_j q_x + \frac{1}{j} (-p b_{j-1,x} + 3b_{j-3,x}) \right] \\ & \times \left[\left(2 - \frac{2}{k-j}\right) b_{k-j-1,x} p_x + \left(1 - \frac{1}{k-j}\right) b_{k-j-1} p_{tt} + b_{k-j} q_{xx} \right. \\ & \left. \left. + \text{frm-e} b_{k-j,x} q_x + \frac{1}{k-j} (-p b_{k-j-1,xx} + 3b_{k-j-3,xx}) \right] \right\}, \quad \forall k \in \mathbb{N}. \end{aligned} \tag{7.12}$$

We underline that functions $\{\hat{b}_k(t, x)\}_{k=0}^{\infty}$ satisfy the initial conditions (7.4b). The convention (7.9) must be employed to interpret correctly the last equation.

The solution of differential Equations (7.10) and (7.12) completes the construction of the fixed point mapping \mathbf{F} (7.8). Then, the connection between the mapping \mathbf{F} and the ramified CAUCHY problem (7.1) is elucidated in the following

Lemma 7.3. *Let us assume that we have fixed point data:*

$$\mathbf{F} \left(p(t, x), q(t, x), \{b_k(t, x)\}_{k=0}^{\infty} \right) = \left(p(t, x), q(t, x), \{b_k(t, x)\}_{k=0}^{\infty} \right)$$

satisfying the ‘initial’ conditions (7.4). Then, the function

$$u(t, x) = \partial_q^{-1} \left(\sum_{k=0}^{+\infty} b_k(t, x) z^k \right), \tag{7.13}$$

with $z^3 = pz + q$ satisfies PDE (7.1a) together with the initial condition (7.1b).

Proof. Recall that by direct computations, we have the following expressions for ansatz (7.13) derivatives:

$$u_{\alpha} = \sum_{k=0}^{+\infty} \left[\left(1 - \frac{1}{k}\right) b_{k-1} p_{\alpha} + b_k q_{\alpha} + \frac{1}{k} (-p b_{k-1,\alpha} + 3b_{k-3,\alpha}) \right] z^k,$$

$$\begin{aligned}
 u_{\alpha\alpha} &= \sum_{k=0}^{+\infty} \left[\left(2 - \frac{2}{k} \right) b_{k-1,\alpha} p_{\alpha} + \left(1 - \frac{1}{k} \right) b_{k-1} p_{\alpha\alpha} + b_k q_{\alpha\alpha} + 2b_{k,\alpha} q_{\alpha} \right. \\
 &\quad \left. + \frac{1}{k} (pb_{k-1,\alpha\alpha} + 3b_{k-3,\alpha\alpha}) \right] z^k \\
 &+ \frac{1}{3z^2 - p} \sum_{k=0}^{+\infty} \left[(k-1)b_{k-1}p_{\alpha}^2 + 2kb_k p_{\alpha} q_{\alpha} + (k+1)b_{k+1}q_{\alpha}^2 \right] z^k,
 \end{aligned}$$

where the symbol $\alpha \in \{t, x\}$.

Now, let us assume that

$$\left(\hat{p}(t, x), \hat{q}(t, x), \{\hat{b}_k(t, x)\}_{k=0}^{\infty} \right) = \left(p(t, x), q(t, x), \{b_k(t, x)\}_{k=0}^{\infty} \right).$$

First, the fact that $(\hat{p}(t, x), \hat{q}(t, x)) = (p(t, x), q(t, x))$ and the eikonal Equation (7.10) imply that the coefficient (7.7) of $\frac{1}{3z^2 - p}$ in $u_{tt} - u_x u_{xx}$ belongs indeed to $\mathcal{O}[z]$. For more details, refer to Appendix B. Then, the fact that $\hat{b}_k \equiv b_k$ for $\forall k \in \mathbb{N}$ and Equation (7.12) imply, by construction, that $u_{tt} - u_x u_{xx} \equiv \underline{0}$. See Appendix B again for more details. \square

To the best of our knowledge, the presented algorithm is completely new.

7.2. Generalizations

We may attempt to generalize the result stated in Theorem 4.1 to the case of the PDEs family (3.1) with $m \geq 2$. However, we stress that at the current stage, it remains at the level of a conjecture:

Conjecture 7.4. Consider the following ramified IVP for Equation (3.1):

$$\mathcal{L}_m(u) := \partial_t^m u - \partial_x^{m-1} u \partial_x^m u = \underline{0}, \tag{7.14a}$$

$$\begin{aligned}
 u(0, x) &= c_1 x^{m-1} + c_2 x^{m-1+\frac{1}{3}} \\
 &+ c_3 x^{m-1+\frac{2}{3}} + x^m \cdot u_0(x), \tag{7.14b}
 \end{aligned}$$

where $c_{1,2} \in \mathbb{C}^\times$, $c_3 \in \mathbb{C}$ and u_0 is a holomorphic function in some neighbourhood of $0 \in \mathbb{C}$. Then, for each choice of a root to the algebraic equation³⁵ $\tau^m - \partial_x^{m-1} u(0, 0) \cdot 1^m = \underline{0}$ (where we replaced $\xi \leftarrow 1 \neq 0$ by homogeneity of the principal symbol), there exist holomorphic functions $p(t, x)$, $q(t, x)$ and $\{a_j(t, x)\}_{j=0}^2$ defined in the vicinity of $(0, 0) \in \mathbb{C}^2$ such that there exists a

³⁵In order to have m distinct roots, we have to choose the initial data so that $\partial_x^{m-1} u(0, 0) \neq 0$. As a result, we may say that the solution procedure is parametrized by the elements of the multiplicative group of the roots of unity $U_m \stackrel{\text{def}}{=} \{\xi^m = 1 \mid \xi \in \mathbb{C}\}$.

local solution u to (7.14) which can be written in the following form:

$$u(t, x) := \partial_q^{-(m-1)} \circ \left(\sum_{j=0}^2 a_j(t, x) z^j \right),$$

where the function z satisfies the following algebraic relation (4.5):

$$z^3 = p(t, x)z + q(t, x).$$

The coefficients p, q verify the ‘initial’ conditions:

$$p(0, x) = \underline{0}, \quad q(0, x) = x.$$

Proof. Open problem. □

Remark 7.5. The holomorphic functions $p(t, x), q(t, x)$ and $\{a_j(t, x)\}_{j=0}^2$ whose existence is conjectured above depend also on $c_{1,2,3}$ and on the function $u_0(x)$ appearing in the formulation of the CAUCHY problem (7.14b).

8. Numerical illustrations

In the absence of rigorous theoretical proof, in this section, we would like to illustrate the practical convergence of the fixed point algorithm described in Section 7.1. Thus, it will provide rational computational support towards various conjectures formulated in this study (in particular, the Conjecture 7.1 and indirectly Conjecture 7.4).

8.1. Test 1

In order to come up with a practical algorithm, we have to truncate the expansion (7.3):

$$u(t, x) \approx \sum_{k=1}^N \frac{1}{k} \left(-p(t, x)b_{k-1}(t, x) + 3b_{k-3}(t, x) \right) z^k,$$

Since we work with holomorphic functions, we shall expand all the functions in double truncated series in the vicinity of the origin $(0, 0) \in \mathbb{C}^2$ as follows:

$$\psi(t, x) = \sum_{\substack{l, m=1 \\ l+m \leq M-1}}^{M-1} \psi_{lm} t^l x^m + \mathcal{O}(t^M + t^{M-1}x + \dots tx^{M-1} + x^M),$$

with $\psi \in \{p, q, b_0, \dots, b_{N-1}\}$. So, in practice, we perform all our computations with such double truncated power series in independent variables $(t, x) \in \mathbb{C}^2$. We have to describe another technical issue towards the practical implementation

Table 1: *Various numerical parameters used in our computations.*

<i>Parameter</i>	<i>Value</i>
Floating point arithmetics, significant digits	30
Truncation degree in t , M	25
Truncation degree in x , M	25
Total number of iterations, I	25
Initial length of data vector, N	$80 = 3 \times I + 5$
Observation point in \mathbb{C} , x_0	$0.1i$
Temporal segment for the error control, $[a, b]$	$[0, 0.1]$

of our algorithm here. Since we work with truncated data, our finitary mapping $\tilde{\mathbf{F}}$ by construction returns a slightly reduced set of data:

$$\left(p(t, x), q(t, x), \{b_k(t, x)\}_{k=0}^N \right) \xrightarrow{\tilde{\mathbf{F}}} \left(\hat{p}(t, x), \hat{q}(t, x), \{\hat{b}_k(t, x)\}_{k=0}^{N-3} \right).$$

Thus, at every iteration, we are practically losing three components of the data. Consequently, if we are willing to obtain N components of data vector $\{b_k(t, x)\}_{k=0}^N$ after I iterations, we have to initiate the iterative process with $\{b_k(t, x)\}_{k=0}^{N+3I}$ initial components. We took into account this observation in our computations. We are going to illustrate the work of the algorithm starting from the following initial guess:

$$p(t, x) := \frac{t}{2}, \quad q(t, x) := t + x, \quad (8.1a)$$

$$b_0 := \underline{1}, \quad b_1 := \underline{1}, \quad b_j := \underline{0}, \quad \forall j \geq 2. \quad (8.1b)$$

It corresponds to the CAUCHY datum (7.5) as it is explained at the beginning of Section 7.1. Throughout all this Section 8 we shall assume $c_1 = 1$ and $c_2 = \frac{3}{4}$. However, three different possibilities are taken for the rest of coefficients³⁶ $\{c_j\}_{j \geq 3} \subseteq \mathbb{C}$. For example, in this Section 8.1 we take all $\{c_j\}_{j \geq 3}$ to be zero.

Remark 8.1. The choice of the initial data (8.1a) implies that $q_t(0, 0) = 1$. This specifies one of two roots in $q_t(0, 0)$ in the eikonal equation. Consequently, in all iterations we shall also have that $\hat{q}_t(0, 0) = 1$. Similarly, we shall also have $\hat{p}_t(0, 0) = \frac{1}{2}$.

In order to speed up the computations, we also turn to the floating-point arithmetics with, possibly, extended precision. The values of all other parameters are reported in Table 1. The proposed algorithm was implemented in the computer algebra system `Maple`TM. The essential part of the employed code is reported in Appendix D. The complete program can be shared upon a simple request by email.

³⁶For the precise meaning of coefficients $\{c_j\}_{j \geq 0}$ see Equation (7.1b).

To summarize, we can say that we perform numeric-symbolic computations in order to observe the convergence towards a fixed point in practice. If we denote by $\psi^{(j)}$ the value of the variable $\psi \in \{p, q, b_0, \dots, b_{N-3j}\}$ after j iterations, the closeness to the fixed point may be appreciated by looking at the norm of the difference between two successive iterations:

$$\|\psi^{(j)} - \psi^{(j-1)}\|.$$

We illustrate the behaviour of the proposed algorithm on the initial data (8.1). We perform I iterations of our mapping. It turns out that even very moderate values of the parameter I (cf. Table 1) are enough to appreciate the convergence of the iterative fixed point process. The whole symbolic/numeric computation lasted about five minutes of the CPU time (on our computers). The precise information is not very important because it may vary from one system to another. In four panels of Figure 3 we show the differences between two last iterations of functions $\{p, q, b_0, b_1\}$, constituting the problem data. It can be clearly seen that the absolute value of the difference is quite small already after $I = 10$ iterations. We checked that subsequent iterations reduced this difference further. The reduction of the L_∞ norm of the difference between two successive iterations is shown in Figure 4. This Figure presents strong empirical evidence for the convergence towards a fixed point, which solves the underlying IVP according to Lemma 7.3, even if the convergence seems to be far from being linear.

In order to present even more convincing numerical evidence, we include in our study another test case with the following initial guess:

$$p(t, x) := \frac{t}{2}, \quad q(t, x) := t + x, \quad (8.2a)$$

$$b_0 := \underline{1} + t, \quad b_1 := \underline{1} - t, \quad b_j := \underline{0}, \quad \forall j \geq 2. \quad (8.2b)$$

It is not difficult to see that these initial data (8.2) (which correspond to a different initialization with respect to (8.1)) verify the same initial conditions (7.4), *i.e.* the CAUCHY datum (7.5). Thus, if the solution we seek is unique³⁷, we may expect the iterations to converge to the same fixed point. This explains the motivation behind this second numerical study. We are using precisely the same numerical parameters as reported above in Table 1 (and, thus, the same parameters as in the previous computations). The convergence of the fixed point iterations is reported in Figure 5. We can observe the same (cf. Figure 4) roughly linear but non-monotonic convergence to a fixed point. Now one may ask a legitimate question whether iterations starting at (8.1) and (8.2) converge to the same point in $\mathcal{O}[[z]]$? We may answer this question by looking at the difference between obtained solutions data at the final iteration I . We can have a look at these differences $t \mapsto |\psi_1^{(I)}(t, x_0) - \psi_2^{(I)}(t, x_0)|$ in Figure 6 where they are represented as

³⁷We may reasonably assume that at least within the solution ansatz we consider. Perhaps the best reason to believe in the uniqueness would be the LAX–GLIMM theory [20], but “we sadly lack a local uniqueness theorem” as the authors of [20] put it themselves (in the real case).




Figure 3: The absolute value of the difference between two successive iterations (I)th and ($I - 1$)th of solution data for $I = 10$. The values of several numerical parameters are reported in Table 1.




Figure 4: The maximum norm $L_{\infty}([a, b])$ of the difference between two successive iterations $(I)^{\text{th}}$ and $(I-1)^{\text{th}}$ as a function of the iteration number in semi-logarithmic coordinates. The initial data is given in Equation (8.1). The values of several numerical parameters are reported in Table 1.

functions of time³⁸ for $t \in [a, b] \equiv \left[0, \frac{1}{10}\right]$. In particular, we would like to underline the fact that the magnitude of these differences is consistent with the results presented in Figures 4 and 5. All these observations provide strong empirical evidence for the convergence of the iterative process described above. The experimental findings of this Section are schematically depicted and summarized in Figure 7.

8.2. Test 2

We shall consider a completely different initial condition from (7.1b):

$$u(0, x) = x + \frac{3}{4}x^{\frac{4}{3}} + \frac{3}{50}x^{\frac{8}{3}} + \frac{1}{20}x^3. \quad (8.3)$$

In terms of the solution data, the last initial datum translates in the following coefficients:

$$p(t, x) := \frac{t}{2}, \quad q(t, x) := t + x, \quad (8.4a)$$

$$b_0 := \underline{1}, \quad b_1 := \underline{1}, \quad (8.4b)$$

$$b_2(t, x) := \frac{x}{10}, \quad b_3(t, x) := \frac{x}{10}, \quad (8.4c)$$

$$b_j := \underline{0}, \quad \forall j \geq 4. \quad (8.4d)$$

The complexity of this test case is much higher than what we did before. In order to make the computational times reasonable, we had to reduce the parameter $M := 10$. All other numerical parameters were kept as in Table 1. We studied the convergence of fixed point iterations under the map \tilde{F} for this IVP. Namely, we monitored the differences between successive iterations for variables p, q, b_0, \dots, b_3 . The results are reported in Figure 8. As we can see, the *practical* linear non-monotonic convergence can be clearly witnessed even in this test case.

As we do it systematically, we also consider another initialization of the fixed point algorithm, which corresponds to the same CAUCHY datum (8.3) as before:

$$p(t, x) := \frac{t}{2}, \quad q(t, x) := t + x, \quad (8.5a)$$

$$b_0 := \underline{1}, \quad b_1 := \underline{1} - \frac{t^2}{10}, \quad (8.5b)$$

$$b_2(t, x) := \frac{x}{10} + \frac{t}{100}, \quad b_3(t, x) := \frac{x}{10} - \frac{t}{100}, \quad (8.5c)$$

$$b_j := \underline{0}, \quad \forall j \geq 4. \quad (8.5d)$$

The algorithm was run with this initialization and all other parameters as above. The convergence of the iterations initialized with the initial date (8.5) is reported in

³⁸It turns out that in this particular case, these differences are only functions of time, *i.e.* they do not depend on x .




Figure 5: The maximum norm $L_\infty([a, b])$ of the difference between two successive iterations $(I)^{\text{th}}$ and $(I - 1)^{\text{th}}$ as a function of the iteration number in semi-logarithmic coordinates. The initial data is given in Equation (8.2). The values of several numerical parameters are reported in Table 1.




Figure 6: Differences between corresponding components of the solution data at $I = 25^{th}$ iteration. Both iterative processes were initialized with data (8.1) and (8.2) respectively. The values of all numerical parameters are given in Table 1.




Figure 7: A schematic representation of the experimental findings of Section 8: $P_1^{(0)}$ stands for the initial guess (8.1) while $P_2^{(0)}$ stands for (8.2). The convergence should be understood in the sense of a numerically-supported conjecture.

Figure 9. We can observe a very similar behaviour to Figure 8 attesting one more time to the convergence of the proposed algorithm. One may ask the legitimate question of whether two different initializations (8.4) and (8.5) converge to the same point. This theoretical question is very complicated. However, what we can do in practice is to measure how close are the corresponding iterates from each other in the spirit of mental representation from Figure 7. This is done in Figure 10 where the difference $(t, x) \mapsto |\psi_1^{(I)}(t, x) - \psi_2^{(I)}(t, x)|$ with $\psi \in \{p, q, b_0, \dots, b_3\}$. In order to make the error plot possible, we considered the real values of independent variables varying in some finite neighbourhood of the origin $\mathbf{0} \in \mathbb{R}^2$:

$$\left\{ (t, x) \in \mathbb{R}^2 \mid t^2 + x^2 \leq \frac{1}{100} \right\}.$$

In particular, Figure 10 shows that the convergence is the slowest for the component b_3 of the solution data but even for b_3 the results are perfectly acceptable.

The Section below contains another couple of computational investigations of the proposed fixed point algorithm.

8.3. Test 3

As a final numerical test case, we consider another IVP (7.1), represented by two different initial guesses, which contains explicitly the fractional power $x^{\frac{4}{3}}$ along with $x^{\frac{5}{3}}$ in perfect agreement with the generalized Conjecture 7.4. Namely, in the CAUCHY datum (7.14b) there is a term $c_3 x^{m-1+\frac{2}{3}}$ (with $c_3 \in \mathbb{C}$), which for $m = 2$ gives $c_3 x^{\frac{5}{3}}$ and it was absent in previous numerical Tests 1 (Section 8.1) and 2 (Section 8.2). As a result, we decided to include an additional numerical test which contains explicitly this term. Namely, we consider the following CAUCHY




Figure 8: The maximum norm $L_\infty([a, b])$ of the difference between two successive iterations (I)th and ($I - 1$)th. The initial data is given in Equation (8.4).




Figure 9: The maximum norm $L_\infty([a, b])$ of the difference between two successive iterations $(I)^{\text{th}}$ and $(I-1)^{\text{th}}$. The initial data is given in Equation (8.5).




Figure 10: The absolute value of the differences between corresponding components of the solution data at $I = 25^{\text{th}}$ iteration. Both iterative processes were initialized with data (8.4) and (8.5) respectively. The values of all numerical parameters are given in Table 1.

datum:

$$u(0, x) = x + \frac{3}{4} x^{\frac{4}{3}} + \frac{3}{50} x^{\frac{5}{3}}. \quad (8.6)$$

Two slightly different initial data can represent this CAUCHY datum. The first initial guess reads:

$$p(t, x) := \frac{t}{2}, \quad q(t, x) := t + x, \quad (8.7a)$$

$$b_0 := \underline{1}, \quad b_1 := \underline{1}, \quad (8.7b)$$

$$b_2(t, x) := \frac{1}{10}, \quad b_j := \underline{0}, \quad \forall j \geq 3. \quad (8.7c)$$

The second one is

$$p(t, x) := \frac{t}{2}, \quad q(t, x) := t + x, \quad (8.8a)$$

$$b_0 := \underline{1}, \quad b_1 := \underline{1} - \frac{t}{10}, \quad (8.8b)$$

$$b_2(t, x) := \frac{1}{10} + \frac{t^2}{5}, \quad b_j := \underline{0}, \quad \forall j \geq 3. \quad (8.8c)$$

The convergence of both initializations has been studied symbolically and numerically using the code provided in Appendix D. The norms of differences between two successive iterations are reported in Figures 11 and 12. Moreover, Figure 13 indicates that both iterations converge to the same element of $\mathcal{O}[[z]]$.

To make an intermediate conclusion, we would like to mention that the Authors tested several other (and even gradually more complicated) configurations, and the proposed algorithm's practical convergence was invariably observed. This constatation gives us a good hope that Conjectures 7.1, 7.4 are true. More precisely, these numerical tests make quite plausible the following (informal) conjecture, which, thanks to PICARD theorem, would imply the Conjecture 7.1:

Conjecture 8.2. There exists a BANACH space E with current point $(p(t, x), q(t, x), \{b_k(t, x)\}_{k=0}^{\infty})$ and a closed subset $B \subseteq E$ satisfying the three following properties:

1. The elements of B satisfy for $t = 0$ the initial condition corresponding to the value $u(0, x)$.
2. $\mathbf{F}_>(\{B\}) \subseteq B$.
3. The restriction $\mathbf{F}|_B$ is a contraction.

8.4. The singular locus $4p^3 - 27q^2 = 0$'s dependence on the Cauchy datum

Recall that the LERAY principle states that singularities of a ramified **linear** CAUCHY problem are determined by the singularity locus of the CAUCHY data. In our genuinely nonlinear case, this principle seems not to be valid anymore.




Figure 11: The maximum norm $L_\infty([a, b])$ of the difference between two successive iterations $(I)^{th}$ and $(I - 1)^{th}$. The initial data is given in Equation (8.7).




Figure 12: The maximum norm $L_\infty([a, b])$ of the difference between two successive iterations $(I)^{\text{th}}$ and $(I - 1)^{\text{th}}$. The initial data is given in Equation (8.8).




Figure 13: Differences between corresponding components of the solution data at $I = 25^{\text{th}}$ iteration. Both iterative processes were initialized with data (8.7) and (8.8) respectively. The values of all numerical parameters are given in Table 1.

Above, we solved (approximately) three different ramified CAUCHY problems (7.5), (8.3) and (8.6). The singularities of the solution are determined by functions p and q . Let us compare them at the end of respective iterative processes:

$$\begin{aligned} p_1^{(I)} &\approx \frac{1}{2}t + 0.0250t^2 + 0.0068t^3 + \dots \\ p_2^{(I)} &\approx \frac{1}{2}t + (0.0250 + 0.0004x + 0.0025x^2)t^2 \\ &\quad + (0.0175 + 0.0070x - 0.0009x^2 + 0.0003x^3)t^3 + \dots \\ p_3^{(I)} &\approx \frac{1}{2}t + 0.0113t^2 + 0.0019t^3 + \dots \end{aligned}$$

Let us have a look at the corresponding q component approximations:

$$\begin{aligned} q_1^{(I)} &\approx t + x - 0.0333t^2 - 0.0050t^3 + \dots \\ q_2^{(I)} &\approx t + x + (-0.0333 + 0.0133x)t^2 \\ &\quad + (0.0082 - 0.0010x + 0.0006x^2)t^3 + \dots \\ q_3^{(I)} &\approx t + x - 0.0200t^2 - 0.0017t^3 + \dots \end{aligned}$$

The lower indices show the test case number (1, 2 or 3 corresponding to Sections 8.1, 8.2 and 8.3 respectively) and we show only four significant digits for the sake of notation compactness. Just from the visual inspection of expressions provided above, we can see that solution data p and q **depend** on the CAUCHY datum. Thus, the LERAY principle seems not to hold in our problem, and it is another sign of the genuinely nonlinear problem.

9. Further generalizations

In this Section, we would like to announce the general conjecture that we would like to formulate as a result of our investigations on this topic. Let us denote by (t, \mathbf{x}) a generic point of \mathbb{C}^{n+1} , where $\mathbf{x} = (x_1, x_2, \dots, x_n) \in \mathbb{C}^n$. We fix also a polydisc $D \subset \mathbb{C}^{n+1}$ centered around the origin $\mathbf{0} \in \mathbb{C}^{n+1}$ and we denote by \mathcal{O}_D the ring of holomorphic functions $A(t, \mathbf{x})$ on D .

Definition 9.1. Denote by \mathcal{S}_{m-1} the vector space of polynomial functions in several variables on \mathcal{O}_D of the following form:

$$F(t, \mathbf{x}, v_{1,\alpha_1}, v_{2,\alpha_2}, \dots, v_{n,\alpha_n}) \stackrel{\text{def}}{=} \sum_{a \in \hat{a}} A_a(t, \mathbf{x}) \prod_{\ell=1}^n \prod_{k=1}^{m-1} v_{\ell,k}^{a_{\ell,k}},$$

where $A_a \in \mathcal{O}_D$, $\forall a \in \hat{a}$ and the collection of maps \hat{a} is a finite set with elements of the type

$$a : n^{\sqsupset} \times (m-1)^{\sqsupset} \longrightarrow \mathbb{N},$$

which do not vanish identically, so that we always have

$$F(t, \mathbf{x}, 0, \dots, 0) \equiv \underline{0}.$$

In the Definition above, $v_{\ell,k}$ (respectively, $v_{\ell,k}^{a(\ell,k)}$) should correspond morally to $\partial_{x_\ell}^k$ (respectively, to the nonlinear operator $u \mapsto (\partial_{x_\ell}^k u)^{a(\ell,k)}$).

Let $u(t, \mathbf{x})$ be a germ of a holomorphic function at a point of the polydisc D . For any $F \in \mathcal{S}_{m-1}$ we shall write:

$$F(t, \mathbf{x}, \partial_{x_1}^{\alpha_1} u, \partial_{x_2}^{\alpha_2} u, \dots, \partial_{x_n}^{\alpha_n} u) \stackrel{\text{def}}{=} \sum_{a \in \hat{a}} A_a(t, \mathbf{x}) \prod_{\ell=1}^n \prod_{k=1}^{m-1} (\partial_{x_\ell}^k u)^{a(\ell,k)}.$$

Notice that in the previous expression, the involved partial derivatives of the germ u are of order at most $m - 1$. Also, as above, we shall use extensively the multi-index notation. If $\beta := (\beta_1, \beta_2, \dots, \beta_n) \in \mathbb{N}^n$, we shall denote $|\beta| := \beta_1 + \beta_2 + \dots + \beta_n$ and $\partial_{\mathbf{x}}^\beta := \partial_{x_1}^{\beta_1} \circ \partial_{x_2}^{\beta_2} \circ \dots \circ \partial_{x_n}^{\beta_n}$.

We also consider the following scalar quasi-linear differential operator of order m :

$$\begin{aligned} P(u) &\stackrel{\text{def}}{=} \partial_t^m u \\ &+ \sum_{j=0}^{m-1} \sum_{\substack{\beta \in \mathbb{N}^n \\ |\beta| = m-j}} F_{j,\beta}(t, \mathbf{x}, \partial_{x_1}^{\alpha_1} u, \partial_{x_2}^{\alpha_2} u, \dots, \partial_{x_n}^{\alpha_n} u) \partial_t^j \circ \partial_{\mathbf{x}}^\beta u \\ &+ G(t, \mathbf{x}, \partial_{x_1}^{\alpha_1} u, \partial_{x_2}^{\alpha_2} u, \dots, \partial_{x_n}^{\alpha_n} u), \end{aligned}$$

where $F_{j,\beta}$ and G both belong to the vector space \mathcal{S}_{m-1} . We are finally led to formulate the following

Conjecture 9.2. Consider $p \in \mathbb{N}^\times$ and $p + 1$ holomorphic functions $\{v_j(\mathbf{x})\}_{j=0}^p$ on $D \cap (\{0\} \times \mathbb{C}^n)$ such that both $v_0(\mathbf{0})$ and $v_1(\mathbf{0})$ are not zero. Let

$$u_0(\mathbf{x}) = \sum_{j=0}^p v_j(\mathbf{x}) x_1^{m-1 + \frac{j}{p+1}} \tag{9.1}$$

be a germ of a holomorphic function ramified around $x_1 = 0$. We make the following two assumptions:

- 1) The following polynomial equation in variable ξ admits m distinct roots:³⁹

$$\begin{aligned} \xi^m + \sum_{j=0}^{m-1} \xi^j F_{j,(m-j,0,\dots,0)}(0, \mathbf{0}, \partial_{x_1}^{\alpha_1} u_0(\mathbf{0}), \partial_{x_2}^{\alpha_2} u_0(\mathbf{0}), \dots, \partial_{x_n}^{\alpha_n} u_0(\mathbf{0})) \\ = 0. \end{aligned} \tag{9.2}$$

- 2) $\partial_{v_1, m-1} F_{0,(m,0,\dots,0)}(0, \mathbf{0}, \partial_{x_1}^{\alpha_1} u_0(\mathbf{0}), \partial_{x_2}^{\alpha_2} u_0(\mathbf{0}), \dots, \partial_{x_n}^{\alpha_n} u_0(\mathbf{0})) \neq 0$.

³⁹This is precisely the place where we see that the initial data enters explicitly into the definition of a suitable differential operator in the context of our study.

Then, the following statements are true. For each choice of a root to the algebraic Equation (9.2), there exist holomorphic functions $\{q_\ell(t, \mathbf{x})\}_{\ell=0}^{p-1}$ defined in a neighbourhood of the origin $\mathbf{0} \in \mathbb{C}^{n+1}$ such that

$$q_0(0, \mathbf{x}) = x_1, \quad q_\ell(0, \mathbf{x}) = \underline{0}, \quad \ell \in (p-1)^\square.$$

We denote by $z(t, \mathbf{x})$ a holomorphic germ satisfying the following algebraic equation outside the swallowtail singularity:

$$z^{p+1}(t, \mathbf{x}) - q_{p-1}(t, \mathbf{x})z^{p-1}(t, \mathbf{x}) - \cdots - q_0(t, \mathbf{x}) = \underline{0}.$$

There exist also the holomorphic functions $\{a_\ell(t, \mathbf{x})\}_{\ell=0}^p$ defined on a neighbourhood of the origin $\mathbf{0} \in \mathbb{C}^{n+1}$ such that the function

$$u(t, \mathbf{x}) = \partial_{q_0}^{-m+1}(a_p(t, \mathbf{x})z^p(t, \mathbf{x}) + \cdots + a_1(t, \mathbf{x})z(t, \mathbf{x}) + a_0(t, \mathbf{x}))$$

satisfies:

$$\begin{aligned} P(u)(t, \mathbf{x}) &\equiv 0, \\ u(0, \mathbf{x}) &\equiv u_0(\mathbf{x}), \end{aligned}$$

where $u_0(\mathbf{x})$ was defined in (9.1).

Let us now discuss the motivation behind this general Conjecture 9.2. The p -tuple (z, z^2, \dots, z^p) is the solution of a holonomic \mathcal{D} -module whose characteristic variety \mathbf{V} is included in the union of the zero section of $\mathbf{T}^*\mathbb{C}^p$ and of the co-normal to the swallow tail singularity associated to the algebraic equation:

$$z^{p+1} - q_{p-1}z^{p-1} - \cdots - q_1z - q_0 = \underline{0}.$$

The fact that this singularity is *stable* plays the crucial rôle in our constructions. Moreover, $\mathbf{T}_0^*\mathbb{C}^p \cap \mathbf{V}$ coincides with the line co-normal to $q_0 = 0$ and $q_0(0, \mathbf{x}) = x_1$. These important facts explain why we consider only the contributions of the monomials $\partial_t^j \circ \partial_{x_1}^{m-j}$ in Equation (9.2).

The assumption (1) means that all the characteristics associated with this geometry are simple. The proof of Conjecture 9.2 should allow to construct m solutions of this type, each corresponding to a choice of a root in Equation (9.2). It is quite clear that holomorphic functions $\{q_\ell(t, \mathbf{x})\}_{\ell=0}^{p-1}$ highly depend on the initial datum $u_0(\mathbf{x})$ and, even more precisely, on $\{v_j(\mathbf{x})\}_{j=0}^p$. The assumption (2) means that the PDE $P(u) = \underline{0}$ is genuinely nonlinear with respect to this geometry. The simplest example of such an operator $P(\cdot)$ from the Conjecture 9.2 statement is given by the following expression:

$$P(u) := \partial_t^m u + (\partial_{x_1}^{m-1} u)^n \partial_{x_1}^m u, \quad n \in \mathbb{N}_{\geq 1}.$$

To make things clearer, let us emphasize that, for instance, the two following quasi-linear operators do not satisfy the assumption (2) of Conjecture 9.2:

- $P_1(u) = \partial_t^m u + (\partial_{x_1}^{m-2} u)^n \partial_{x_1}^m u,$
- $P_2(u) = \partial_t^m u + (\partial_{x_1}^m u)^n \partial_{x_2}^m u.$

We summarize below the proposed strategy to attack this problem:

1. We seek a solution of $P(u) = \underline{0}$ under the form

$$u(t, \mathbf{x}) = \partial_{q_0}^{-m+1} \left(\sum_{j=0}^{+\infty} b_j(t, \mathbf{x}) z^j \right).$$

Since it does not seem to be possible to construct a solution directly, we proceed as in Section 7, *i.e.* we try to find an algorithm with an underlying map \mathbf{F} such that a fixed point of \mathbf{F} gives a solution. The fact that the swallow tail singularity should be characteristic for the PDE will lead to the eikonal equation. So, in some sense, we use an ansatz and the method of the geometric optics.⁴⁰ A heuristic (but crucial) underlying idea is that the HAMILTONIAN flow of the principal symbol of the linearized operator should propagate the singularities along the co-normal of the swallow tail.

2. Find suitable semi-norms allowing to construct a BANACH space so that one might apply the fixed point theorem to the mapping \mathbf{F} .

We hope to motivate and stimulate the research in this direction. Of course, this programme is extremely difficult. The best rational reasons to believe that it should work are the validity of Theorem 2.8 (the case of algebraic equations of the second degree) and the numerical convergence results for the algorithm of Section 7.

10. Conclusions and perspectives

Above, we presented the main results of the present manuscript. The main conclusions and perspectives of this study are outlined below.

10.1. Conclusions

In this article, we considered several ramified CAUCHY problem (CP)s for the first and second-order genuinely nonlinear PDEs in the full complex setting. The term CP should be understood in the context because we specify only one initial condition even for PDEs (3.1) with $m > 1$. We succeeded in understanding completely the case of the iBE (1.1) using the methods of CAUCHY–KOVALEVSKAYA and those of the contact geometry. Both theories give the same singularities for the iBE solutions to the IVP (4.2). Then, we switched to the second-order IVP (7.1) for the genuinely nonlinear PDE (1.2). However, this problem cannot be addressed with the same methods. This observation led us to consider some holonomic \mathcal{D} -modules and their geometry, which gave us the solution ansatz. The closed-form expression

⁴⁰This method is also known as the HAMILTON–JACOBI theory.

is not possible even within the proposed ansatz. Consequently, we devised an iterative method whose fixed point gives the desired solution. Unfortunately, we were not able to prove rigorously its convergence. Nevertheless, we performed the practical computations with the celebrated computer algebra software `Maple`[™] in order to check in practice the behaviour of the proposed algorithm. Indeed, our symbolic computations clearly indicate the convergence of the proposed method. The main part of the employed code is provided in Appendix D. On the basis of this empirical evidence, we were able to formulate several new conjectures which remain open for the moment, the main one being the Conjecture 9.2.

10.2. Perspectives

We already mentioned in Remark 5.5 the need to study the micro-local singularity of distributions z and z^2 . The Authors are currently working to establish a rigorous convergence proof for the iterative scheme proposed in Section 7. The underlying theoretical setting was briefly mentioned in Section 2. This result will allow us to establish the existence and uniqueness result for the considered IVP to Equation (1.2). For the moment, the appropriate BANACH spaces seem to have been identified (see Section 2). Of course, the application of these methods to other fully nonlinear and even higher order PDEs (such as the family of PDEs (3.1) with $m > 2$) is to be expected. Finally, we would like to see the connections between our theory and the more classical theory of shock waves for p -systems as presented in [49, Chapter 12]. We shall investigate the interplay and connections between both theories. As the first glance at possible connections, we invite the reader to consult Appendix C. We would like to explore other types of singularities as a separate research direction. According to the classification of catastrophes [3], the cusp and swallowtail belong to the class A_n for some $n \in \mathbb{N}$. We believe that it would be interesting to explore other types of singularities D_n and E_n and the solutions they generate in the current genuinely nonlinear PDE framework.

For nonlinear PDEs of order two or higher, the theory of shock waves is deep, intricate, and quite elaborated (see, for instance, [49]). However, the specific type of algebraic singularities that we consider in this work offers a promising avenue for a more simplified treatment. It is in this context that we expect that it should be possible to explain the appearance of shocks for the real solution by holomorphic ramified singularities, as we pointed out in Section 4.3.

Acknowledgements. E. LEICHTNAM would like to acknowledge the hospitality of LAMA UMR #5127 (University SAVOIE MONT BLANC), where this work took a physical place. Both authors would like to thank F. GOLSE (CMLS, École Polytechnique) and D. SERRE (UMPA, ENS de LYON) for very helpful discussions. The second Author (EL) thanks G. LEBEAU for fruitful discussions about this topic some time ago (during the mid-eighties). Finally, we would like to express our gratitude to the Referee for their insightful remarks, which significantly enhanced the quality of our manuscript.

Funding. This publication is based upon work supported by the Khalifa University

of Science and Technology under Award No. FSU-2023-014.

References

- [1] Arnold, V. I.: Singularities of Caustics and Wave Fronts, vol. 62 of *Mathematics and Its Applications*. Springer Netherlands, Dordrecht, 1990.
- [2] Arnold, V. I.: *Mathematical Methods of Classical Mechanics*, 2nd ed. Springer, New York, 1997.
- [3] Arnold, V. I., Gusein-Zade, S. M., Varchenko, A. N.: *Singularities of Differentiable Maps. Volume 1*. Birkhäuser, Boston, 1985.
- [4] Banner, M. L., Peregrine, D. H.: Wave Breaking in Deep Water. *Ann. Rev. Fluid Mech.* 25 (1993), 373–397.
- [5] Boldrighini, C., Frigio, S., Maponi, P.: Exploding solutions of the complex two-dimensional Burgers equations: Computer simulations. *J. Math. Phys.* 53, 8 (aug 2012), 083101.
- [6] Bony, J.-M., Schapira, P.: Propagation des singularités analytiques pour les solutions des équations aux dérivées partielles. *Annales de l'institut Fourier* 26, 1 (1976), 81–140.
- [7] Burgers, J. M.: A mathematical model illustrating the theory of turbulence. *Advances in Applied Mechanics* 1 (1948), 171–199.
- [8] Caffisch, R. E., Gargano, F., Sammartino, M., Sciacca, V.: Complex singularities and PDEs. *Riv. Mat. Univ. Parma* 6, 1 (2015), 69–133.
- [9] Dafermos, C. M.: *Hyperbolic Conservation Laws in Continuum Physics*, 3 ed., vol. 325 of *Grundlehren der mathematischen Wissenschaften*. Springer, Berlin, Heidelberg, 2010.
- [10] D'Agnolo, A.: Problème de Cauchy ramifié en théorie des faisceaux (d'après un travail avec P. Schapira). *Séminaire Équations aux dérivées partielles (Polytechnique)*, 7 (1991), 7.
- [11] D'Agnolo, A., Schapira, P.: An inverse image theorem for sheaves with applications to the Cauchy problem. *Duke Mathematical Journal* 64, 3 (dec 1991), 451–472.
- [12] Eggers, J., Fontelos, M. A.: A catalogue of singularities. *Arxiv 0711.0442* (2007), 1–24.
- [13] Fontelos, M. A., Velazquez, J. J. L.: Fractal-like singularities for an inviscid one-dimensional model of fluid jets. *Eur. J. Appl. Math.* 11, 1 (feb 2000), 29–60.
- [14] Forster, O.: *Lectures on Riemann Surfaces*, 1 ed., vol. 81 of *Graduate Texts in Mathematics*. Springer New York, New York, NY, 1981.
- [15] Gabber, O.: The Integrability of the Characteristic Variety. *American Journal of Mathematics* 103, 3 (jun 1981), 445–468.
- [16] Gal, C. G., Gal, S. G., Goldstein, J. A.: Wave and telegraph equations with real time variable and complex spatial variables. *Complex Var. Elliptic Eqns.* 57, 1 (jan 2012), 91–109.
- [17] Gal, C. G., Gal, S. G., Goldstein, J. A.: Burgers and Black-Merton-Scholes equations with real time variable and complex spatial variable. *Applicable Analysis* 92, 8 (aug 2013), 1766–1786.
- [18] Gelfand, I. M., Kapranov, M. M., Zelevinsky, A. V.: *Discriminants, resultants and multidimensional determinants*. Birkhäuser, Boston, 1994.
- [19] Gilbert, A. D., Pauls, W.: Complex manifolds for the Euler equations: a hierarchy of ODEs and the case of vanishing angle in two dimensions. *Fluid Dyn. Res.* 43, 2 (apr 2011), 025505.
- [20] Glimm, J., Lax, P. D.: *Decay of Solutions of Systems of Nonlinear Hyperbolic Conservation Laws*. American Mathematical Society, Providence, Rhode Island, 1970.
- [21] Hamada, Y.: The singularities of the solutions of the Cauchy problem. *Publications of the Research Institute for Mathematical Sciences* 5, 1 (1969), 21–40.
- [22] Hamada, Y., Leray, J., Wagschal, C.: Systemes d'équations aux dérivées partielles à caractéristiques multiples: problème de Cauchy ramifié, hyperbolicité partielle. *J. Math. Pures Appl.* IX, 55 (1976), 297–352.
- [23] Hou, T. Y.: Blow-up or no blow-up? A unified computational and analytic approach to 3D incompressible Euler and Navier–Stokes equations. *Acta Numerica* 18 (may 2009), 277–346.

- [24] Ishii, T.: On a representation of the solution of the Cauchy problem with singular initial data. *Proc. Japan Acad., Ser. A* 56 (1980), 59–61.
- [25] Kashiwara, M.: On the maximally overdetermined system of linear differential equations. I. *Publ. Res. Inst. Math. Sci., Kyoto Univ.* 10, 2 (1975), 563–579.
- [26] Kashiwara, M., Schapira, P.: Problème de Cauchy pour les systèmes microdifférentiels dans le domaine complexe. *Inventiones Mathematicae* 46, 1 (feb 1978), 17–38.
- [27] Kenyon, R., Okounkov, A.: Limit shapes and the complex Burgers equation. *Acta Mathematica* 199, 2 (2007), 263–302.
- [28] Kovalevskaya, S.: Zur Theorie der partiellen Differentialgleichungen. *J. Reine Angew. Math.* 80 (1875), 1–32.
- [29] Ladyzhenskaya, O. A.: The Mathematical Theory of Viscous Incompressible Flow. Martino Publishing, Eastford (CT), USA, 2014.
- [30] Larios, A., Petersen, M. R., Titi, E. S., Wingate, B.: A computational investigation of the finite-time blow-up of the 3D incompressible Euler equations based on the Voigt regularization, *Theoretical and Computational Fluid Dynamics* 32 (1), (2018), 23–34.
- [31] Lax, P. D.: Hyperbolic Systems of Conservation Laws and the Mathematical Theory of Shock Waves. SIAM, Philadelphia, Penn., 1973.
- [32] Leichtnam, E.: Front d’onde d’une sous-variété: applications aux équations aux dérivées partielles non linéaires. *Comm. Partial Diff. Eqns.* 10, 11 (jan 1985), 1229–1260.
- [33] Leichtnam, E.: Construction de solutions singulières pour des équations aux dérivées partielles non linéaires. *Ann. scient. Ec. Norm. Sup.* 20, 2 (1987), 137–170.
- [34] Leichtnam, E.: Le problème de Cauchy ramifié. *Ann. scient. Ec. Norm. Sup.* 23, 3 (1990), 369–443.
- [35] Leichtnam, E.: Le problème de Cauchy ramifié semi-linéaire d’ordre deux. *Ann. scient. Ec. Norm. Sup.* 24, 2 (1991), 189–214.
- [36] Leichtnam, E.: Le problème de Cauchy ramifié linéaire pour des données à singularités algébriques. *Bulletin de la S.M.F.* 121, 2 (1993), 1–128.
- [37] Li, D., Sinai, Y. G.: Complex singularities of the Burgers system and renormalization group method. *Current Developments in Mathematics 2006*, 1 (2006), 181–210.
- [38] Li, D., Sinai, Y. G.: Blow ups of complex solutions of the 3D Navier-Stokes system and renormalization group method. *J. Eur. Math. Soc.* 10 (2008), 267–313.
- [39] Li, D., Sinai, Y. G.: Singularities of complex-valued solutions of the two-dimensional Burgers system. *J. Math. Phys.* 51, 1 (jan 2010), 015205.
- [40] Luo, G., Hou, Th. Y.: Toward the Finite-Time Blowup of the 3D Axisymmetric Euler Equations: A Numerical Investigation, *Multiscale Modeling & Simulation* 12 (1), (2014), 1722–1776.
- [41] Lutz, D. A., Miyake, M., Schäfke, R.: On the Borel summability of divergent solutions of the heat equation. *Nagoya Mathematical Journal* 154 (jan 1999), 1–29.
- [42] Malgrange, B.: L’involutivité des caractéristiques des systèmes différentiels et microdifférentiels. *Séminaire Bourbaki* 20 (1977), 277–289.
- [43] Matytsin, A.: On the large-N limit of the Itzykson-Zuber integral. *Nuclear Physics B* 411, 2-3 (jan 1994), 805–820.
- [44] Narasimhan, R.: Introduction to the Theory of Analytic Spaces, vol. 25 of *Lecture Notes in Mathematics*. Springer Berlin Heidelberg, Berlin, Heidelberg, 1966.
- [45] Poláčik, P., Šverák, V.: Zeros of complex caloric functions and singularities of complex viscous Burgers equation. *J. Reine Angew. Math.* 2008, 616 (jan 2008), 205–215.
- [46] Pomeau, Y., Sciamarella, D.: An unfinished tale of nonlinear PDEs: Do solutions of 3D incompressible Euler equations blow-up in finite time?, *Physica D: Nonlinear Phenomena* 205 (1-4), (2005), 215–221.
- [47] Popivanov, P. R.: Nonlinear PDE. Singularities, Propagation, Applications. In *Nonlinear Hyperbolic Equations, Spectral Theory, and Wavelet Transformations*. Birkhäuser Basel, Basel, 2003, pp. 1–94.
- [48] Rosenkranz, M., Wang, D. (Eds): Gröbner Bases in Symbolic Analysis. Walter De Gruyter, Berlin and New York, dec 2007.
- [49] Serre, D.: Systems of Conservation Laws 2: Geometric Structures, Oscillations, and Initial-Boundary Value Problems. Cambridge University Press, Cambridge, 2000.

-
- [50] Singh, J., Kumar, S. D.: On the Involutivity of the Characteristic Variety. *Communications in Algebra* 42, 8 (aug 2014), 3607–3618.
 - [51] Van der Put, M.: Some properties of the ring of germs of C^∞ -functions. *Compositio Mathematica* 34, 1 (1977), 99–108.
 - [52] Viswanath, D., Sahutoglu, S.: Complex Singularities and the Lorenz Attractor. *SIAM Review* 52, 2 (jan 2010), 294–314.
 - [53] Wagschal, C.: Sur le problème de Cauchy ramifié. *J. Math. Pures Appl.* 53, 9 (1974), 147–163.
 - [54] Weideman, J. A. C.: Computing the Dynamics of Complex Singularities of Nonlinear PDEs. *SIAM Journal on Applied Dynamical Systems* 2, 2 (jan 2003), 171–186.
 - [55] Woodcock, A. E. R., Poston, T.: A Geometrical Study of the Elementary Catastrophes, vol. 373 of *Lecture Notes in Mathematics*. Springer Berlin Heidelberg, Berlin, Heidelberg, 1974.

A. Nomenclature

$f : A \rightarrow B$ A map from the domain $\text{dom } f = A$ to the co-domain (or target) $\text{codom } f = B$.

$f_{>}$ Image of a map. Consider a map $f : A \rightarrow B$. Then, the image $f_{>} : \wp(A) \rightarrow \wp(B)$ is defined in the usual way, where $\wp(\cdot)$ denotes the power set.

$f_{<}$ Pre-image of a map. Consider a map $f : A \rightarrow B$. Then, the pre-image $f_{<} : \wp(B) \rightarrow \wp(A)$ is defined in the usual way, where $\wp(\cdot)$ denotes the power set.

○ Composition operation for functions of differential operators.

$f^m \circ (-)$ The m^{th} iterate of a mapping $f : A \rightarrow A$ for some $m \in \mathbb{N}^{\times}$, *i.e.*

$$f^m \circ (-) \stackrel{\text{def}}{=} \underbrace{(f \circ f \circ \dots \circ f)}_{m \text{ times}}(-)$$

df The differential of a smooth function f .

$j^1(u)$ 1-jet of a smooth function u .

\mathbb{N} The set of natural numbers starting from 0.

\mathbb{N}^{\times} The set of strictly positive natural numbers.

\mathbb{R} The set of real numbers.

\mathbb{R}^{\times} The multiplicative group of real numbers.

$\mathbb{R}_{\geq 0}$ The set of non-negative real numbers.

\mathbb{C} The set of complex numbers.

\mathbb{C}^{\times} The multiplicative group of complex numbers.

$|\cdot|$ The absolute value of a real or complex number.

$\|\cdot\|$ The norm function on a BANACH space.

\mathcal{O}_D The ring of holomorphic functions defined on the polydisc D .

$\mathcal{O}[z]$ The ring of polynomials in the formal variable z with coefficients being holomorphic germs at the origin.

$\mathcal{O}[[z]]$ The ring of formal power series in the formal variable z with coefficients being holomorphic germs at the origin.

n^{\sqsupset} Finite set $\{1, 2, \dots, n\}$, $\forall n \in \mathbb{N}_{\geq 1}$.

n^{\sqsubset} Finite set $\{0, 2, \dots, n-1\}$, $\forall n \in \mathbb{N}_{\geq 1}$.

-
- r A function taking the constant value $r \in \mathbb{R}$. The domain and co-domain of this function should be clear from the context (usually, from the left-hand side of the equation, where it appears).
- \mathbf{r} A function taking the constant vector value $(r, r, \dots, r) \in \mathbb{R}^m$ with $r \in \mathbb{R}$ and $m \in \mathbb{N}_{\geq 2}$. The domain and co-domain of this function should be clear from the context (usually, from the left-hand side of the equation, where it appears).
- $(-)^{\top}$ The transposition operator acting on a vector or on a linear operator. Strictly speaking, the result of this operation belongs to the dual vector space.
- \mathbf{x} Element of the vector space $\mathbf{x} \stackrel{\text{def}}{:=} (x_1, x_2, \dots, x_n) \in \mathbb{C}^n$.
- $\partial_{x_j}^m$ Partial derivative operator of the order m with respect to the independent variable x_j . Sometimes, we also use the index notation $(\cdot)_{x_i}$ to denote the first-order partial derivative.
- E A BANACH space.
- F Fixed point mapping.
- \tilde{F} Finitary truncation of the fixed point mapping.
- $\psi^{(I)}$ Function ψ on the $(I)^{\text{th}}$ iteration.
- $:=$ Assignment of the right-hand side to the left-hand side.
- $\stackrel{\text{def}}{:=}$ The left-hand side is defined by the right-hand side.
- $\stackrel{\text{def}}{=}$ The right-hand side is defined by the left-hand side.
- $:\Rightarrow$ By definition.
- \equiv Equal identically.
- \cong Isomorphic.
- \mathbb{k}^n The standard n -dimensional vector space with the base field \mathbb{k} .
- \mathcal{U}_z Some open neighbourhood of a point $z \in \mathbb{C}^m$.
- $\bar{\xi}$ Bold version of the Greek letter ξ .
- $Z(G)$ The center of the group G .
- U_n The multiplicative group of the roots of unity of degree n .
- S_n The symmetric group with n symbols.

$\mathbf{GL}_n(\mathbb{k})$ The general linear group of the square $n \times n$ matrices with entries belonging to the field \mathbb{k} .

$\mathbf{SL}_n(\mathbb{k})$ The special linear group of the square $n \times n$ matrices with entries belonging to the field \mathbb{k} .

$\mathbf{PSL}_n(\mathbb{k})$ The projective special linear group defined as $\mathbf{SL}_n(\mathbb{k}) / Z(\mathbf{SL}_n(\mathbb{k}))$. For $n = 2$ and $\mathbb{k} = \mathbb{Z}$ it is called the modular group.

\mathbf{Br}_n The braid group with n strands.

\mathbf{M} Holomorphic, smooth or topological manifold, depending on the context.

$\pi_1(\mathbf{M})$ The fundamental (or the first homotopy) group of a topological manifold \mathbf{M} .

$\mathbf{T}_x \mathbf{M}$ Tangent space to a smooth manifold \mathbf{M} at point $x \in \mathbf{M}$.

ω The standard symplectic 2-form on $\mathbf{T}_x \mathbf{M}$.

$\{-, -\}$ The standard POISSON bracket on $C^\infty(\mathbf{M})$.

$\mathbf{T}\mathbf{M}$ Tangent bundle of a smooth manifold \mathbf{M} .

$\mathbf{T}_x^* \mathbf{M}$ Co-tangent space to a smooth manifold \mathbf{M} at point $x \in \mathbf{M}$.

$\mathbf{T}^* \mathbf{M}$ Co-tangent bundle of a smooth manifold \mathbf{M} .

$\mathcal{N}(\mathbf{M})$ Co-normal to a smooth sub-manifold \mathbf{M} .

$\text{sing supp}(\cdot)$ The singular support of a distribution.

$\text{WF}(\cdot)$ The wave front set of a distribution.

\mathbf{X} The complex manifold.

$\{X_\lambda\}_{\lambda \in \Lambda}$ The WHITNEY stratification of the complex manifold \mathbf{X} .

\mathcal{D}_X The sheaf of differential operators of *finite order* on the complex manifold \mathbf{X} .

\mathfrak{M} The coherent \mathcal{D}_X -module.

\mathcal{O}_X The sheaf of holomorphic functions on \mathbf{X} , which is a left coherent \mathcal{D}_X -module.

Ext The right derived functor of the hom-functor.

$\mathbf{Gal}(-)$ GALOIS group of a field extension.

B. Details of computations

In this Appendix, we explain more precisely how we construct the iterations from Section 7.1. Namely, we specify how the expression $(\widehat{\mathfrak{X}})$ is constructed from (\mathfrak{X}) .

Recall, that the solution ansatz is given by

$$u(t, x) = \sum_{k=1}^{\infty} \frac{1}{k} \left(-p(t, x) b_{k-1}(t, x) + 3b_{k-3}(t, x) \right) z^k.$$

One checks by direct computation that

$$\begin{aligned} u_{tt} = & \sum_{k=0}^{+\infty} \left[\left(2 - \frac{2}{k} \right) b_{k-1,t} p_t + \left(1 - \frac{1}{k} \right) b_{k-1} p_{tt} + b_k q_{tt} + \underline{2b_{k,t} q_t} + \right. \\ & \left. \frac{1}{k} (p b_{k-1,tt} + 3b_{k-3,tt}) \right] z^k + \\ & \frac{1}{3z^2 - p} \sum_{k=0}^{+\infty} \left[(k-1) b_{k-1} p_t^2 + 2k b_k p_t q_t + (k+1) b_{k+1} q_t^2 \right] z^k, \end{aligned} \tag{B.1}$$

Moreover, one checks that the coefficient in front of $\frac{1}{3z^2 - p}$ in $(\mathfrak{X}) \Rightarrow u_{tt} - u_x u_{xx}$ is equal to

$$\begin{aligned} & \sum_{k=0}^{+\infty} \left[(k-1) p_t^2 b_{k-1} + \underline{2k b_k p_t q_t} + \underline{(k+1) b_{k+1} q_t^2} \right. \\ & \left. - \sum_{j=0}^k \left(\left(1 - \frac{1}{j} \right) p_x b_{j-1} + q_x b_j + \frac{1}{j} (-p b_{j-1,x} + 3b_{j-3,x}) \right) \right. \\ & \left. \left((k-j-1) p_x^2 b_{k-j-1} + 2(k-j) p_x q_x b_{k-j} \right. \right. \\ & \left. \left. + (k-j+1) q_x^2 b_{k-j+1} \right) \right] z^k. \end{aligned} \tag{B.2}$$

Now, we construct $(\widehat{\mathfrak{X}})$ from $(\mathfrak{X}) \equiv u_{tt} - u_x u_{xx}$ in the following way. First, we replace in (B.1) the term $2b_{k,t} q_t$ by $2\hat{b}_{k,t} \hat{q}_t$. Then, we replace in (B.2) the term $2k b_k p_t q_t$ by $2k b_k \hat{p}_t \hat{q}_t$ and $(k+1) b_{k+1} q_t^2$ by $(k+1) b_{k+1} \hat{q}_t \hat{q}_t$. Then, the eikonal Equation (7.10) is obtained by writing that $(\widehat{\mathfrak{X}}) \in \mathcal{O}[[z]]$, namely, the coefficient (B.2) of $\frac{1}{3z^2 - p}$ in $(\widehat{\mathfrak{X}})$ is divisible by $3z^2 - p$. Finally, Equation (7.12) is obtained by fulfilling the requirement that $(\widehat{\mathfrak{X}}) \equiv 0$.

Remark B.1. In Equation (B.2) we could have replaced the term $2k b_k p_t q_t$ (respectively, $(k+1) b_{k+1} q_t^2$) by $2k b_k \hat{p}_t \hat{q}_t$ (respectively $(k+1) b_{k+1} \hat{q}_t^2$) and

thus get a nonlinear eikonal equation. What we did is roughly a simple linearisation of this, which is more convenient for implementing in practice. In other words, no matter the eikonal equation's form, one has to perform the infinity of fixed point iterations. We took special care to make each such iterative step as easy as possible.

C. A simple explicit example of shock wave formation in the iBE

In this Appendix, we would like to provide one simple explicit example and discuss some relations between our theory of ramified solutions and the classical theory of shock waves. This Appendix should be perceived only as an appetizer. The deeper connections will be revealed in the subsequent works.

Let us consider the following IVP for the iBE:

$$u_t - u u_x = \underline{0}, \quad (\text{C.1})$$

$$u(0, x) = x^{\frac{1}{2}}, \quad (\text{C.2})$$

where we took the square root singularity to simplify algebraic computations which follow. In the real case, we would consider the IVP on a half-line $(t, x) \in \mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0}$, but there are no such restrictions in the complex case. So, in agreement with our line of thinking, we assume that $(t, x) \in \mathbb{C}^2$. Using the methods of contact geometry described in Section 4.3, we introduce the auxiliary variable

$$y \stackrel{\text{def}}{:=} x - x^{\frac{1}{2}} t. \quad (\text{C.3})$$

The solution u verifies the following second-order algebraic equation:

$$u^2 - t u - y = \underline{0}. \quad (\text{C.4})$$

The last algebraic Equation (C.4) can be also rewritten in an equivalent form:

$$\left(u - \frac{t}{2}\right)^2 - \left(y + \frac{t^2}{4}\right) = \underline{0}.$$

Now it is obvious that the discriminant of Equation (C.4) is

$$\delta \stackrel{\text{def}}{:=} y + \frac{t^2}{4}.$$

Our theory states that the complex solution u will be ramified around the locus $\delta = \underline{0}$. Taking into account (C.3), two solutions to Equation (C.4) read

$$r_{\pm} \stackrel{\text{def}}{:=} \frac{t}{2} \pm \sqrt{\left(\frac{t}{2} - \sqrt{x}\right)^2}.$$

We take two points $x_0 \neq x_1$ and we consider two characteristics (C.3) passing through these points at $t = 0$. If $x_{0,1}$ are real, we may assume that $0 < x_0 < x_1$

to fix the ideas. Along the characteristic $y_j \stackrel{\text{def}}{=} x_j - x_j^{\frac{1}{2}} t$ the solution u takes the value $x_j^{\frac{1}{2}}$, $j \in 2^{\square}$. A shock wave forms when two characteristics cross each other since in the crossing point, the solution value is obviously contradictory:

$$x_0 \neq x_1 \quad \implies \quad x_0^{\frac{1}{2}} \neq x_1^{\frac{1}{2}}.$$

The crossing will take place when

$$x_0 - x_0^{\frac{1}{2}} t = x_1 - x_1^{\frac{1}{2}} t.$$

It can be easily checked that the characteristic $y = x_0 - x_0^{\frac{1}{2}} t$ meets the singular locus $\delta = 0$ in the real plane at time instance $t_0 \stackrel{\text{def}}{=} 2x_0^{\frac{1}{2}}$. Indeed,

$$y_0 = x_0 - x_0^{\frac{1}{2}} t_0 \equiv x_0 - 2x_0 = -x_0.$$

Henceforth,

$$\delta(t_0, x_0) \implies y_0 + \frac{t_0^2}{4} = -x_0 + \frac{(2x_0^{\frac{1}{2}})^2}{4} \equiv 0$$

and we deduce that $(t_0, x_0) \in \delta^<(\{0\})$. Moreover, it is not difficult to compute exactly the crossing time t_* :

$$t_* = x_0^{\frac{1}{2}} + x_1^{\frac{1}{2}} > 2x_0^{\frac{1}{2}} \implies t_0.$$

We would like to explain what happens precisely at time $t = t_0$. Until the crossing time $t = t_*$, the solution value along the characteristic y_0 is $x_0^{\frac{1}{2}}$. The point is that at time $t = t_0$, we change the solution branch. Indeed, for $0 < t < t_0$:

$$r_+ \implies \frac{t}{2} + x_0^{\frac{1}{2}} - \frac{t}{2} \equiv x_0^{\frac{1}{2}},$$

while for $t_0 < t < t_*$ we have

$$r_- \implies \frac{t}{2} - \left(\frac{t}{2} - x_0^{\frac{1}{2}} \right) \equiv x_0^{\frac{1}{2}}.$$

In other words, at the discriminant contact point $t = t_0$, we have no other choice except to jump from one solution branch to another in order to keep the constant solution value along the given characteristic. The change in the solution branch precedes and explains the apparition of the shock. This situation is illustrated in Figure 14.




Figure 14: A schematic graphical illustration of the shock wave formation in the iBE for an IVP (C.1), (C.2) with the square root algebraic singularity.

D. Maple code

In this Appendix we provide the Maple™ code which implements one iteration of the mapping \tilde{F} used in Section 8:

```

restart :
with(LinearAlgebra) :
Digits := 30;
DEG := 25;

[...]

OneIter := proc(v::Array)
  description "One iteration of the fixed point algorithm";
  local N::integer ,K::integer ,l::integer ,p::symbol ,q::
    symbol;
  local px::symbol ,pt::symbol ,qx::symbol ,qt::symbol;
  local qtt::symbol ,qxx::symbol ,b::Array ,A::Array ,B::Array;
  local BB::Array ,C::Array ,M::Matrix ,Minv::Matrix ,B0::
    symbol;
  local B1::symbol ,ptt::symbol ,B0rhs::symbol ,B1rhs::symbol;

```

```

local k::integer , j::integer , P::symbol , Q::symbol , Pt::
    symbol;
local Qt::symbol , RHS::symbol , rhs1::symbol , rhs2::symbol;
local M11::symbol , M12::symbol , M21::symbol , M22::symbol;
p := convert(v[1], float);
q := convert(v[2], float);
pt := convert(diff(p, t), float);
px := convert(diff(p, x), float);
qt := convert(diff(q, t), float);
qx := convert(diff(q, x), float);
ptt := convert(diff(p, t$2), float);
qtt := convert(diff(q, t$2), float);
qxx := convert(diff(q, x$2), float);
N := ArrayNumElems(v) - 2;
b := Array(-3 .. N - 1);
b[-3 .. -1] := Array([0., 0., 0.]);
b[0 .. N - 1] := convert(v[3 .. N + 2], float);
K := ceil((N - 2)/2);
B := Array(0 .. K);
for k from 0 to K do
    B[k] := convert(b[0]*diff(q, x)*((k - 1.0)*px^2*b[k -
        1] + 2.0*k*px*qx*b[k] + (k + 1)*qx^2*b[k + 1]),
        float);
    for j from 1 to k do
        B[k] := B[k] + convert((((1.0 - 1.0/j)*px*b[j - 1] +
            qx*b[j] + (-p*diff(b[j - 1], x) + 3.0*diff(b[j -
                3], x))/j)*((k - j - 1.0)*px^2*b[k - j - 1] +
                2.0*(k - j)*px*qx*b[k - j] + (k - j + 1.0)*qx^2*b
                    [k - j + 1]), float);
    end do;
    B[k] := convert(B[k] - (k - 1.0)*pt^2*b[k - 1], float);
end do;
rhs1 := convert(add((p/3.0)^k*B[2*k], k = 0 .. floor(K/2)
    ), float);
rhs2 := convert(add((p/3.0)^k*B[2*k + 1], k = 0 .. floor
    ((K - 1)/2)), float);
M11 := convert(4.0*qt*add((p/3.0)^k*k*b[2*k], k = 0 ..
    floor((N - 1)/2)), float);
M12 := convert(qt*add((p/3.0)^k*(2.0*k + 1.0)*b[2*k + 1],
    k = 0 .. floor((N - 2)/2)), float);
M21 := convert(2.0*qt*add((p/3.0)^k*(2.0*k + 1.0)*b[2*k +
    1], k = 0 .. floor((N - 2)/2)), float);
M22 := convert(2.0*qt*add((p/3.0)^k*(k + 1.0)*b[2*k + 2],
    k = 0 .. floor((N - 3)/2)), float);
M := convert(Matrix([[M11, M12], [M21, M22]]), float);

```

```

Minv := MatrixInverse(M);
Pt := mtaylor(rhs1*Minv[1, 1] + rhs2*Minv[1, 2], [t, x],
  DEG);
Qt := mtaylor(rhs1*Minv[2, 1] + rhs2*Minv[2, 2], [t, x],
  DEG);
P := dsolve({P(0.) = 0., diff(P(t), t) = Pt});
Q := dsolve({Q(0.) = 1.0*x, diff(Q(t), t) = Qt});
A := Array(0 .. N - 2);
for k from 0 to N - 2 do
  A[k] := -b[0]*diff(q, x)*((k - 1.0)*px^2*b[k - 1] +
    2.0*k*px*qx*b[k] + (k + 1.0)*qx^2*b[k + 1]);
  for j from 1 to k do
    A[k] := A[k] - ((1.0 - 1.0/j)*px*b[j - 1] + qx*b[j] +
      (-p*diff(b[j - 1], x) + 3.0*diff(b[j - 3], x))/j
      )*((k - j - 1.0)*px^2*b[k - j - 1] + 2.0*(k - j)*
      px*qx*b[k - j] + (k - j + 1.0)*qx^2*b[k - j + 1])
    ;
  end do;
  A[k] := convert(A[k] + (k - 1.0)*pt^2*b[k - 1] + 2.0*k*
    Pt*qt*b[k] + (k + 1.0)*Qt*qt*b[k + 1], float);
end do;
C := Array(0 .. N - 4);
for k from 0 to N - 4 do
  C[k] := convert(add((p/3.0)^1*A[k + 2 + 2*1], 1 = 0 ..
    floor((N - k - 4)/2)), float);
end do;
B0rhs := mtaylor(1/(2.0*qt)*(-b[0]*qtt - C[0] + b[0]*qx*(
  b[0]*qxx + 2.0*qx*diff(b[0], x))), [t, x], DEG);
B0 := dsolve({B0(0.0) = 1.0, diff(B0(t), t) = B0rhs});
B1rhs := mtaylor(1/(2.0*qt)*(-b[1]*qtt + p*diff(b[0], t $
  2) - C[1] + b[0]*qx*(b[1]*qxx + 2.0*qx*diff(b[1], x)
  - p*diff(b[0], x $ 2)) + (b[1]*qx - p*diff(b[0], x))
  *(b[0]*qxx + 2*qx*diff(b[0], x))), [t, x], DEG);
B1 := dsolve({B1(0.0) = 1.0, diff(B1(t), t) = B1rhs});
BB := Array(2 .. N - 4);
for k from 2 to N - 4 do
  RHS := -(2.0 - 2.0/k)*pt*diff(b[k - 1], t) - (1.0 -
    1.0/k)*ptt*b[k - 1] - qtt*b[k] - 1.0/k*(3.0*diff(b[
    k - 3], t $ 2) - p*diff(b[k - 1], t $ 2)) - C[k] +
    qx*b[0]*((2.0 - 2.0/k)*px*diff(b[k - 1], x) + (1.0
    - 1.0/k)*ptt*b[k - 1] + qxx*b[k] + 2.0*qx*diff(b[k
    ], x) + 1.0/k*(-p*diff(b[k - 1], x $ 2) + 3.0*diff(
    b[k - 3], x $ 2)));
  for j from 1 to k-1 do

```

```

RHS := RHS + ((1.0 - 1.0/j)*px*b[j - 1] + qx*b[j] +
  1.0/j*(-p*diff(b[j - 1], x) + 3.0*diff(b[j - 3],
  x)))*((2.0 - 2.0/(k - j))*px*diff(b[k - j - 1], x
  ) + (1.0 - 1.0/(k - j))*ptt*b[k - j - 1] + qxx*b[
  k - j] + 2.0*qx*diff(b[k - j], x) + 1.0/(k - j)
  *(-p*diff(b[k - j - 1], x $ 2) + 3.0*diff(b[k - j
  - 3], x $ 2)));
end do;
RHS := RHS + ((1.0 - 1.0/k)*px*b[k - 1] + qx*b[k] +
  1.0/k*(-p*diff(b[k - 1], x) + 3.0*diff(b[k - 3], x)
  ))*(qxx*b[0] + 2.0*qx*diff(b[0], x));
RHS := mtaylor(RHS/(2.0*qt), [t, x], DEG);
BB[k] := dsolve({beta(0.0) = 0.0, diff(beta(t), t) =
  RHS});
end do;
return convert(Array([convert(series(rhs(P), t = 0, DEG),
  polynomial), convert(series(rhs(Q), t = 0, DEG),
  polynomial), convert(series(rhs(B0), t = 0, DEG),
  polynomial), convert(series(rhs(B1), t = 0, DEG),
  polynomial), seq(convert(series(rhs(BB[j]), t = 0, DEG),
  polynomial), j = 2 .. N - 4)]), float);
end proc;

```

The code provided above was used to study the convergence of initial data (8.1), (8.2) and all the others from Section 8. A few modifications are needed to run the IVP (8.4). The complete programs can be obtained under a simple request by email.

E. An example of a genuinely nonlinear problem

Sometimes, an example is worth one thousand words. In our terminology, the following CAUCHY problem is nonlinear (quasi-linear) but not genuinely nonlinear:

$$u_{tt} - uu_{xx} = \underline{0}, \quad u(0, x) = c_1 x + c_2 x^{1+\frac{1}{3}},$$

for some real non-zero constants $c_{1,2} \in \mathbb{R}^\times$. In contrast, the following PDE is genuinely nonlinear:

$$u_{tt} - u_x u_{xx} = \underline{0}, \quad u(0, x) = c_1 x + c_2 x^{1+\frac{1}{3}}.$$

In reality, this notion depends on the operator *and* on the initial condition as well. That is why we speak above about the CAUCHY problem instead of just a PDE. The notion of a genuinely nonlinear problem was made more precise in Section 9.

Denys Dutykh

Mathematics Department, Khalifa University of Science and Technology, PO Box 127788, Abu Dhabi, United Arab Emirates

Denys.Dutykh@ku.ac.ae

<http://www.denys-dutykh.com/>

Éric Leichtnam

CNRS, Institut de Mathématiques de Jussieu–Paris Rive Gauche UMR 7586, UP7D – Bâtiment Sophie Germain, 56-58 avenue de France — BC 7012, 75205 Paris Cedex 13

Eric.Leichtnam@imj-prg.fr

© The copyright of this article is retained by the Author(s).

Open Access. This article is published in open access form and licensed under the terms of the Creative Commons Attribution–NonCommercial–ShareAlike 4.0 International (CC BY-NC-SA 4.0) (<http://creativecommons.org/licenses/by-nc-sa/4.0/>).